www.bjcancer.com

Network-based analysis of calcium-binding protein genes identifies Grp94 as a target in human oral carcinogenesis

H Nomura¹, K Uzawa^{*,1,2}, Y Yamano¹, K Fushimi¹, T Ishigami¹, Y Kato¹, K Saito¹, D Nakashima¹, M Higo¹, Y Kouzu¹, K Ono², K Ogawara¹, M Shiiba¹, H Bukawa², H Yokoe² and H Tanzawa^{1,2,3}

¹Department of Clinical Molecular Biology, Graduate School of Medicine, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba 260-8670, Japan; ²Division of Dentistry and Oral-Maxillofacial Surgery, Chiba University Hospital, 1-8-1 Inohana, Chuo-ku, Chiba 260-8670, Japan; ³Center of Excellence (COE) Program in the 21st Century, Graduate School of Medicine, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba 260-8670, Japan;

To characterise Ca^{2+} -binding protein gene expression changes in oral squamous cell carcinomas (OSCCs), we compared the gene expression profiles in OSCC-derived cell lines with normal oral tissues. One hundred Ca^{2+} -binding protein genes differentially expressed in OSCCs were identified, and genetic pathways associated with expression changes were generated. Among genes mapped to the network with the highest significance, glucose-regulated protein 94 kDa (Grp94) was evaluated further for mRNA and protein expression in the OSCC cell lines, primary OSCCs, and oral premalignant lesions (OPLs). A significant (P < 0.001) overexpression of Grp94 protein was observed in all cell lines compared to normal oral epithelium. Immunohistochemical analysis showed highly expressed Grp94 in primary OSCCs and OPLs, whereas most of the corresponding normal tissues had no protein immunoreaction. Real-time quantitative reverse transcriptase-PCR data agreed with the protein expression status. Moreover, overexpression of Grp94 in primary tumours was significantly (P < 0.001) correlated with poor disease-free survival. The results suggested that Grp94 may have potential clinical application as a novel diagnosis and prognostic biomarker for human OSCCs. *British Journal of Cancer* (2007) **97**, 792–801. doi:10.1038/sj.bjc.6603948 www.bjcancer.com

© 2007 Cancer Research UK

Keywords: oral squamous cell carcinoma; Ca²⁺-binding protein; pathway analysis; glucose-regulated protein 94 kDa; poor prognosis

Cytosolic Ca^{2+} , a basic parameter of intracellular homeostasis, has a wide variety of important cellular functions, that is, cell growth and proliferation, processing and folding of endoplasmic reticulum (ER)-translated proteins, and Ca^{2+} -mediated signalling in response to extracellular stimuli (Berridge *et al*, 2000; Kunzelmann, 2005). Under physiologic conditions, many aspects of these processes are mediated by members of the Ca^{2+} -binding protein (Nelson and Chazin, 1998). Since considerable evidence shows that intracellular Ca^{2+} deregulation is crucial for tumour growth and development (Durham and Walton, 1982; Huang *et al*, 2005; Ding *et al*, 2006; Thebault *et al*, 2006), understanding the genetic mechanisms of the Ca^{2+} -binding protein genes involved in carcinogenesis is imperative for the development of new therapeutic strategies.

 Ca^{2+} -binding protein is characterised by the presence of the putative Ca^{2+} -binding domain; EF-hand structure is an essential mediator of intracellular Ca^{2+} control elements, such as the plasma membrane Ca^{2+} pumps or ER (Koch *et al*, 1986; Nelson and Chazin, 1998; Haeseleer *et al*, 2000). A number of Ca^{2+} -binding proteins are thought to be implicated in establishing the malignant and metastatic phenotypes of various tumours (Chen *et al*, 1997; Takenaga *et al*, 1997; Van Ginkel *et al*, 1998), and recent studies have reported altered expression of several Ca^{2+} -binding protein genes in a wide range of human malignancies (Bustin *et al*, 2001; Pietas *et al*, 2002; Lakshmikuttyamma *et al*,

2004; Imazawa *et al*, 2005). Regarding oral squamous cell carcinomas (OSCCs), aberrant expression of S100 Ca²⁺-binding protein families seen in tumour sites has been correlated with cancer cell invasion or metastasis (Moriyama-Kita *et al*, 2004; Tsai *et al*, 2005). We recently identified certain Ca²⁺-binding protein genes differentially expressed in OSCCs compared with normal oral squamous epithelium, including *PMCA1* (Saito *et al*, 2006), *ITPKA* (Kato *et al*, 2006), *ATP2A2* (Endo *et al*, 2004), *CALR* (Shimada *et al*, 2005), and *E-cadherin* (Saito *et al*, 1998).

We recently developed the strategy of using a knowledge-based network approach to search for candidate genes relevant to the treatment of oral cancers (Kasamatsu *et al*, 2005; Higo *et al*, 2006; Ishigami *et al*, 2007). Thus, this study aimed to explore further the global changes in Ca^{2+} -binding protein genes associated with OSCCs by applying network development tools to expand on microarray analysis methods. In addition, a candidate gene for therapeutic targeting and follow-up studies was evaluated further for the expression status of the mRNA and protein in a large series of OSCCs and oral premalignant lesions (OPLs).

MATERIALS AND METHODS

Tissue specimens

Eighty pairs of primary OSCC samples and corresponding normal oral epithelial tissues or 20 OPLs (diagnosed as oral leukoplakias) were obtained at the time of surgery performed at Chiba University Hospital between 1998 and 2005. All patients provided informed

^{*}Correspondence: Dr K Uzawa; E-mail: uzawak@faculty.chiba-u.jp Revised 29 June 2007; accepted 31 July 2007; published online 28 August 2007

consent according to the protocol that was reviewed and approved by the Institutional Review Board of Chiba University before any procedures were performed. Postoperative follow-up data were collected until April 2006 or until the day of the patient's death, metastasis, or local recurrence. The median follow-up time was 2.1 years (range: 3 months to 8 years).

The resected tissues were divided into two parts: one was frozen immediately after removal of the surrounding normal tissue and stored at -80° C until RNA extraction, and another was fixed in 10% buffered formaldehyde solution for pathologic diagnosis and immunohistochemical staining. Histopathologic diagnosis of each tumour specimen was performed according to the International Histological Classification of Tumours by the Department of Pathology, Chiba University Hospital. Clinicopathologic staging was determined by the TNM classification of the International Union against Cancer. All OSCC samples were histologically confirmed and checked to ensure the presence of tumour in greater than 80% of specimens.

Formalin-fixed, paraffin-embedded human breast cancer tissue (positive control for Grp94 expression) was purchased from Lab Vision Co., Fremont, CA, USA (GRP94, Ab-1, positive control for immunohistochemistry).

Cell culture

The OSCC-derived cell lines used in this study were HSC-2, HSC-3, Ca9-22 (Human Science Research Resources Bank, Osaka, Japan), H-1, Sa-3 (provided by Dr Fujita, Wakayama Medical University, Wakayama, Japan), and OK-92 (established from carcinoma of the tongue in our department) (Takahashi *et al*, 1989). All OSCC-derived cell lines were cultured in Dulbecco's modified Eagle's medium F-12 HAM (Sigma-Aldrich Co., St Louis, MO, USA) supplied with 10% heat-inactivated fetal bovine serum (Sigma) and 50 U ml⁻¹ penicillin and streptomycin (Sigma). The human normal oral keratinocytes (HNOKs) cell line was cultured and maintained in defined keratinocyte-SFM (Gibco BRL, Gaithusberg, Germany). All cell lines were incubated at 37°C in a humidified atmosphere with 5% CO₂.

Protein and mRNA extraction

Protein was extracted when the cells reached 80-90% confluence; they were washed two times with phosphate-buffered saline (PBS), scraped into a tube, and centrifuged briefly. The cell pellets were incubated for 30 min in a lysis buffer (LB) containing 7 M urea, 2 M thiourea, 4% wv^{-1} CHAPS, and 10 mM Tris pH 8.0, and lysed by sonication $(3 \times 10 \text{ s pulses on ice})$. The sample was centrifuged at 13000 r.p.m. for 20 min. The supernatant containing the cell proteins then was recovered and the protein concentration was measured with a Protein Assay Kit (Bio-Rad Laboratories, Hercules, CA, USA) and adjusted to 1 mg ml⁻¹ with LB. The pH of the protein sample was adjusted to 8.5 with 30 mM Tris-HCl. Total RNA was extracted using Trizol reagent (Invitrogen Life Technologies, Carlsbad, CA, USA) according to the manufacturer's instructions. Total protein extracted from a human breast cancer cell line, SKBR3 (positive control for Grp94 expression), was obtained from Lab Vision (GRP94, Ab-1, positive control for Western blotting). Each extracted RNA or protein was stored separately at -80° C until use.

Affymetrix GeneChip hybridisation

Double-stranded cDNA was synthesised from 20 μ g of total RNA using the Superscript Choice system (Invitrogen). After phenol/ chloroform extraction and ethanol precipitation, a biotin-labelled *in vitro* transcription reaction was carried out using the cDNA template (Enzo Bioarray, Farmingdale, NY, USA). cRNA (7 μ g) was fragmented according to Affymetrix protocols and added to the



recommended hybridisation mixture. Expression profiles were created using the Human Genome U 133 Plus 2.0 arrays containing 54 675 probe sets (Affymetrix, Santa Clara, CA, USA). Arrays were stained with phycoerythrin-streptavidin, and the signal intensity was amplified by treatment with a biotin-conjugated anti-streptavidin antibody followed by a second staining with phycoerythrin-streptavidin. Arrays stained a second time were scanned using the Affymetrix GeneChip Scanner 3000 (Affymetrix). Expression data were analysed using GeneChip Operating Software 1.1 (Affymetrix), and Ca²⁺-binding protein genes were then classified by GeneSpring 6.1 (Silicon Genetics, Redwood City, CA, USA).

Network and gene ontology analysis

A list of Ca²⁺-binding protein genes identified by microarray analysis was used for network and gene ontology analyses. Gene accession numbers were imported into the Ingenuity Pathway Analysis (IPA) Software (Ingenuity Systems, Mountain View, CA, USA). The IPA database consists of proprietary ontology representing 300 000 biologic objects ranging from genes, proteins, and molecular and cellular processes. More than 11 200 human genes are currently represented in the database. The genes were categorised based on location, cellular components, and reported or suggested biochemical, biologic, and molecular functions using the software. The identified genes also were mapped to genetic networks available in the Ingenuity database and then ranked by score. The score is the probability that a collection of genes equal to or greater than the number in a network could be achieved by chance alone. A score of 3 indicates a 1/1000 chance that the focus genes are in a network due to random chance. Therefore, a score of 3 or higher has a 99.9% confidence level of not being generated by random chance alone. This score was used as the cutoff for identifying gene networks.

Western blot analysis

Among genes identified, glucose-regulated protein 94 kDa (Grp94) was selected for further analyses. We carried out Western blot analysis to investigate Grp94 protein expression in normal oral epithelium and OSCC-derived cell lines. Protein extracts were electrophoresed on 11% sodium dodecyl sulphate-polyacrylamide gel electrophoresis gels, transferred to polyvinylidene difluoride (PVDF) membranes (Bio-Rad Laboratories, Hercules, CA, USA), and blocked for 1 h at room temperature in 5% skim milk. Immunoblot PVDF membranes were washed with 0.1% Tween-20 in TBS (TBS-T) five times, and $2 \mu \text{g ml}^{-1}$ affinity-purified rat antihuman Grp94 monoclonal antibody (Lab Vision) was added directly to the TBS-T solution for 2 h at room temperature. The PVDF membranes were washed again and incubated with 1:10000 ratio of horseradish peroxidase (HRP)-conjugated antirat IgG Ab (Santa Cruz Biotechnology, Santa Cruz, CA, USA) as a secondary antibody for 20 min at room temperature. The membranes were then incubated with enhanced chemiluminescence (ECL) + HRP substrate solution included in the ECL+ kit (Amersham Biosciences, Buckinghamshire, UK), and immunoblotting was visualised by exposing the membrane to Hyperfilm (Amersham Biosciences). Triplicate examinations were performed to confirm the specificity of the antibody.

Immunofluorescence

The cells grown on glass coverslips were washed with PBS, fixed in 4% paraformaldehyde for 10 min at 37° C followed by absolute methanol for 10 min at 4° C, and blocked in PBS containing 1% skimmed milk for 10 min. The samples then were incubated with anti-Grp94 antibody (Lab Vision) at a dilution of 1:100 for 2 h, rinsed two times with PBS, and incubated with goat antirat

secondary antibody labelled with Alexa Fluor 546 (Molecular Probes, Leiden, The Netherlands) for 1 h. For counterstaining of the nucleus, the dishes were then incubated with $1 \,\mu \mathrm{g}\,\mathrm{m}\mathrm{l}^{-1}$ Cellstain-DAPI (Dojindo Laboratories, Kumamoto, Japan) in PBS for 10 min. The samples were observed under a Leica TCS2-MP confocal system (Leica Laserteknik, Mannheim, Germany) and Coherent Mira tunable pulsed titanium sapphire laser (Coherent Laser Group, Santa Clara, CA, USA).

Immunohistochemistry

To examine the cellular distribution of Grp94 protein in normal and OSCCs, we carried out immunohistochemical staining on 4- μ m sections of paraffin-embedded specimens. Briefly, after deparaffinisation and hydration, the slides were pretreated in 10 mM sodium citrate buffer (pH 6.0) in a microwave oven for 5 min at 95°C. The endogenous peroxidase activity was quenched

Table I Genetic networks in the OSCC-derived cell lines

Network	Genes in ingenuity networks ^a	Function	Score ^b
I	ATGRI, BDKRB2, CACNAIB, CACNAIC, CACNAIS, CACNBI, CACNGI, CAPN3, CAPN9, CCL2, CCL3, CCL5, CCL7, CCL13, CCL14, CCL15, CCR1, CCR5, CSPG2, CXCL12, DSG2, EGF, EPS15, F10, MYL2, NOS2A, PLA2G10, PLA2G2A, PLCD1, PLEK, PRKCA, RASGRP3, RYR1, TGM1, GEP94	Cell signalling, cellular function, and maintenance	71
2	AGTI, ANXAI, ATFI, CALML3, CDHII, COPS5, DIABLO, DGS3, FGFIO, FPRI, ILII, IRIRN, INHBA, LTBP2, MGP, MMP8, MMP13, MTPN, NOTCH4, PCDH7, RXRA, SI00A7, SI00A8, SI00A9, SAAI, SCARBI, SCUBEI, SCUBE9, SERPINA3, SODI, TNF, TNRSF4, TNRSF9, TPO, UMOD	Lipid metabolism, molecular transport	20
3	AKTI, ATP2A3, BAG3, BBC3, BCL2, CALU, CCLI9, CCR7, CDH5, CHGB, CHP, CIBI, CKS2, CLCN3, E2F4, EGF, ETSI, ITPR3, JUB, KRAS, LOCG3928, MIG-6, MYLI, PPP2R3A, PRKCSH, RALGDS, RASGRPI, RBLI, ROCKI, RPS6KB2, RRM1, RRM2, SIC3A2, SIC9A1, TERT	Cancer, cell morphology, cellular compromise	20
4	AHNAK, ATP2AI, BBC3, BGN, BMPI, CANX, CCR6, CDKN IB, CHGA, CHRD, COL5AI, CTSD, CUBN, DDX5, DMPI, FBLNS, FREQ, GFAP, GHR, IER3, IGFI, INPPLI, LGALS3, LOX, LPL, METAP2, MRCI, MTPN, MYC, PLSI, SI00AI, SI00B, SLC2A4, TLLI	DNA replication, recombination, and developmental disorder, carbohydrate	17
5	C2, C3, C1QA, C1S, C3AR1, CCL2, DSC1, DSG1, F13A1, H0XD3, IGFBP2, IGFBP5, IL13, ITIH1, JUP, KLK5, MMP2, MMP13, MMP14, MYOD1, PROC, RP6SKA2, S100A4, SERPINA1, SLIT2, SPOK3, TGM2, THBS1, THBS2, TIMP1, TKT, TNNC1, TNNC2, TNNI1, TNNT2	Organism, injury and abnormalities, cardiovascular system	17
6	ACTN2, ACTN3, CALM1, CASQ1, CD47, CDK6, CDKN1A, CTSK, CXCL12, DCN, ERBB2, ERBB3, ERBB4, EREG, FBLN2, FBN1, GHR, GRB7, GRK1, HAS2, HBEGF, HRC, IFNA2, IL6R, MTPN, MYL9, MYOZ1, PTGS1, RCV1, S100A4, SERPINA3, TGM1, TIMP3, TNC, TRDN	Cellular growth and proliferation, cellular movement, cell death	17

^aGenes in boldface were identified by microarray analysis to be expressed differentially more than two-fold in OSCCs. Other genes were either not on the expression array or not significantly regulated. ^bA score > 3 was considered significant.



Figure 1 Network with the highest score (network 1). Expression levels of all 35 genes (100%) are altered significantly in the oral squamous cell carcinomas (OSCCs). Functional relationships between gene products based on known interactions in ingenuity pathway analysis (IPA) knowledge are described.

by 30-min incubation in a mixture of 0.3% hydrogen peroxide solution in 100% methanol. After being washed with PBS buffer, sections then were incubated with primary antibody antiGrp94 antibody (1:100 dilution) at room temperature in a moist chamber for 2 h. After being washed with PBS buffer, the slides were treated with biotinylated secondary antibody for 1 h followed by colour development in 3,3'-diaminobenzidine tetrahydrochloride (Dako Japan Inc., Kyoto, Japan). Finally, the slides were lightly counterstained with haematoxylin. A known positive breast cancer control section for Grp94 was clearly stained. While the breast cancer cells were immunohistochemically positive for Grp94, those of the negative control prepared by omitting the primary antibody were negative, thus confirming the staining specificity. To quantitate the state of the Grp94 protein expression, the mean percentage of positive tumour cells was determined in at least five random fields at \times 400 magnification in each section. The intensity of the Grp94 immunoreaction was scored as follows: 1 +, weak; 2 +, moderate; and 3+, intense. The percentage of positive tumour cells and the staining intensity were then multiplied to produce a Grp94immunohistochemical staining score. Cases with a Grp94-immunohistochemical staining score exceeding 65.56 (maximum score

 Table 2
 Thirty-five focus genes in network |

705

of normal tissues) were considered positive. Two independent pathologists, neither of whom had knowledge of the patients' clinical status, made these judgments.

mRNA expression analysis

The expression levels of *Grp94* mRNA were examined in 50 OSCC specimens from patients with primary tumours among the OSCC cases studied by immunohistochemical staining. Control reactions were prepared in parallel without reverse transcriptase (RT). Before cDNA synthesis, residual genomic DNA was removed from the total RNA using DNase I treatment (DNA-free; Ambion, Austin, TX, USA). The primer sequences used to analyse *Grp94* mRNA expression were 5'-AGCAAGACGTGTTCGATTC-3' (nucleotides 1392–1410) and 5'-CCTCAATTTTGTCAAGGGTG-3' (nucleotides 1607–1626). The sequences of specific primers were checked before use to avoid amplification of genomic DNA or pseudogenes by the Primer3 program (available at http:www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi). Amplified products were analysed by 3% agarose gel electrophoresis to ascertain size and purity. Real-time quantitative RT (qRT)–PCR

Affymetrix no.	Gene Molecular function		Location	Fold change ^a	
217901_at	DSG2	Cell adhesion	Plasma membrane	85.187	
200599_s_at	Grp94	Protein folding and sorting, antigen presentation	Plasma membrane	5.479	
210037_s_at	NÖS2A	Nitric oxide synthase activity	Cytoplasm	5.072	
1560074_at	PRKCA	Protein kinase C activity	Cytoplasm	3.781	
210641_at	CAPN9	Calpain activity	Unknown	3.003	
207162_s_at	CACNAIB	Voltage-gated calcium channel activity	Plasma membrane	2.894	
207222_at	PLA2G10	Unknown	Extracellular space	2.749	
207225_at	EPS15	Protein binding	Plasma membrane	0.4539	
201511_at	BDKRB2	Receptor activity	Plasma membrane	0.4464	
205434_s_at	CACNAIC	Voltage-gated calcium channel activity	Plasma membrane	0.4288	
218075_at	AGTRI	Receptor activity	Plasma membrane	0.3305	
1558450_at	CCRI	Chemokine receptor activity	Plasma membrane	0.2580	
207798_s_at	CACNBI	Voltage-gated calcium channel activity	Plasma membrane	0.2346	
228767_at	CCL13	Chemokine receptor activity	Extracellular space	0.2200	
235070_at	PLEK	Reorganisation	Cytoplasm	0.1697	
229819_at	CCR5	Chemokine receptor activity	Plasma membrane	0.1640	
228376_at	CCL7	Chemokine receptor activity	Extracellular space	0.1611	
218529_at	FIO	Peptidase activity	Extracellular space	0.1583	
224938_at	PLA2G2A	Protein binding	Extracellular space	0.1339	
224939_at	CACNAIS	Voltage-gated calcium channel activity	Plasma membrane	0.1256	
2 47 _at	CCL5	Chemokine receptor activity	Extracellular space	0.1172	
214319_at	CCL3	Chemokine receptor activity	Extracellular space	0.1071	
1555904_at	CCL15	Chemokine receptor activity	Extracellular space	0.0868	
209665_at	RASGRP3	Protein binding	Cytoplasm	0.0810	
220781_at	CACNGI	Voltage-gated calcium channel activity	Plasma membrane	0.0744	
1553086_at	CSPG2	Unknown	Extracellular space	0.0736	
212698_s_at	EGF	Protein binding	Extracellular space	0.0611	
226627_at	PLCD I	Unknown	Cytoplasm	0.0381	
213666_at	CXCL12	Chemokine receptor activity	Extracellular space	0.0350	
211890_x_at	CAPN3	Calpain activity	Cytoplasm	0.0227	
216598_s_at	CCL2	Chemokine receptor activity	Extracellular space	0.0198	
235340_at	CCL14	Chemokine receptor activity	Extracellular space	0.0165	
205485_at	RYRI	Voltage-gated calcium channel activity	Plasma membrane	0.0126	
206008_at	TGMI	Cell formation	Plasma membrane	0.0109	
209742_s_at	MYL2	Motor activity	Cytoplasm	0.0100	

AGTR1 = angiotensin II receptor, type 1; BDKRB2 = bradykinin receptor B2; CACNA1B = calcium channel, voltage-dependent, L type, α 1B- subunit; CACNA1C = calcium channel, voltage-dependent, L type, α 1C subunit; CACNA1S = calcium channel, voltage-dependent, L type, α 1S subunit; CACNB1 = calcium channel, voltage-dependent, β 1 subunit; CACNG1 = calcium channel, voltage-dependent, γ subunit 1; CAPN3 = calpain 3, (p94); CAPN9 = calpain 9; CCL2 = chemokine (C-C motif) ligand 2; CCL3 = chemokine (C-C motif) ligand 3; CCL5 = chemokine (C-C motif) ligand 5; CCL7 = chemokine (C-C motif) ligand 7; CCL13 = chemokine (C-C motif) ligand 13; CCL14 = chemokine (C-C motif) ligand 14; CCL15 = chemokine (C-C motif) ligand 15; CCR1chemokine (C-C motif); CCR5 = chemokine (C-C motif) receptor 5; CSPG2 = chondroitin sulfate proteoglycan 2; CXCL12 = chemokine (C-C-C motif) ligand 12; DSG2 = desmoglein 2; EGF = epidermal growth factor; EPS15 = epidermal growth factor; cereptor pathway substrate 15; F10 = coagulation factor X; Grp94 = glucose regulated protein; 94 kDa; MYL2 = myosin, light polypeptide 2, regulatory, cardiac, slow; NOS2A = nitric oxide synthase 2A; PLA2G2A = phospholipase A2, group IIA; PLA2G10 = phospholipase A2, group X; PLCD1 = phospholipase C, delta 1; PLEK = pleckstrin; PRKCA = protein kinase C, α ; RASGRP3 = RAS guanyl releasing protein 3; RYR1 = ryanodine receptor 1; TGM1 = transglutaminase 1. ^aFold overexpression for microarray data of OSCC-derived cell lines compared to normal control.

Ca²⁺-binding protein genes in OSCCs H Nomura et al

was performed with a single method using the LightCycler FastStart DNA Master SYBR Green I kit (Roche, Mannheim, Germany). To prepare the standard curve, $3 \mu g$ of total RNA from normal oral tissue was reverse-transcribed with Superscript RT (Life Technologies, Grand Island, NY, USA) and oligo-d(T)12-18 primer, after which serial dilutions were made corresponding to cDNA transcribed from 300, 30, 3.0, and 0.3 ng of total RNA. PCRs using LightCycler apparatus were carried out in a final volume of $20\,\mu$ l of reaction mixture consisting of $2\,\mu$ l of FirstStart DNA Master SYBR Green I mix, 3 mM MgCl₂, and 0.2 µl of primers, according to the manufacturer's instructions. The reaction mixture then was loaded into glass capillary tubes and submitted to an initial denaturation at 95°C for 10 min, followed by 45 rounds of amplification at 95°C 10s) for denaturation, 58°C (10s) for annealing and 72°C for extension, with a temperature slope of 20°C s⁻¹, performed in the LightCycler. The transcript amount for Grp94 was estimated from the respective standard curves and normalised to the GAPDH transcript amount determined in corresponding samples.

Statistical analysis

Differences in gene expression levels between Grp94-positive and Grp94-negative cases were calculated with the Mann-Whitney's U-test. Correlations between Grp94-immunohistochemical staining scores and clinicopathologic features were evaluated by Fisher's exact test. Overall survival time was defined as the interval between the date of treatment and the date of death or until the last objective follow-up information was obtained. Disease-free survival time was regarded as the time interval between tumour treatment and detection of the first locoregional recurrence, distant metastasis, or both or the date of the last follow-up, whichever occurred first. Patients without evidence of disease (local recurrence or metastasis) during follow-up were considered to have a good prognosis; patients with local recurrence, distant metastasis during follow-up were regarded as having a poor prognosis. Survival curves were obtained by the Kaplan-Meier method and differences in survival rates between Grp94-positive and Grp94-negative cases were compared by log-rank tests with 95% significance. The criterion for statistical significance was P < 0.05. The data are expressed as the mean values \pm s.e.

RESULTS

Altered Ca²⁺⁻binding protein genes expression in **OSCC-derived cell lines**

To investigate the expression profile changes of Ca^{2+} -binding protein genes in OSCC, we initially performed oligonucleotide microarray analyses using the Affymetrix U133A chips, which contained 54 675 probe sets with RNAs isolated from four OSCCderived cell lines (HSC-2, HSC-3, H-1, and Sa-3). Control RNAs consisted of a pool made of normal tongue tissue RNAs extracted from three patients. Expression data were analysed using the GeneChip Operating Software 1.1 (Affymetrix) and GeneSpring 6.1 (Silicon Genetics). Of the 2241 Ca²⁺-binding protein gene transcripts analysed, 100 (4.4%) were identified as genes that were differentially expressed at least two-fold in all OSCC-derived cell lines examined compared with controls. Of these, 24 genes were found to be upregulated and 76 genes were downregulated in OSCC (data not shown).

Network and gene ontology analysis

On the basis of all genes identified as described previously (referred to as focus genes), new and expanded pathway maps and connections and specific gene-gene interactions were inferred, functionally analysed, and used to build on the existing pathway

using the IPA knowledge base. To generate networks, the knowledge base was queried for interactions between focus genes and all other gene objects stored in the database. Six networks were found to be significant in OSCC in that they had more of the identified genes present than would be expected by chance (Table 1). Of them, the network with the highest score (network 1, score = 71) was generated with all focus genes (Figure 1; Table 2). We also performed gene ontology analyses of 100 focus genes using the IPA tool. Seventy-six upregulated genes were associated with cancer-related functions, cell death, and the cell cycle; 24 downregulated genes were associated with cellular movement, signalling, and function and maintenance (Table 3). In particular, we found that the cellular signalling was contained in both the upregulated and downregulated groups. Six upregulated genes were associated with a variety of cellular signalling functions, including induction/production of nitric oxide, inhibition of cyclic adenosine monophosphate, and inactivation of mitogen-activated protein kinase. In contrast, 31 downregulated groups played a critical role in cellular signalling, which is related

Table 3 Functional characterisation of Ca²⁺-binding protein genes altered expressed in OSCC-derived cell lines

Relevant function and disease	P-value	Gene, no ^a	
Cancer ^b	1.15E-3 to	7	
	4.27E-2		
Cell death	1.15E-3 to	4	
	6.42E-2		
Cell cycle	1.86E-3 to	6	
	6.95E-2		
Cell-to-cell signalling and interaction	3.94E-3 to	7	
0 0	6.16E-2		
Cell signalling	3.29E-3 to	6	
	6.42E-2		
Molecular transport	4.29E-3 to	5	
	7.47E-2		
Small molecule biochemistry	4.29E-3 to	9	
	7.47E-2		
Nucleic acid metabolism	5.07E-3 to	4	
	2.89E-2		
Cellular growth and proliferation	5.07E-3 to	3	
0	6.95E-2		
Cell morphology	8.24E-3 to	I	
	6.95E-2		

Top-10 functions associated with downregulated genes in OSCC-derived cell lines Cellular movement^c 1.60E-13 to 33

		8.53E-3	
	Cellular signalling	1.52E-11 to	31
		8.53E-3	
	Cellular function and maintenance	1.52E-11 to	16
		8.53E-3	
	Small molecule biochemistry	1.52E-11 to	20
		8.53E-3	
	Vitamin and mineral metabolism	1.52E—11 to	32
		7.02E-3	
	Tissue morphology	2.10E-11 to	33
		8.53E-3	
	Molecular transport	2.69E—11 to	18
		8.53E-3	
	Skeletal and vascular system development and	2.08E-10 to	21
fu	nction	8.53E-3	
	Cancer	5.13E—9 to	27
		8.53E—3	
	Cell-to-cell signalling and interaction	1.01E-8 to	26
		8531-3	

^aNumber of associated genes. ^bApoptosis, cell movement, proliferation, cell cycle progression, migration, binding quantity of cancer cell line. ^cChemotaxis, migration, cell movement, mobilisation of haematologic cyte.

Ca²⁺-binding protein genes in OSCCs H Nomura et *al*

<u>Molecular Diagnostics</u>

to intracellular Ca^{2+} homeostasis, suggesting that genes associated with cell signalling may act in a different manner during development of OSCC.

Among genes mapped to the network with the highest significance, *Grp94* was analysed further.

GRP94 expression in OSCCs

The state of Grp94 protein expression in six OSCC-derived cell lines (HSC-2, HSC-3, H-1, Sa-3, Ca9-22, and OK-92) (n = 6) was evaluated by Western blot analysis. Figure 2A shows representative results. The band size was found to be 94 kDa as a single band, as reported by Little *et al* (1994). A significant (P<0.001) over-expression of Grp94 protein was observed in all OSCC-derived cell lines compared with normal oral epithelium. We also assessed the level of Grp94 protein expression in an OSCC-derived cell line (HSC-3) and HNOKs by immunofluorescence analysis. Represen-

tative cases of immunofluorescence are shown in Figure 2B. Strong immunoreactivity of Grp94 protein was detected in the cytoplasm of the OSCC-derived cell line HSC-3 compared with the HNOKs.

Immunohistochemical staining was performed using a series of surgical OSCC specimens, including 80 OSCCs with corresponding normal tissues and 20 OPLs. Grp94 overexpression was found not only in OSCCs (P < 0.001) but also in OPLs (P < 0.001). Among the OSCCs, 46 had significantly increased expression of Grp94 (immunohistochemical staining score, > 65.56). In contrast, the normal tissues had no or significant downregulation of Grp94 expression and were considered Grp94 negative (Fisher's exact test). There was no significant difference between the frequency of Grp94-positive cases and clinicopathologic features (Table 4).

Thirteen of 20 (65%) OPLs were defined as Grp94 positive. The Grp94-immunohistochemical staining scores for normal tissues, OPLs, and OSCCs, respectively, ranged from 0 to 65 (mean, 12.173), 12 to 191 (mean, 85.27), and 0 to 262 (mean, 106.511)



Figure 2 Representative results of expression of Grp94 protein in oral squamous cell carcinomas (OSCC)-derived cell lines. (A) Western blot analysis of Grp94 protein in OSCC-derived cell lines and normal oral epithelium. All OSCC-derived cell line extracts exhibit a single band for Grp94 protein expression at high levels. In contrast, normal oral epithelium shows a low level of Grp94 protein expression. (B) Immunocytochemical analysis shows strong immunoreactivity of Grp94 in an OSCC-derived cell line (HSC-3) compared with human normal oral keratinocytes (HNOKs). DAPI staining was used to stain DNA. Bar, $100 \,\mu$ m. (C) Immunohistochemical staining of Grp94 in normal tissue, oral premalignant lesion (OPL), and primary OSCC. (a) Normal oral tissue exhibits negative Grp94 protein expression. (b) Grp94-negative case of OSCC. (c) Grp94-positive case of OPL. The immunoreaction is enhanced in the spinous layer. (d) Grp94-positive case of OSCC. Strong positive immunoreactivity for Grp94 is detected in the cytoplasm.



 Table 4
 Correlation between Grp94 expression and clinical classification in OSCCs

		Result of imm number of	Result of immunostaining: number of patients(%)	
Clinical classification	Total	Grp94 (–)	Grp94 (+)	P-value ^a
Age at surgery (years)				
<60	23	8 (30)	15 (70)	
60≤, <70	26	11 (42)	15 (58)	0.431974
70≼	31	15 (48)	16 (52)	
Gender				
Male	33	13 (39)	20 (61)	0.654387
Female	47	21 (45)	26 (55)	
T-primary tumour				
TI	5	2 (40)	3 (60)	
T2	35	15 (43)	20 (57)	0.55542
Т3	17	5 (29)	12 (71)	
T4	23	12 (52)	11 (48)	
N-regional lymph node				
N(-)	52	23 (44)	29 (56)	0.8 3 85
N(+)	28	(39)	17 (61)	
Stage				
I	5	2 (40)	3 (60)	
II	16	7 (44)	9 (56)	0.756529
III	17	7 (41)	10 (59)	
IV	52	28 (54)	24 (46)	
Histopathologic type				
Well differentiated	51	24 (47)	27 (53)	
Moderately differentiated	22	9 (41)	13 (59)	0.300615
Poorly differentiated	7	(4)	6 (86)	
Tumour site				
Tongue	34	12 (35)	22 (65)	
Gingiva	29	13 (45)	16 (55)	0.90283
Oral floor	7	3 (43)	4 (57)	
Buccal mucosa	6	3 (50)	3 (50)	
Oropharynx	3	2 (67)	l (33)	
Lip	1	0 (0)	(100)	
Leukoplakia	20	7 (35)	13 (65)	

 $^{a}P < 0.05$ was considered significant.



Figure 3 State of Grp94 protein expression in normal tissues (n = 80), oral premalignant lesions (OPLs) (n = 20) and primary oral squamous cell carcinomas (OSCCs) (n = 80). Grp94 protein expression in OPLs and OSCCs is significantly higher than in normal oral tissues (P < 0.001, Mann–Whitney's U-test). The results are expressed as the mean ± s.d.



Figure 4 *Grp94* mRNA expression status in primary oral squamous cell carcinomas (OSCCs) and OSCC-derived cell lines. (**A**) Comparison of *Grp94* mRNA expression levels between Grp94-positive and Grp94-negative cases classified by immunohistochemical analysis. There is a significant difference in the *Grp94* mRNA expression levels between the negative and positive cases (P < 0.001, Mann–Whitney's *U*-test). (**B**) Quantification of mRNA levels in OSCC-derived cell lines by qRT–PCR analysis. Significant upregulation of *Grp94* mRNA expression is seen in all OSCC-derived cell lines examined compared to normal oral epithelium. Data are expressed as the means \pm s.d.

(Figure 2C). Grp94 expression levels in primary OSCCs and OPLs were significantly (P < 0.001) higher than those in normal oral tissues (Figure 3). In contrast, we found no significant (P = 0.448334) difference in Grp94-immunohistochemical staining scores between OSCCs and OPLs.

Grp94 mRNA expression levels were significantly upregulated in primary tumours of randomly selected Grp94-positive cases (n = 29) compared to selected Grp94-negative cases (n = 21,Mann-Whitney U-test, P < 0.001; Figure 4A). mRNA expression levels were normalised to GAPDH. The relative mRNA expression levels in positive and negative cases ranged from 113 to 297 (mean, 191.3) and 9 to 88 (mean, 56.6), respectively. Grp94 expression was upregulated significantly in all OSCC cell lines examined compared to normal oral epithelium used as a control (Figure 4B).

Prognostic significance of GRP94 expression in OSCCs

To assess whether Grp94 expression also had a prognostic impact on patients with OSCC, clinical postoperative data from patients





Figure 5 Kaplan-Meier curves for disease-free (A) or overall survival (B) according to Grp94-positive expression in patients with oral squamous cell carcinoma (OSCC) (log-rank test (A), P=0.011 (B), P=0.024). Cases with significantly increased expression of Grp94 protein (immunohistochemical staining score, >65.56; maximum score of normal tissues) are considered Grp94 positive. Overexpression of Grp94 protein is significantly associated with a poor outcome in patients with OSCC.

whose tissue samples were investigated for Grp94 protein expression by immunohistochemical analysis were statistically analysed. Tumours with significantly increased expression levels of Grp94 protein (immunohistochemical staining score, >65.56; maximum score of normal tissues) were defined as Grp94 positive (n = 46). Cases with no or significant downregulation of Grp94 protein expression (immunohistochemical staining score, <65.56) were considered Grp94 negative (n = 34).

Survival curves in relation to Grp94 protein expression are shown in Figure 5. Log-rank survival analysis indicated that Grp94-positive expression was significant both for disease-free survival (P = 0.011; Figure 5A) and overall survival (P = 0.024; Figure 5B), which suggested that high expression of Grp94 protein in patients with OSCCs was significantly associated with a poor prognosis.

DISCUSSION

Using a microarray technique, we identified a total of 100 Ca^{2+} binding protein genes aberrantly expressed in OSCC cells. Pathway analysis could further characterise six networks from the 100 genes (Table 1). In addition, the network of the highest significance was generated entirely from 35 of the 100 focus genes (Figure 1), including a number of the CCL and CXC families. Previous studies have shown that chemokine-mediated JAK/STAT activation is critical for phospholipase C- β dependent Ca²⁺ flux (Li *et al*, 2000; Lewis, 2001; Thelen, 2001; Soriano et al, 2003). This evidence together with the current results suggests that reduced expression of the main chemokine families, CCL and CXC, disrupts intracellular Ca²⁺ homeostasis and initiates oral tumorigenesis. The network with the highest significance also contained several cancer-related genes with high expression levels, including DSG2, PRKCA, PLA2G10, and Grp94.

Considerable evidence has shown a significant association between the heat-shock protein 90 (Hsp90) and a wide range of human malignancies, including head and neck cancers (Ito et al, 1998; Baker et al, 2005; Yin et al, 2005). Grp94, also known as gp96, is the ER-resident member of the Hsp90 family constitutively expressed in virtually all cell types (Van et al, 1989) and the most abundant ER chaperon protein showing high homology (50%) to cytosolic counterpart Hsp90 (Sorger and Pelham, 1987). Increased expression of Grp94 both at the mRNA and protein levels also has been reported in several types of human cancers, such as oesophageal cancer (Wang et al, 2005b), lung cancer (Wang et al, 2005a), breast cancer (Gazit et al, 1999), liver cancer (Lim et al, 2005), and colon cancer (Wang et al, 2005c). We hypothesised that Grp94 also has potential as an emerging therapeutic target of interest for the treatment of oral cancer. However, the status of Grp94 in OSCC remains unclear and, therefore, we selected it for further investigation.

To confirm our hypothesis, we determined the protein/mRNA expression in a series of OSCC-derived cell lines and human primary OSCCs using immunofluorescence, Western blot analysis, qRT-PCR, and immunohistochemistry. Significant increases in Grp94-protein- and mRNA-expression levels were observed in the OSCC-derived cell lines examined compared with the normal oral epithelium. We also detected a comparatively strong tumour celllocalised cytoplasmic Grp94-immunoreaction in primary OSCCs. By evaluating the Grp94 immunohistochemistry scores, significant upregulation was evident in the primary OSCCs compared with normal tissues. While we could not find a significant correlation between Grp94 protein expression status and any clinicopathologic features examined, high levels of Grp94 protein expression was detected even in the OPLs examined. In addition, the current study showed that Grp94 overexpression is closely related to the diseasefree/overall survival (P = 0.011 and 0.024, respectively). Thus, we suggest that Grp94 could be associated with not only early-stage OSCC development but also tumour prognosis.

Interestingly, a recent study has shown that inhibition of Grp94 expression by geldanamycin in chronic lymphocytic leukaemia cells induces apoptosis with modest cytoprotective effects of primary haematopoietic progenitors from normal bone marrow (Jones *et al*, 2004). In addition, targeting cancer cells with an antisense or RNAi procedure against Grp94 has shown increased chemosensitivity or radiosensitivity (Reddy *et al*, 1999; Kubota *et al*, 2005). From the therapeutic standpoint, this evidence may provide a novel/effective approach for treating human OSCCs.

In summary, we found novel specific networks of Ca^{2+} -binding protein genes in OSCC cells and identified several candidate genes for molecular targeting, especially for *Grp94*. Our findings may contribute to an understanding of key biologic functions and pathways of certain Ca^{2+} -binding protein genes associated with

REFERENCES

- Baker H, Patel V, Molinolo AA, Shillitoe EJ, Ensley JF, Yoo GH, Meneses-Garcia A, Myers JN, El-Naggar AK, Gutkind JS, Hancock WS (2005) Proteome-wide analysis of head and neck squamous cell carcinomas using laser-capture microdissection and tandem mass spectrometry. Oral Oncol 2: 183-199
- Berridge MJ, Lipp P, Bootman MD (2000) The versatility and universality of calcium signalling. Nat Rev Mol Cell Biol 1: 11-21
- Bustin SA, Li SR, Dorudi S (2001) Expression of the Ca2+-activated chloride channel genes CLCA1 and CLCA2 is downregulated in human colorectal cancer. DNA Cell Biol 20: 331-338
- Chen D, Davies MP, Rudland PS, Barraclough R (1997) Transcriptional down-regulation of the metastasis-inducing S100A4 (p9Ka) in benign but not in malignant rat mammary epithelial cells by GC-factor. *J Biol Chem* **272**: 20283-20290
- Ding Y, Robbins J, Fraser SP, Grimes JA, Djamgoz MB (2006) Comparative studies of intracellular Ca²⁺ in strongly and weakly metastatic rat prostate cancer cell lines. *Int J Biochem Cell Biol* **38**: 366–375
- Durham AC, Walton JM (1982) Calcium ions and the control of proliferation in normal and cancer cells. *Biosci Rep* 2: 15-30
- Endo Y, Uzawa K, Mochida Y, Shiiba M, Bukawa H, Yokoe H, Tanzawa H (2004) Sarcoendoplasmic reticulum Ca(2+) ATPase type 2 downregulated in human oral squamous cell carcinoma. *Int J Cancer* 110: 225-231
- Gazit G, Lu J, Lee AS (1999) De-regulation of GRP stress protein expression in human breast cancer cell lines. *Breast Cancer Res Treat* 2: 135-146
- Haeseleer F, Sokal I, Verlinde CLMJ, Erdjumnet-Bromage H, Tempst P, Pronin AN (2000) Ca (2+)-binding proteins in the retina: structure, function, and the etiology of human visual diseases. J Biol Chem 275: 1247-1260
- Higo M, Uzawa K, Kawata T, Kato Y, Kouzu Y, Yamamoto N, Shibahara T, Mizoe JE, Ito H, Tsujii H, Tanzawa H (2006) Enhancement of SPHK1 *in vitro* by carbon ion irradiation in oral squamous cell carcinoma. *Int J Radiat Oncol Biol Phys* **65**: 867–875
- Huang JK, Huang CJ, Chen WC, Liu SI, Hsu SS, Chang HT, Tseng LL, Chou CT, Chang CH, Jan CR (2005) Independent [Ca²⁺]i increases and cell proliferation induced by the carcinogen safrole in human oral cancer cells. Naunyn Schmiedebergs Arch Pharmacol 372: 88-94
- Imazawa M, Hibi K, Fujitake S, Kodera Y, Ito K, Akiyama S, Nakao A (2005) S100A2 overexpression is frequently observed in esophageal squamous cell carcinoma. Anticancer Res 25: 1247-1250
- Ishigami T, Uzawa K, Higo M, Nomura H, Saito K, Kato Y, Nakashima D, Shiiba M, Bukawa H, Yokoe H, Kawata T, Ito H, Tanzawa H (2007) Genes and molecular pathways related to radioresistance of oral squamous cell carcinoma cells. *Int J Cancer* **120**: 2262–2270
- Ito T, Kawabe R, Kurasono Y, Hara M, Kitamura H, Fujita K, Kanisawa M (1998) Expression of heat shock proteins in squamous cell carcinoma of the tongue: an immunohistochemical study. J Oral Pathol Med 1: 18-22
- Jones DT, Addison E, North JM, Lowdell MW, Hoffbrand AV, Mehta AB, Ganeshaguru K, Folarin NI, Wickremasinghe RG (2004) Geldanamycin and herbimycin A induce apoptotic killing of B chronic lymphocytic leukemia cells and augment the cells' sensitivity to cytotoxic drugs. *Blood* 5: 1855-1861
- Kasamatsu A, Endo Y, Uzawa K, Nakashima D, Koike H, Hashitani S, Numata T, Urade M, Tanzawa H (2005) Identification of candidate genes associated with salivary adenoid cystic carcinomas using combined comparative genomic hybridization and oligonucleotide microarray analyses. Int J Biochem Cell Biol 37: 1869-1880

OSCC and should stimulate further investigation into Ca^{2+} -binding protein genes relevant to oral carcinogenesis.

ACKNOWLEDGEMENTS

We thank Lynda C Charters for editing this article. This study was partly supported by a Grant-in-Aid Scientific Research (no. 16591820) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

- Kato H, Uzawa K, Onda T, Kato Y, Saito K, Nakashima D, Ogawara K, Bukawa H, Yokoe H, Tanzawa H (2006) Down-regulation of 1D-myoinositol 1,4,5-trisphosphate 3-kinase A protein expression in oral squamous cell carcinoma. *Int J Oncol* **28:** 873-881
- Koch G, Smith M, Macer D, Webster P, Mortara R (1986) endoplasmic reticulum contains a common, abundant calcium-binding glycoprotein, endoplasmin. J Cell Sci 86: 217-232
- Kubota H, Suzuki T, Lu J, Takahashi S, Sugita K, Sekiya S, Suzuki N (2005) Increased expression of GRP94 protein is associated with decreased sensitivity to X-rays in cervical cancer cell lines. *Int J Radiat Biol* 81: 701-709

Kunzelmann K (2005) Ion channels and cancer. J Membr Biol 205: 159-173

- Lakshmikuttyamma A, Selvakumar P, Kanthan R, Kanthan SC, Sharma RK (2004) Overexpression of m-calpain in human colorectal adenocarcinomas. *Cancer Epidemiol Biomarkers Prev* 13: 1604–1609
- Lewis RS (2001) Calcium signaling mechanisms in T lymphocytes. Annu Rev Immunol 19: 497-521
- Li Z, Jiang H, Xie W, Zhang Z, Smrcka AV, Wu D (2000) Roles of PLC-2 and -3 and the PI3K in chemoattractant-mediated signal transduction. *Science* 287: 1046–1049
- Lim SO, Park SG, Yoo JH, Park YM, Kim HJ, Jang KT, Cho JW, Yoo BC, Jung GH, Park CK (2005) Expression of heat shock proteins (HSP27, HSP60, HSP70, HSP90, GRP78, GRP94) in hepatitis B virus-related hepatocellular carcinomas and dysplastic nodules. 14: 2072-2079
- Little E, Ramakrishnan M, Roy B, Gazit G, Lee AS (1994) The glucoseregulated proteins (GRP78 and GRP94): functions, gene regulation, and applications. *Crit Rev Eukaryot Gene Expr* **4**: 1–18
- Moriyama-Kita M, Endo Y, Yonemura Y, Heizmann CW, Schafer BW, Sasaki T, Yamamoto E (2004) Correlation of S100A4 expression with invasion and metastasis in oral squamous cell carcinoma. *Oral Oncol* **40**: 496-500
- Nelson MR, Chazin WJ (1998) Calmodulin and Signal Transduction. Academic Press: London
- Pietas A, Schluns K, Marenholz I, Schafer BW, Heizmann CW, Petersen I (2002) Molecular cloning and characterization of the human S100A14 gene encoding a novel member of the S100 family. *Genomics* 79: 513-522
- Reddy RK, Lu J, Lee AS (1999) The endoplasmic reticulum chaperone glycoprotein GRP94 with Ca(2+)-binding and antiapoptotic properties is a novel proteolytic target of calpain during etoposide-induced apoptosis. J Biol Chem 274: 28476-28483
- Saito K, Uzawa K, Endo Y, Kato Y, Nakashima D, Ogawara K, Shiba M, Bukawa H, Yokoe H, Tanzawa H (2006) Plasma membrane Ca2+ ATPase isoform 1 down-regulated in human oral cancer. Oncol Rep 15: 49-55
- Saito Y, Takazawa H, Uzawa K, Tanzawa H, Sato K (1998) Reduced expression of E-cadherin in oral squamous cell carcinoma: relationship with DNA methylation of 5' CpG island. *Int J Oncol* **12**: 293–298
- Shimada K, Uzawa K, Kato M, Endo Y, Shiiba M, Bukawa H, Yokoe H, Seki N, Tanzawa H (2005) Aberrant expression of RAB1A in human tongue cancer. *Br J Cancer* **92:** 1915–1921
- Sorger PK, Pelham HRB (1987) The glucose-regulated protein grp94 is related to heat shock protein hsp90. *J Mol Biol* **2:** 341-344
- Soriano SF, Serrano A, Hernanz-Falcon P, Martin de Ana A, Monterrubio M, Martinez C, Rodriguez-Frade JM, Mellado M (2003) Chemokines integrate JAK/STAT and G-protein pathways during chemotaxis and calcium flux responses. *Eur J Immunol* **33**: 1328-1333
- Takahashi K, Kanazawa H, Akiyama Y, Tazaki S, Takahara M, Muto T, Sato K (1989) Establishment and characterization of a cell line (SAS) from

poorly differentiated human squamous-cell carcinoma of the tongue. J Jpn Stomatol Soc 38: 20-28

- Takenaga K, Nakamura Y, Sakiyama S (1997) Expression of antisense RNA to S100A4 gene encoding an S100-related calcium-binding protein suppresses metastatic potential of high-metastatic Lewis lung carcinoma cells. *Oncogene* 14: 331-337
- Thebault S, Flourakis M, Vanoverberghe K, Vandermoere F, Roudbaraki M, Lehen'kyi V, Slomianny C, Beck B, Mariot P, Bonnal JL, Mauroy B, Shuba Y, Capiod T, Skryma R, Prevarskaya N (2006) Differential role of transient receptor potential channels in Ca2+ entry and proliferation of prostate cancer epithelial cells. *Cancer Res* **66**: 2038-2047
- Thelen M (2001) Dancing to the tune of chemokines. Nat Immunol 12: 129-134
- Tsai ST, Jin YT, Tsai WC, Wang ST, Lin YC, Chang MT, Wu LW (2005) S100A2, a potential marker for early recurrence in early-stage oral cancer. Oral Oncol **41**: 349-357
- Van Ginkel PR, Gee RL, Walker TM, Hu D-N, Heizmann CW, Polans AS (1998) The identification and differential expression of calcium-binding proteins associated with ocular melanoma. *Biochim Biophys Acta* 1448: 290-297

- Van PN, Peter F, Soling HD (1989) Four intracisternal calcium-binding glycoproteins from rat liver microsomes with high affinity for calcium. No indication for calsequestrin-like proteins in inositol 1,4,5-trisphosphate-sensitive calcium sequestering rat liver vesicles. *J Biol Chem* **264**: 17494-17501
- Wang Q, He Z, Zhang J, Wang Y, Wang T, Tong S, Wang L, Wang S, Chen Y (2005a) Overexpression of endoplasmic reticulum molecular chaperone GRP94 and GRP78 in human lung cancer tissues and its significance. *Cancer Detect Prev* 6: 544-551
- Wang XP, Liu GZ, Song AL, Chen RF, Li HY, Liu Y (2005b) Expression and significance of heat shock protein 70 and glucose-regulated protein 94 in human esophageal carcinoma. *World J Gastroenterol* 11: 429–432
- Wang XP, Qiu FR, Liu GZ, Chen RF (2005c) Correlation between clinicopathology and expression of heat shock protein 70 and glucoseregulated protein 94 in human colonic adenocarcinoma. *World J Gastroenterol* 7: 1056–1059
- Yin X, Zhang H, Burrows F, Zhang L, Shores CG (2005) Potent activity of a novel dimeric heat shock protein 90 inhibitor against head and neck squamous cell carcinoma *in vitro* and *in vivo*. *Clin Cancer Res* **10**: 3889–3896