Letter to the Editor

Human breast cancer and sexual activities

N Akil¹, A Kassab², A Yasmeen², A D Darnel^{2,3,4}, T A Bismar^{2,3,4} and A-E Al Moustafa^{*,1,2,3}¹Faculty of Medicine/University of Aleppo, Aleppo, Syria; ²Montréal Center for Experimental Therapeutics in Cancer, Lady Davis Institute for Medical Research of the Sir Mortimer B Davis-Jewish General Hospital, McGill University, Montreal, Quebec, Canada; ³Program in Cancer Genetics, Department of Oncology, McGill University, Montreal, Quebec, Canada; ⁴Department of Pathology Jewish General Hospital, McGill University, Montreal, Quebec, Canada

British Journal of Cancer (2008) 98, 508–509. doi:10.1038/sj.bjc.6604103 www.bjcancer.com

Published online 22 January 2008

© 2008 Cancer Research UK

Sir,

Several studies have reported that high-risk human papilloma-viruses (HPVs) are present in more than 50% of human breast cancers (Yu *et al*, 2000; Liu *et al*, 2001; Damin *et al*, 2004; de Villiers *et al*, 2005; Kan *et al*, 2005; Lawson *et al*, 2006a, b; Yasmeen *et al*, 2007a, b). Furthermore, high-risk HPVs of the same type were found in both cervical and breast cancer of the same patients (Hennig *et al*, 1999; Widschwendter *et al*, 2004). This finding has led to the hypothesis that HPVs could be transmitted to the breast through sexual activities (Kan *et al*, 2005; Lawson *et al*, 2006a, b). Consequently, it is possible that the incidence of HPV-positive breast cancer in young women is related to high-risk HPV genital infections, which are much more common in young women who have multiple sexual partners (IARC, 1995).

There are three studies concerning the age of women in relation to the incidence of high-risk HPV-positive breast cancer. While there were no differences in the average age of women with either HPV-positive or -negative breast cancer in Brazilian women (Damin *et al*, 2004). In contrast, two recent studies reported that the average age of HPV-positive breast cancer in Greek and Australian women are 38 and 55.6 years in comparison with HPV-

negative breast cancer, which are 53 and 63.8 years ($P=0.001$ and 0.049), respectively, (Kroupis *et al*, 2006; Lawson *et al*, 2006a, b).

We recently reported that E6/E7 of HPV type 16 is present in the majority of invasive and metastatic breast cancer in comparison to normal mammary tissues in young Canadian women (Yasmeen *et al*, 2007a, b). Moreover, we were able to demonstrate that the presence of E6/E7 of high-risk HPVs is correlated with invasive breast cancer in young Syrian women (Akil *et al*, in preparation); in this study, we documented the ages of Canadian and Syrian women in relation to their HPVs status. We found a statistically significant difference between the average age of women with high-risk HPV-positive breast cancers, which was 46.5 years in comparison with HPV-negative breast cancers of 57.5 years ($P=0.05$). Our data suggest that high-risk HPV-positive breast cancers are more frequent in younger women who are probably sexually more active than older women who have less sexual activity; thus, we firmly believe that high-risk HPV infections can be transmitted by sexual activity and could play an important role in the progression of human breast cancer. However, future epidemiological studies are necessary to confirm these findings.

REFERENCES

- Damin APS, Karam R, Zettler CG, Caleffi M, Alexandre COP (2004) Evidence for an association of human papillomavirus and breast carcinomas. *Breast Cancer Res Treatment* 84: 131–137
- de Villiers EM, Sandstrom RE, zur Hausen H, Buck CE (2005) Presence of papillomavirus sequences in condylomatous lesions of the mamillae and in invasive carcinoma of the breast. *Breast Cancer Res* 1: R1–R11
- Hennig EM, Suo Z, Thoresen S, Holm R, Kvinnsland S, Nesland JM (1999) Human papillomavirus 16 in breast cancer of women treated for high grade cervical intraepithelial neoplasia (CIN III). *Breast Cancer Res Treatment* 53: 121–135
- International Agency for Research on Cancer (IARC) (1995) *Monographs on the Evaluation of Carcinogenic Risks to Humans: Human Papillomaviruses* Vol 64. Lyon, France: WHO
- Kan C-Y, Iacopetta BJ, Lawson JS, Whitaker NJ (2005) Identification of human papillomavirus DNA gene sequences in human breast cancer. *Br J Cancer* 93: 946–948
- Kroupis C, Markou A, Vourlidis N, Dionysiou-Asteriou A, Lianidou ES (2006) Presence of high-risk human papillomavirus sequences in breast cancer tissues and association with histopathological characteristics. *Clin Biochem* 39: 727–731
- Lawson JS, Guenzburg W, Whitaker NJ (2006a) Viruses and breast cancer. *Future Microbiol* 1: 33–51
- Lawson JS, Kan CY, Iacopetta BJ, Whitaker NJ (2006b) Are some breast cancers sexually transmitted? *Br J Cancer* 95: 1708
- Liu Y, Klimberg VS, Andrews NR, Hicks CR, Peng H, Chiriva-Internati M, Henry-Tillman R, Hermonat PL (2001) Human papillomavirus DNA is present in a subset of unselected breast cancers. *J Hum Virol* 4: 329–334

*Correspondence: Dr A-E Al Moustafa, Lady Davis Institute for Medical Research of the Sir Mortimer B Davis-Jewish General Hospital, 3755 Cote Ste-Catherine Montreal, Quebec, Canada H3T 1E2;
E-mail: ala-eddin.almoustafa@mcgill.ca
Published online 22 January 2008

- Widschwendter A, Brunhuber T, Wiedemair A, Mueller-Holzner E, Marth C (2004) Detection of human papillomavirus DNA in breast cancer of patients with cervical cancer history. *J Clin Virol* **31**: 292–297
- Yasmeen A, Bismar TA, Kandouz M, Foulkes W, Desprez P-Y, Al Moustafa A-E (2007a) E6/E7 of HPV type 16 promotes cell invasion and metastasis of human breast cancer cells. *Cell Cycle* **6**: 2033–2042
- Yasmeen A, Ricciardi R, Kassab A, Bismar TA, Al Moustafa A-E (2007b) High-risk HPVs in human breast cancer and normal mammary tissues. *The Breast J* **16**: 445
- Yu Y, Morimoto T, Sasa M, Okazaki K, Harada Y, Fujiwara T, Irie Y, Takahashi E, Tanigami A, Izumi K (2000) Human papillomavirus type 33 DNA in breast cancer in Chinese. *Breast Cancer* **7**: 33–36