Letter to the Editor

Molecular determinants of prognosis in oesophageal cancer patients treated with chemoradiation

JA Ajani^{*,1}

¹Department of GI Medical Oncology, University of Texas MD Anderson Cancer Center, Gastroesophageal Multidisciplinary Translation Research, Houston, TX 77030, USA

British Journal of Cancer (2008) 99, 1195. doi:10.1038/sj.bjc.6604627 www.bjcancer.com
Published online 2 September 2008
© 2008 Cancer Research UK

Sir,

The paper by Yoshikawa *et al* (2008) is of considerable interest to our group. The authors assessed the expression of Gli-1 in the residual squamous cell carcinoma of the oesophagus in the resected specimens of the patients treated with preoperative chemoradiation. The expression of Gli-1 correlated with poor prognosis. When employing the strategy of preoperative chemoradiation in oesophageal carcinoma, the prognosis of the patients can be determined by assessing the degree of residual cancer (Chirieac *et al*, 2005; Wu *et al*, 2007). In addition, the residual cancer correlates with metastatic potential (Rohatgi *et al*, 2005). It would be of interest to document that Gli-1 expression has an independent prognostic value. The

authors did not report a multivariate analysis. Our group first documented the expression and temporal kinetics of sonic hedgehog ligand and Gli-1 protein expression in a xenograft model using oesophageal cancer cell lines (Sims-Mourtada *et al*, 2006). Both biomarkers correlated with chemoradiation resistance and dynamics of tumour repopulation after chemoradiation (Sims-Mourtada *et al*, 2006). Furthermore, our group documented that sonic hedgehog signalling mediates chemotherapy resistance through multiple drug transporter mechanisms and that this resistance can be overcome by inhibitors of the sonic hedgehog pathway (Sims-Mourtada *et al*, 2007). Unfortunately, Yoshikawa *et al* (2008) have neglected to discuss these two publications.

REFERENCES

- Chirieac LR, Swisher SG, Ajani JA, Komaki RR, Correa AM, Morris JS, Roth JA, Rashid A, Hamilton SR, Wu TT (2005) Posttherapy pathologic stage predicts survival in patients with esophageal carcinoma receiving preoperative chemoradiation. *Cancer* 103: 1347–1355
- Rohatgi PR, Swisher SG, Correa AM, Wu TT, Liao Z, Komaki R, Walsh G, Vaporciyan A, Lynch PM, Rice DC, Roth JA, Ajani JA (2005) Failure patterns correlate with the proportion of residual carcinoma after preoperative chemoradiotherapy for carcinoma of the esophagus. *Cancer* 104: 1349–1355
- Sims-Mourtada J, Izzo JG, Ajani J, Chao KS (2007) Sonic Hedgehog promotes multiple drug resistance by regulation of drug transport. *Oncogene* 26: 5674–5679
- Sims-Mourtada J, Izzo JG, Apisarnthanarax S, Wu TT, Malhotra U, Luthra R, Liao Z, Komaki R, van der Kogel A, Ajani J, Chao KS (2006) Hedgehog: an attribute to tumor regrowth after chemoradiotherapy and a target to improve radiation response. *Clin Cancer Res* 12: 6565–6572
- Wu TT, Chirieac LR, Abraham SC, Krasinskas AM, Wang H, Rashid A, Correa AM, Hofstetter WL, Ajani JA, Swisher SG (2007) Excellent interobserver agreement on grading the extent of residual carcinoma after preoperative chemoradiation in esophageal and esophagogastric junction carcinoma: a reliable predictor for patient outcome. *Am J Surg Pathol* 31: 58–64
- Yoshikawa R, Nakano Y, Tao L, Koishi K, Matsumoto T, Sasako M, Tsujimura T, Hashimoto-Tamaoki T, Fujiwara Y (2008) Hedgehog signal activation in oesophageal cancer patients undergoing neoadjuvant chemoradiotherapy. 98: 1670–1674

*Correspondence: Professor JA Ajani;
E-mails: jajani@mdanderson.org or jafferajani@mac.com
Published online 2 September 2008