Dietary glycemic load and gastric cancer risk in Italy

P Bertuccio^{*,1}, D Praud^{1,2}, L Chatenoud¹, E Lucenteforte¹, C Bosetti¹, C Pelucchi¹, M Rossi¹, E Negri¹ and C La Vecchia^{1,3}

¹Istituto di Ricerche Farmacologiche 'Mario Negri', Via Giuseppe La Masa, 19, 20156 Milan, Italy; ²Institut d'Informatique et de Mathématiques Appliquées, Université Joseph Fourier, BP 53, 38041 Grenoble Cedex 9, France; ³Istituto di Statistica Medica e Biometria 'Giulio A. Maccacaro', Università degli Studi di Milano, 20133 Milan, Italy

We investigated gastric cancer risk in relation to dietary glycemic index (GI) and glycemic load (GL), which represent indirect measures of carbohydrate absorption and consequently of dietary insulin demand, in a case-control study conducted in northern Italy between 1997 and 2007, including 230 patients with the incident, histologically confirmed gastric cancer and 547 frequency matched controls, admitted to the same hospitals as cases with acute non-neoplastic conditions. We used conditional logistic regression models, including terms for major recognised gastric cancer risk factors and non-carbohydrate energy intake. The odds ratios (ORs) in the highest vs lowest quintile were 1.9 (95% CI: 1.0–3.3) for GI and 2.5 (95% CI: 1.3–4.9) for GL. Compared with participants reporting low GL and high fruits/vegetables intake, the OR rose across strata of high GL and low fruits/vegetables, to reach 5.0 (95% CI: 2.2–11.5) for those reporting low fruits/vegetables intake and high GL. Our study may help to explain the direct relation observed in several studies between starchy foods and gastric cancer risk.

British Journal of Cancer (2009) 100, 558–561. doi:10.1038/sj.bjc.6604894 www.bjcancer.com

© 2009 Cancer Research UK

Keywords: case-control studies; dietary glycemic load; glycemic index; gastric cancer

A direct relation between starchy foods – particularly refined cereals – and gastric cancer has long been observed in studies conducted in Greece (Trichopoulos *et al*, 1985), Italy (La Vecchia *et al*, 1987) and Belgium (Tuyns *et al*, 1992), and has been confirmed in most subsequent studies (World Cancer Research Fund/American Institute for Cancer Research, 2007; Lucenteforte *et al*, 2008; Navarro Silvera *et al*, 2008). Refined cereals and starchy foods, particularly in southern Europe, considered indicators of a poorer diet, have been associated with increased gastric cancer risk (La Vecchia and Franceschi, 2000); they may be associated with a diet high in glycemic index (GI) and glycemic load (GL).

The GI is an indicator of the absorption rate of carbohydrates and ranks their effect on blood glucose concentrations. It compares available carbohydrates gram-for-gram in individual foods, providing a numerical, evidence-based index of postprandial glycemia (Jenkins *et al*, 1981, 1984; Gnagnarella *et al*, 2004). As the GL combines measures of carbohydrate (both qualitative and quantitative) and of dietary insulin demand, the overall GI reflects the average quality of carbohydrates consumed, whereas the total dietary GL reflects both their average quantity and quality (Foster-Powell *et al*, 2002).

GI and, particularly, GL have been related to excess risk of colorectal (Augustin *et al*, 2002), breast (Augustin *et al*, 2001), oral and oesophageal (Augustin *et al*, 2003) cancers. Gastric cancer has also been associated with GI and GL in a case-control study in Italy (Augustin *et al*, 2004), though not in a Swedish cohort study

(Larsson *et al*, 2006). We have further considered the relation between GL, GI and gastric cancer risk using data from another Italian case-control study, based on a more detailed, reproducible and validated food frequency questionnaire (FFQ) (Lucenteforte *et al*, 2008).

MATERIALS AND METHODS

We derived data from a case-control study of gastric cancer conducted in 1997–2007 in the greater Milan area, Italy, the design of which has been described earlier (Lucenteforte *et al*, 2008). Briefly, cases were 230 patients (143 men, 87 women; median age 63 years, range 22–80 years) admitted to major teaching and general hospitals in the study area with the incident, histologically confirmed stomach cancer (ICD IX 151.0–151.9), diagnosed no longer than 1 year before the interview, and with no earlier diagnosis of cancer. The control group included 547 patients (286 men, 261 women; median age 63 years, range 22–80 years) frequency matched to cases by age and sex (ratio of 2:1 for men and 3:1 for women), admitted to the same hospitals as cases for a wide spectrum of acute, non-neoplastic conditions, unrelated to risk factors for stomach cancer and long-term diet modification. Of these 20% were admitted for traumas, 23% for other orthopaedic conditions, 22% for acute surgical and 35% for other miscellaneous disorders. Less than 5% of cases and controls who were approached refused to be interviewed.

For both cases and controls, data were collected during their hospital stay by trained interviewers using a structured questionnaire covering socio-demographic characteristics,

*Correspondence: Dr P Bertuccio; E-mail: bertuccio@marionegri.it

Received 11 November 2008; revised 19 December 2008; accepted 19 December 2008

anthropometric measures, selected lifestyle habits, including tobacco and alcohol consumption, personal medical and family history of cancer. The patients' usual diet during the two years before diagnosis or hospital admission (for controls) was assessed through a reproducible (Franceschi *et al*, 1995) and valid (Decarli *et al*, 1996) FFQ covering 78 foods and beverages, including a range of the most common recipes in Italian diet. Participants indicated the average weekly frequency of consumption for each dietary item; intakes lower than once a week but at least once a month were coded as 0.5 per week. Energy and carbohydrate intake were computed from the FFQ using an Italian food composition database, integrated with other sources when needed (Salvini *et al*, 1998; Gnagnarella *et al*, 2004).

GI values were assigned to each food item using international tables (Foster-Powell *et al*, 2002). Daily average GI was calculated by summing the products of the carbohydrate content per serving for each food or food group, times the average number of servings of that food per week, times its GI (Wolever *et al*, 1994), all divided by the total amount of available carbohydrates. The daily average GL was computed as the GI, but without dividing by the total amount of carbohydrates.

Odds ratios (ORs) and their corresponding 95% confidence intervals (CIs) for subsequent quintiles of GI and GL were estimated using conditional multiple logistic regression models, conditioned on age and sex (Breslow and Day, 1980). We considered two models: the first included terms for period of interview, education, body mass index, tobacco smoking, intake of fruits and vegetables, and family history of stomach cancer; the second also included non-carbohydrate energy intake, to allow for bias due to systematic over- or under-reporting (Willett and Stampfer, 1986).

RESULTS

Table 1 gives the distribution of gastric cancer cases and controls according to quintiles of GI and GL, and the corresponding multivariate ORs. The OR for the highest vs the lowest quintile of GI was 2.1 (95% CI: 1.2–3.6), with a significant trend in risk ($P = 0.034$). For GL, ORs were significantly above unity in the third (OR = 2.5, 95% CI: 1.4–4.5), fourth (OR = 2.7, 95% CI: 1.5–4.9) and highest quintile (OR = 2.7; 95% CI: 1.5–4.8), with a significant

trend in risk ($P < 0.001$). Continuous ORs for the difference between the 80th and 20th percentile (based on the control distribution) were 1.4 (95% CI: 1.1–1.9) for GI and 1.5 (95% CI: 1.2–2.0) for GL. After further adjustment for non-carbohydrate energy intake, the OR estimates were similar for both GI and GL.

Table 2 gives the continuous ORs of gastric cancer for GI and GL in strata of sex, age and major established risk factors for gastric cancer. No significant differences emerged in any strata of the covariates considered.

We further considered the combined effect of fruits/vegetables intake and GL on gastric cancer risk. Compared with participants reporting low GL and high fruits and vegetables intake, the OR rose for increasing levels of GL, particularly in participants with low fruits/vegetables intake, to reach 5.0 (95% CI: 2.2–11.5) for those reporting high GL and low fruits/vegetables intake.

DISCUSSION

This study, based on a valid and detailed FFQ, confirms and further quantifies the existence of a direct relation between GI, and mostly GL, and gastric cancer risk. We were able to allow in the analysis not only for major risk factors for gastric cancer, such as education, tobacco smoking and vegetables and fruits intake, but also for non-carbohydrate energy intake. Our results indicate that some characteristics of carbohydrates, in particular of such refined carbohydrates as white bread, pasta and rice, have an appreciable influence on gastric cancer risk in this population.

The findings are consistent with those of an earlier Italian case-control study, based on a shorter questionnaire, which found an approximately two-fold risk of gastric cancer for the highest quartile of GL, and only a modest association with GI (Augustin *et al*, 2004). They also agree with several earlier studies, indicating that a diet rich in cereals or starches is associated with an increased gastric cancer risk (Trichopoulos *et al*, 1985; La Vecchia *et al*, 1987; Tuyns *et al*, 1992; World Cancer Research Fund/American Institute for Cancer Research, 2007; Lucenteforte *et al*, 2008; Navarro Silvera *et al*, 2008). They are, however, at variance with the findings of the Swedish Mammography Cohort (Larsson *et al*, 2006), which found no relation between GI and GL and gastric cancer risk.

Table 1 Odds ratios (ORs) and 95% confidence interval (CI) of gastric cancer among 230 cases and 547 controls, according to glycemic index and glycemic load. Italy, 1997–2007

	Upper limit	Cases n (%)	Controls n (%)	OR ^a (95% CI)	OR ^b (95% CI)
Glycemic index					
I (low)	73	28 (12.2)	110 (20.1)	1.0 ^c	1.0 ^c
II	77	44 (19.1)	109 (19.9)	1.5 (0.9–2.7)	1.5 (0.9–2.6)
III	80	44 (19.1)	110 (20.1)	1.5 (0.8–2.6)	1.4 (0.8–2.4)
IV	83	47 (20.4)	108 (19.7)	1.5 (0.8–2.6)	1.4 (0.8–2.5)
V (high)	—	67 (29.1)	110 (20.1)	2.1 (1.2–3.6)	1.9 (1.0–3.3)
χ^2_1 trend (P-value)				4.5 (P-value = 0.034)	3.0 (P-value = 0.083)
Continuous ^d				1.4 (1.1–1.9)	1.4 (1.0–1.9)
Glycemic load					
I (low)	147	24 (10.4)	109 (19.9)	1.0 ^c	1.0 ^c
II	183	30 (13.0)	110 (20.1)	1.4 (0.7–2.5)	1.3 (0.7–2.5)
III	217	57 (24.8)	110 (20.1)	2.5 (1.4–4.5)	2.4 (1.3–4.5)
IV	263	58 (25.2)	108 (19.7)	2.7 (1.5–4.9)	2.6 (1.4–5.1)
V (high)	—	61 (26.5)	110 (20.1)	2.7 (1.5–4.8)	2.5 (1.3–4.9)
χ^2_1 trend (P-value)				13.4 (P-value < 0.001)	8.9 (P-value = 0.003)
Continuous ^d				1.5 (1.2–2.0)	1.4 (1.1–1.9)

^aEstimates from conditional logistic regression models, conditioned on age and sex and adjusted for period of interview, only when indicated, education, body mass index, tobacco smoking, intake of fruits and vegetables, and family history of stomach cancer. ^bFurther adjusted for non-carbohydrate energy intake. ^cReference category. ^dThe unit is the difference between the 80th and the 20th percentile.

Table 2 Odds ratios (ORs) and 95% confidence intervals (CI) of gastric cancer among 230 cases and 547 controls, according to glycemic index and glycemic load in strata of selected covariates. Italy, 1997–2007

Covariates (no. of cases/no. of controls)	OR (95 % CI) ^{a,b}	
	Glycemic index	Glycemic load
Sex		
Men (143/286)	1.1 (0.7–1.6)	1.5 (1.0–2.1)
Women (87/261)	1.8 (1.1–2.9)	1.3 (0.8–2.3)
Age (years)		
< 65 (125/297)	1.5 (1.0–2.3)	1.3 (0.9–2.0)
≥ 65 (105/250)	1.3 (0.8–2.1)	1.7 (1.1–2.7)
Education ^c (years)		
< 7 (95/236)	1.7 (1.0–3.0)	1.4 (0.9–2.4)
≥ 7 (132/307)	1.1 (0.8–1.6)	1.4 (0.9–2.0)
Body mass index ^c (kg m ⁻²)		
< 25 (118/248)	1.2 (0.8–1.8)	1.3 (0.8–2.0)
≥ 25 (108/297)	1.6 (1.0–2.6)	1.6 (1.0–2.4)
Smoking status ^c		
Non-smoker (171/428)	1.5 (1.0–2.2)	1.3 (0.9–1.8)
Smoker (58/118)	1.2 (0.6–2.3)	2.1 (1.1–3.9)
Intake of fruits (portions/week)		
≤ 15 (117/273)	1.5 (1.0–2.3)	1.5 (1.0–2.2)
> 15 (113/274)	1.6 (1.0–2.5)	1.4 (0.9–2.2)
Intake of vegetables (portions/week)		
≤ 9 (141/274)	1.4 (0.9–2.1)	1.3 (0.9–2.0)
> 9 (89/273)	1.2 (0.8–1.9)	1.4 (0.9–2.2)

^aThe unit is the difference between the 80th and the 20th percentile. ^bEstimates from conditional logistic regression models, conditioned on age and sex and adjusted for period of interview, only when indicated, education, body mass index, tobacco smoking, intake of fruits and vegetables, family history of stomach cancer and non-carbohydrate energy intake. ^cThe sum does not add up to the total because of some missing values.

We also found that the role of GL is independent of other major dietary correlates for gastric cancer, including vegetables and fruits consumption, and that there was a five-fold difference in gastric cancer risk between participants reporting low GL and high fruits and vegetables intake and those reporting high GL and low fruits and vegetables consumption. Thus, two simple diet indicators determine substantial variation in gastric cancer risk.

A high GL diet may increase gastric cancer risk through the modulation of the insulin-like growth factors (IGFs). Insulin increases the activity of IGFs, such as IGF-1 (Gnagnarella *et al*, 2008), and suppresses hepatic secretion of IGF-binding protein-1. IGF-1 stimulates cell proliferation and differentiation, inhibits apoptosis (Kaaks and Lukanova, 2001; Gnagnarella *et al*, 2008) and increases production of vascular endothelial growth factors, important in tumour angiogenesis, and the expression of glucose

transporters, glycolytic enzymes, and growth factors, which may promote tumor cell survival under hypoxic conditions (Akakura *et al*, 2001; Oh *et al*, 2008). Higher IGF-1 concentrations have been observed in patients with gastric cancer than in healthy controls (Franciosi *et al*, 2003). Insulin can also influence sex hormone concentrations and reduce the levels of their binding protein (Garzo and Dorrington, 1984; Poretsky and Kalin, 1987; Gnagnarella *et al*, 2008). Sex hormones have been linked to gastric cancer risk (La Vecchia *et al*, 1994), though this has been disputed (Bahmanyar *et al*, 2008; Persson *et al*, 2008).

A limitation of our study is the lack of information on *Helicobacter pylori* (*H. pylori*) infection. A recent study observed a positive association between plasma glucose concentration and risk of stomach cancer only among *H. pylori*-positive participants (Yamagata *et al*, 2005; Larsson *et al*, 2006). Though the prevalence of *H. pylori* infection is declining (Malaty, 2007), it was relatively high (about 45%) in Italy in the mid 1990's. It increased with age and was more frequent in men than in women (Russo *et al*, 1999). Thus, a large proportion of the population studied is likely to be *H. Pylori*-positive.

Participation among eligible patients was almost complete, and the catchment areas for cases and controls were comparable. Hospital controls may differ in their dietary habits from the general population. However, we excluded control participants admitted for conditions associated with dietary modifications (diabetes mellitus, cardiovascular diseases, etc.). Recall bias is also unlikely, given that the association between various types of starches and gastric cancer was not evident at the time of interviews. Moreover, the similar hospital setting for cases and controls likely increased the comparability of dietary histories, and the questionnaire was reproducible and valid (Franceschi *et al*, 1995; Decarli *et al*, 1996; D'Avanzo *et al*, 1997).

With respect to confounding results, we adjusted for non-carbohydrate energy intake to control for potential systematic over- or under-reporting between cases and controls (Willett and Stampfer, 1986); further adjustment for several covariates failed to explain the association between GI and GL and gastric cancer.

Our study suggests that a diet high in GL and GI is directly related to gastric cancer risk, and may therefore help to explain the relation observed in several studies with starchy foods (Trichopoulos *et al*, 1985; La Vecchia *et al*, 1987; Buiatti *et al*, 1990; Tuyns *et al*, 1992).

ACKNOWLEDGEMENTS

This work was conducted with contribution from the Italian Association for Cancer Research (AIRC) and the Italian League Against Cancer. PB was supported by a fellowship from the Italian Foundation for Cancer Research (FIRC). The work of this paper was undertaken while CLV was a senior fellow at the International Agency for Research on Cancer, Lyon, France. The authors thank Ms I Garimoldi for editorial assistance.

REFERENCES

- Akakura N, Kobayashi M, Horiuchi I, Suzuki A, Wang J, Chen J, Niizeki H, Kawamura K, Hosokawa M, Asaka M (2001) Constitutive expression of hypoxia-inducible factor-1 α renders pancreatic cancer cells resistant to apoptosis induced by hypoxia and nutrient deprivation. *Cancer Res* **61**: 6548–6554
- Augustin LS, Dal Maso L, La Vecchia C, Parpinel M, Negri E, Vaccarella S, Kendall CW, Jenkins DJ, Franceschi S (2001) Dietary glycemic index and glycemic load, and breast cancer risk: a case-control study. *Ann Oncol* **12**: 1533–1538
- Augustin LS, Franceschi S, Jenkins DJ, Kendall CW, La Vecchia C (2002) Glycemic index in chronic disease: a review. *Eur J Clin Nutr* **56**: 1049–1071
- Augustin LS, Gallus S, Franceschi S, Negri E, Jenkins DJ, Kendall CW, Dal Maso L, Talamini R, La Vecchia C (2003) Glycemic index and load and risk of upper aero-digestive tract neoplasms (Italy). *Cancer Causes Control* **14**: 657–662
- Augustin LS, Gallus S, Negri E, La Vecchia C (2004) Glycemic index, glycemic load and risk of gastric cancer. *Ann Oncol* **15**: 581–584

- Bahmanyar S, Lambe M, Zendehdel K, Nyren O, Boffetta P, Ye W (2008) Parity and risk of stomach cancer by sub-site: a national Swedish study. *Br J Cancer* **98**: 1295–1300
- Breslow NE, Day NE (1980) *Statistical Methods in Cancer Research. Vol. I. The Analysis of Case-Control Studies*. IARC Sci Publ No. 32. IARC: Lyon, France
- Buiatti E, Palli D, Decarli A, Amadori D, Avellini C, Bianchi S, Bonaguri C, Cipriani F, Cocco P, Giacosa A, Marubini E, Minacci C, Puntoni R, Russo A, Vindigni G, Fraumeni Jr JF, Blot WJ (1990) A case-control study of gastric cancer and diet in Italy: II. Association with nutrients. *Int J Cancer* **45**: 896–901
- D'Avanzo B, La Vecchia C, Katsouyanni K, Negri E, Trichopoulos D (1997) An assessment, and reproducibility of food frequency data provided by hospital controls. *Eur J Cancer Prev* **6**: 288–293
- Decarli A, Franceschi S, Ferraroni M, Gnagnarella P, Parpinel MT, La Vecchia C, Negri E, Salvini S, Falcini F, Giacosa A (1996) Validation of a food-frequency questionnaire to assess dietary intakes in cancer studies in Italy. Results for specific nutrients. *Ann Epidemiol* **6**: 110–118
- Foster-Powell K, Holt SH, Brand-Miller JC (2002) International table of glycemic index and glycemic load values: 2002. *Am J Clin Nutr* **76**: 5–56
- Franceschi S, Barbone F, Negri E, Decarli A, Ferraroni M, Filiberti R, Giacosa A, Gnagnarella P, Nanni O, Salvini S, La Vecchia C (1995) Reproducibility of an Italian food frequency questionnaire for cancer studies. Results for specific nutrients. *Ann Epidemiol* **5**: 69–75
- Franciosi CM, Piacentini MG, Conti M, Romano F, Musco F, Caprotti R, Rovelli F, Uggeri F (2003) IGF-1 and IGF-1BP3 in gastric adenocarcinoma. Preliminary study. *Hepatogastroenterology* **50**: 297–300
- Garzo VG, Dorrington JH (1984) Aromatase activity in human granulosa cells during follicular development and the modulation by follicle-stimulating hormone and insulin. *Am J Obstet Gynecol* **148**: 657–662
- Gnagnarella P, Gandini S, La Vecchia C, Maisonneuve P (2008) Glycemic index, glycemic load, and cancer risk: a meta-analysis. *Am J Clin Nutr* **87**: 1793–1801
- Gnagnarella P, Parpinel M, Salvini S, Franceschi S, Palli D, Boyle P (2004) The update of the Italian food composition database. *J Food Comp Analysis* **17**: 509–522
- Jenkins DJ, Wolever TM, Jenkins AL, Josse RG, Wong GS (1984) The glycaemic response to carbohydrate foods. *Lancet* **2**: 388–391
- Jenkins DJ, Wolever TM, Taylor RH, Barker H, Fielden H, Baldwin JM, Bowling AC, Newman HC, Jenkins AL, Goff DV (1981) Glycemic index of foods: a physiological basis for carbohydrate exchange. *Am J Clin Nutr* **34**: 362–366
- Kaaks R, Lukanova A (2001) Energy balance and cancer: the role of insulin and insulin-like growth factor-I. *Proc Nutr Soc* **60**: 91–106
- La Vecchia C, D'Avanzo B, Franceschi S, Negri E, Parazzini F, Decarli A (1994) Menstrual and reproductive factors and gastric-cancer risk in women. *Int J Cancer* **59**: 761–764
- La Vecchia C, Franceschi S (2000) Nutrition and gastric cancer with a focus on Europe. *Eur J Cancer Prev* **9**: 291–295
- La Vecchia C, Negri E, Decarli A, D'Avanzo B, Franceschi S (1987) A case-control study of diet and gastric cancer in northern Italy. *Int J Cancer* **40**: 484–489
- Larsson SC, Bergkvist L, Wolk A (2006) Glycemic load, glycemic index and carbohydrate intake in relation to risk of stomach cancer: a prospective study. *Int J Cancer* **118**: 3167–3169
- Lucenteforte E, Scita V, Bosetti C, Bertuccio P, Negri E, La Vecchia C (2008) Food groups and alcoholic beverages and the risk of stomach cancer: a case-control study in Italy. *Nutr Cancer* **60**: 577–584
- Malaty HM (2007) Epidemiology of *Helicobacter pylori* infection. *Best Pract Res Clin Gastroenterol* **21**: 205–214
- Navarro Silvera SA, Mayne ST, Risch H, Gammon MD, Vaughan TL, Chow WH, Dubrow R, Schoenberg JB, Stanford JL, West AB, Rotterdam H, Blot WJ, Fraumeni Jr JF (2008) Food group intake and risk of subtypes of esophageal and gastric cancer. *Int J Cancer* **123**: 852–860
- Oh SY, Kwon SH, Kim SH, Jang JS, Kim MC, Kim KH, Han JY, Kim CO, Kim SJ, Jeong JS, Kim HJ (2008) Clinicopathologic significance of HIF-1 α , p53, and VEGF expression and preoperative serum VEGF level in gastric cancer. *BMC Cancer* **8**: 123
- Persson C, Inoue M, Sasazuki S, Kurahashi N, Iwasaki M, Ye W, Tsugane S (2008) Female reproductive factors and the risk of gastric cancer in a large-scale population-based cohort study in Japan (JPHC study). *Eur J Cancer Prev* **17**: 345–353
- Poretsky L, Kalin MF (1987) The gonadotropic function of insulin. *Endocr Rev* **8**: 132–141
- Russo A, Eboli M, Pizzetti P, Di Felice G, Ravagnani F, Spinelli P, Hotz AM, Notti P, Maconi G, Franceschi S, Ferrari D, Bertario L (1999) Determinants of *Helicobacter pylori* seroprevalence among Italian blood donors. *Eur J Gastroenterol Hepatol* **11**: 867–873
- Salvini S, Parpinel MT, Gnagnarella P, Maisonneuve P, Turrini A (1998) *Banca dati di composizione degli alimenti per studi epidemiologici in Italia*. Istituto Europeo di Oncologia: Milano
- Trichopoulos D, Ouranos G, Day NE, Tzonou A, Manousos O, Papadimitriou C, Trichopoulou A (1985) Diet and cancer of the stomach: a case-control study in Greece. *Int J Cancer* **36**: 291–297
- Tuyns AJ, Kaaks R, Haelterman M, Riboli E (1992) Diet and gastric cancer. A case-control study in Belgium. *Int J Cancer* **51**: 1–6
- Willett W, Stampfer MJ (1986) Total energy intake: implications for epidemiologic analyses. *Am J Epidemiol* **124**: 17–27
- Wolever TM, Nguyen PM, Chiasson JL, Hunt JA, Josse RG, Palmason C, Rodger NW, Ross SA, Ryan EA, Tan MH (1994) Determinants of diet glycemic index calculated retrospectively from diet records of 342 individuals with non-insulin-dependent diabetes mellitus. *Am J Clin Nutr* **59**: 1265–1269
- World Cancer Research Fund/American Institute for Cancer Research (2007) *Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Prospective*. American Institute for Cancer Research: Washington, DC
- Yamagata H, Kiyohara Y, Nakamura S, Kubo M, Tanizaki Y, Matsumoto T, Tanaka K, Kato I, Shirota T, Iida M (2005) Impact of fasting plasma glucose levels on gastric cancer incidence in a general Japanese population: the Hisayama study. *Diabetes Care* **28**: 789–794