microangiopathy, particularly in the presence of diabetes, and with compressive lesions caused by intracranial aneurysms. They are a rare presenting sign in bilateral subdural haematoma. A review of 114 cases of bilateral subdural haematoma identified no cases in which the presentation involved an oculomotor nerve palsy.¹ Two cases have been published describing oculomotor nerve palsies in association with bilateral subdural haematoma. One describes a case similar to that reported here in which CT investigation of a patient presenting with an isolated oculomotor nerve palsy revealed bilateral chronic subdural haematoma, which also resolved following surgical evacuation.² The other describes a case of oculomotor nerve palsy and hemiparesis complicating the surgical drainage of subdural haematoma.³

Subdural haematoma rarely affects the anterior visual pathways, more commonly affecting the posterior pathways in the form of homonymous hemianopia. However, this case involved in addition to the oculomotor nerve palsy, an ipsilateral reduction in visual acuity, which subsequently resolved upon drainage of the haematoma. Both signs may have arisen by the same mechanism. This patient's CT brain revealed bilateral subdural haematoma causing downward displacement of the cerebral hemispheres with herniation of the uncus portion of the temporal lobe. This in turn is likely to have led to compression and paresis of the oculomotor nerve. The downward displacement may have led to traction of the optic nerve, which in this case, with a history of ischaemic optic neuropathy, is known to be vascularly compromised, resulting in reversible reduction in visual acuity. Following evacuation of the haematoma, with release of the traction on the optic nerve, normal visual acuity was restored. Hollander and Stewart⁴ describe a similar case in which sudden painless visual loss occurred in the presence of subdural haematoma with partial resolution following drainage. They similarly speculate that midline shift caused by subdural haematoma led to optic nerve traction and vascular compromise that was not detectable on imaging.

The patient in this case also described formed visual hallucinations occurring upon closing his eyes. Manford and Andermann⁵ describe three mechanisms by which complex visual hallucinations may occur: epileptic hallucinations due to direct irritation of cortical centres; visual pathway lesions resulting in defective visual processing with abnormal cortical release phenomena (Charles-Bonnet hallucinations) and disturbance of ascending cholinergic and serotonergic pathways in brainstem lesions. In this case the visual hallucinations described were most likely caused by direct stimulation of cortical centres by the

presence of the subdural haematoma. Charles-Bonnet hallucinations are less likely, given the relatively short duration of this patient's visual loss and the fact that these hallucinations were present only upon closing his eyes.

In summary, this case represents an unusual presentation of bilateral subacute subdural haematoma with oculomotor palsy, visual loss, and hallucinations.

References

- 1 Cameron MM. Chronic subdural haematoma: a review of 14 cases. *J Neurol Neurosurg Psychiatr* 1978; **41**(9): 834–839.
- 2 Phookan G, Cameron M. Bilateral chronic subdural haematoma: an unusual presentation with isolated oculomotor nerve palsy. *J Neurol Neurosurg Psychiatr* 1994; **57**(9): 1146.
- 3 Okuchi K, Fujioka M, Maeda Y, Kagoshima T, Sakaki T. Bilateral chronic subdural haematomas resulting in unilateral oculomotor nerve paresis and brain stem symptoms after operation—case report. *Neurologica Medico-Chirurgica* 1999; **39**(5): 365–371.
- 4 Hollander DA, Stewart MJ. Anterior pathway vision loss due to subdural haematoma. *Br J Ophthalmol* 2003; **87**(11): 1423–1424.
- 5 Manford M, Andermann F. Complex visual hallucinations—clinical and neurobiological insights. *Brain* 1998; **121**: 1819–1840.

C Mulholland and FA Knox

Ward 28, Department of Ophthalmology,
The Royal Victoria Hospital, Grosvenor Road,
Belfast, BT12 6BA, UK

Correspondence: C Mulholland

Tel: +44 28 90632392;

Fax: +44 28 90330744.

E-mails: conormulholland@doctors.org.uk;

angelaknox@ireland.com

This work has not previously been presented at any meeting

Eye (2006) **20**, 125–126. doi:10.1038/sj.eye.6701809;
published online 21 January 2005

Sir,

Orbital well-differentiated liposarcoma demonstrating chromosomal imbalances

Although adipose tissue comprises a large portion of the orbit, liposarcomas consisted of only 2% of primary

orbital tumours, and were hardly ever included in the clinical differential diagnosis of orbital lesions.¹

We report a 24-year-old woman who suffered from recurrent orbital liposarcoma with unusual findings. She first presented with right 3 mm exophthalmos and limitation of eye movements that developed within 2 weeks. Her visual acuity was 6/6. CT scan and MRI revealed an extraconal lesion, consisting of fat, soft tissue, and calcifications at the roof of the right orbit, pushing down the superior rectus. The patient postponed surgery for 5 months and the proptosis progressed to 6 mm. She underwent anterior and lateral orbitotomy with incomplete debulking of the tumour found between the levator and the superior rectus muscles. After surgery, she felt well for 8 months, when proptosis recurred (Figure 1a) with complete limitation of upward gaze. The visual acuity was 6/6. CT and MRI revealed a heterogenic tissue filling the right upper orbit (Figure 1b). Another operation was performed 16 months after the first one and the tumour was excised through a lid-crease incision, using a cryo-probe. At follow-up 1.5 years later, ptosis and hypoglobus persisted, but with no signs of tumour recurrence.

Paraffin sections from the primary tumour revealed a lipoma-like well-differentiated liposarcoma (WDL) composed of adipose lobules of mature adipocytes with variations in cell size and occasional cells with hyperchromatic nuclei, and scattered bizarre, multinucleated cells. Scattered lipoblasts were identified within fibrotic septa. A focus of metaplastic bone was present (Figure 1c). The relapsing lesion was an

inflammatory WDL with lipoma-like and sclerosing areas. It contained a prominent chronic inflammatory component with the formation of reactive lymphoid follicles (Figure 1d). The lipomatous component contained adipocytes varying in size, with focally hyperchromatic nuclei and scattered multivacuolar cells. The sclerosing component was paucicellular with spindle fibroblastic cells and occasional atypical cells with hyperchromatic nuclei. Immunohistochemical examination showed that the lipocytes, some of the lipoblast-like cells and some of the spindle cells were positive for S-100 protein and negative for alpha smooth muscle actin, desmin, and CD68. Immunostaining for CD3, CD20, CD68, and CD138 demonstrated a reactive immunoarchitecture of the lymphoid component.

Comparative genomic hybridization (CGH) analysis that was performed, as far as we are aware, for the first time for an orbital liposarcoma, detected same chromosomal aberrations in both the primary and relapsing tumors. Changes in the DNA sequence copy number were: +1p34–q32, +2p21–q33, +4q22–q28, +5p13–q31, +6q14–23, +8q, +12q12–q22, +13q12–q22 and –Xq. All aberrations were of a low level.

Discussion

While rare, the prognosis of orbital liposarcomas is not known. WDL outside the retroperitoneum is usually cured by wide resection with a 20–30% risk of local recurrence and up to 5% risk of dedifferentiation and metastasis.^{2,3}

CGH might help indicate more malignant phenotypes.^{4–6} While gains within chromosomes 1,4,5,6,8, 12, and 13 were found in WDL, gains within chromosomes 2 have not been reported before. Loss of chromosomal material, such as the loss of Xq found in our case, is generally detected less frequently in sarcomas than gains.⁴ It was suggested that an increased number of aberrations, over-representations of chromosomal subregion 12q24, and amplification of 1q may indicate a more malignant phenotype.^{4,6}

The commonly accepted classification of liposarcoma by the World Health Organization⁷ distinguishes between well-differentiated, myxoid, round cell, pleomorphic, and dedifferentiated groups. WDL is further classified into lipoma-like, sclerosing, and inflammatory subgroups.^{3,7} Unusually, metaplasia into cartilaginous, osseous, and smooth muscle elements may present in liposarcoma.⁸ However, metaplastic bone in orbital liposarcoma, to the best of our knowledge, has not been previously described. Inflammatory WDL is rare and almost always retroperitoneal. Inflammatory WDL,

Figure 1 (a) Recurrent orbital liposarcoma causes fullness of the right upper eyelid, ptosis, and hypoglobus. (b) CT scans of the recurring tumour. Heterogenic tissue fills the right orbit at the upper retrobulbar space. (c) A focus of metaplastic bone was seen in the primary lipoma-like liposarcoma (asterisk) (H&E; original magnification $\times 40$). (d) Formations of reactive lymphoid follicles (arrows) in the relapsing, inflammatory WDL (H&E; original magnification, $\times 40$). (The images were digitally processed.)

as found in the recurrent lesion of our patient, has been described only once as an orbital primary tumour³ and has not been previously reported as a recurrent orbital liposarcoma.

A recent classification from The Armed Forces Institute of Pathology⁸ groups together the WDL, atypical lipoma, and pleomorphic lipoma into one category of 'atypical lipomatous tumor' located between the benign and malignant groups, as it contains cells with irregular, hyperchromatic nuclei but does not metastasize. It was also argued that the differentiation between lipoma-like, sclerosing, and inflammatory subgroups have no significance and may occur in the same tumour. Our finding of the same chromosomal imbalances in the primary lipoma-like tumour and in the inflammatory relapsing tumour also suggests that these subtypes should be grouped together.

In summary, liposarcoma with metaplastic bone and recurrent liposarcoma with inflammatory component should be added to the differential diagnosis of rare orbital lesions.

Acknowledgements

Financial support/Proprietary interest: None.

References

- Jakobiec FA, Rini F, Char D, Orcutt J, Rootman J, Baylis H *et al*. Primary liposarcoma of the orbit. Problems in the diagnosis and management of five cases. *Ophthalmology* 1989; **96**: 180–191.
- Kraus MD, Guillou L, Fletcher CDM. Well-differentiated inflammatory liposarcoma is an uncommon and easily overlooked variant of common sarcoma. *Am J Surg Pathol* 1997; **21**: 518–527.
- Cai YC, McMenamin ME, Rose G, Sandy CJ, Cree IA, Fletcher CDM. Primary liposarcoma of the orbit: a clinicopathologic study of seven cases. *Ann Diagn Pathol* 2001; **5**: 255–266.
- Rieker RJ, Joos S, Bartsch C, Willeke F, Schwarzbach M, Otano-Joos M *et al*. Distinct chromosomal imbalances in pleomorphic and in high-grade dedifferentiated liposarcomas. *Int J Cancer* 2002; **99**: 68–73.
- Szymanska J, Virolainen M, Tarkkanen M, Wiklund T, Asko-Seljavaara S, Tukiainen E *et al*. Overrepresentation of 1q21–23 and 12q13–21 in lipoma like liposarcomas but not in benign lipomas: a comparative genomic hybridization study. *Cancer Genet Cytogenet* 1997; **99**: 14–18.
- Forus A, Larramendy ML, Meza-Zepeda LA, Bjerkehagen B, Godager LH, Dahlberg AB *et al*. Dedifferentiation of a well-differentiated liposarcoma to a highly malignant metastatic osteosarcoma: amplification of 12q14 at all stages and gain of 1q22–q24 associated with metastases. *Cancer Genet Cytogenet* 2001; **125**: 100–111.
- Weiss SW, Sobin LH. *International Histological Classification of Tumors. Histological Typing of Soft Tissue Tumors*, 2nd ed. Springer: Berlin, 1994.
- Kempson RL, Fletcher CDM, Evans HL, Hendrickson MR, Sibley RK. *Tumors of the soft tissues*. Armed Forces Institute of Pathology: Washington DC, 2001, pp. 188–217.

M Rosner¹, A Yosepovich², M Paul³, N Rosen¹ and M Perelman²

¹Goldschleger Eye Institute, Tel-Aviv University, Sackler School of Medicine, Sheba Medical Center, Tel-Hashomer, Israel

²Pathologic Institute, Tel-Aviv University, Sackler School of Medicine, Sheba Medical Center Tel-Hashomer, Israel

³Eye Department, Edith Wolfson Medical Center, Holon, Israel

Correspondence: M Rosner, Goldschleger Eye Research Institute, Sheba Medical Center, 52621, Tel-Hashomer, Israel
Tel: +972 3 5302957;
Fax: +972 3 5351577.
E-mail: mrosner@post.tau.ac.il

Part of this work was presented at the Joint European Research Meeting in Ophthalmology and Vision (EVER), 2002, and at the Stein Update Seminar, 2003

Eye (2006) **20**, 126–128. doi:10.1038/sj.eye.6701812;
published online 28 January 2005

Sir, Simultaneous bilateral acute angle closure glaucoma following venlafaxine treatment

A 49-year-old woman presented to eye casualty with blurred vision and pain in both eyes. She had been diagnosed with irritable bowel syndrome (IBS) 4 years earlier and her gastroenterologist had prescribed venlafaxine earlier that day. At 4 h after taking the first oral dose, she developed symptoms of pain and blurred vision simultaneously in both eyes. She also reported nausea and vomiting.

On examination, visual acuities were 6/18 in the right eye and 6/12 in the left. Both pupils were fixed and mid-dilated. There was bilateral corneal oedema and shallow anterior chambers were evident. Gonioscopy revealed angle closure in both eyes. Intraocular pressures (IOPs) were 49 mmHg in the right eye and 54 mmHg in the left. A diagnosis of bilateral acute angle closure glaucoma was made.