

Quality of life and visual rehabilitation: an observational study of low vision in three general practices in West Glamorgan

GP Williams, V Pathak-Ray, MW Austin, AP Lloyd, IM Millington and A Bennett

Correction to: Eye (2006) E-pub ahead of print: February 3 2006; doi:10.1038/sj.eye.6702256

Eye (2007) 21, 575. doi:10.1038/sj.eye.6702537; published online 11 August 2006

In the above article, the reference in Table 5 is incorrect. The correct table is shown below:

Table 5 Comparison of mean NEI-VFQ subscale scores; present study, depressed AMD, and diabetic retinopathy sufferers

Questionnaire category	Percentage mean score for blind partially sighted in this study (SD)	Percentage mean score for depressed adults with AMD (SD) ^a	Percentage mean score for type 1 diabetes mellitus (SD) ^b
General health	57.16 (24.65)	59.08 (19.83)	60.5 (23.8)
General vision	27.80 (11.93)	36.33 (17.66)	79.7 (16.1)
Ocular pain	83.90 (23.08)	79.59 (20.99)	92.6 (87.6)
Near activities	17.58 (15.31)	23.52 (11.45)	87.6 (17.3)
Distance activities	23.30 (19.48)	32.56 (18.66)	86.6 (17.7)
Social functioning	45.52 (30.70)	54.34 (26.83)	94.7 (12.4)
Mental health	38.11 (22.87)	46.07 (20.58)	84.6 (18.2)
Role difficulties	37.50 (23.23)	34.06 (15.58)	91.8 (18.1)
Dependency	37.50 (25.66)	45.66 (25.2)	93.3 (16.5)
Driving	0	63.89 (31.55)	85.1 (22.9)
Colour vision	55.68 (35.30)	68.37 (30.52)	94.8 (14.9)
Peripheral vision	32.95 (26.37)	61.98 (28.71)	89.3 (20.9)

^aFrom Brody *et al.*¹⁶

^bFrom Klein *et al.*⁶

The authors would like to apologise for this mistake.