

commonly associated with it. *Candida* has also been found to be a cause of post-surgical ICK.³ This is the first reported case of non-tuberculous mycobacteria without iatrogenic or accidental corneal trauma. Similarly,⁴ we found fourth generation fluoroquinolones effective in treatment and recommend first line use for these cases. As *M. abscessus* can be found in soil and water, patients at risk of infection should be warned of hygiene to prevent the severe consequences of ICK.

References

- Huang SC, Soong HK, Chang JS, Liang YS. Non-tuberculous mycobacterial keratitis: a study of 22 cases. *Br J Ophthalmol* 1996; **80**: 962–968.
- Alvarenga L, Freitas D, Hofling-Lima AL, Belfort R, Sampaio J, Sousa L *et al.* Infectious post-LASIK crystalline keratopathy caused by nontuberculous mycobacteria. *Cornea* 2002; **21**: 426–429.
- Wilhelmus KR, Robinson NM. Infectious crystalline keratopathy caused by *Candida albicans*. *Am J Ophthalmol* 1991; **112**: 322–325.
- Chung SH, Roh MI, Park MS, Kong YT, Lee HK, Kim EK. Mycobacterium abscessus keratitis after LASIK with IntraLase femtosecond laser. *Ophthalmologica* 2006; **220**: 277–280.

R Wong¹, YH Luo¹, MF Al-Zahawi² and TRG Poole¹

¹Ophthalmology Department, Frimley Park Hospital NHS Trust, Surrey, UK

²Microbiology Department, Frimley Park Hospital NHS Trust, Surrey, UK
E-mail: drrogerwong@aol.com

Eye (2007) **21**, 1123–1124; doi:10.1038/sj.eye.6702892;
published online 25 May 2007

Sir, Operating microscope use in scleral buckling Correspondence

Raman *et al*¹ advocate the use of the operating microscope for scleral buckling surgery. Iatrogenic scleral perforation is a recognized complication of scleral buckling surgery, and the authors elucidate manoeuvres that may decrease the likelihood of this intraoperative complication.

Time is of the essence when surgery is required to repair a macula-attached retinal detachment, as progression of the detachment to the macula will influence visual outcome significantly.² When non-elective scleral buckling surgery is undertaken for these patients, ideal surgical facilities may not be available to the surgeon—for example, the eye operating room might be unavailable, while the general operating facility is more accessible. In the absence of the operating microscope under such circumstances, the surgeon will have to rely on loupes or the naked eye to pass partial thickness scleral sutures.

The authors suggest the use of the non-dominant hand for passing scleral sutures in certain positions to access parts of the scleral quadrants. In our view, the use of the

non-dominant hand leads to considerable loss of control and proprioception, increasing the possibility of an iatrogenic perforation. Also, the frequency of scleral buckling surgery has decreased somewhat, following the universal adoption of extracapsular cataract extraction techniques.^{3,4} This has diminished the opportunity for trainee surgeons to develop ambidexterity for scleral suturing manoeuvres.

The authors assert that using the operating microscope promotes comfort and good posture, since the surgeon remains seated 'all the time' during the operation. We believe that scleral buckling surgery is necessarily a dynamic operation in which the surgeon can remain seated only some of the time, because indirect ophthalmoscopy must still be performed at various points during the surgery when the surgeon will stand and the microscope will be removed from the field.

References

- Raman SV, Smith M, Simcock PR. Use of the operating microscope for scleral buckling. *Eye* 2007; **21**: 103–104.
- Ross WH. Visual recovery after macula-off retinal detachment. *Eye* 2002; **16**: 440–446.
- Naeser K, Nielsen NE. Retinal detachment following intracapsular and extracapsular cataract extraction. *J Cataract Refract Surg* 1995; **21**: 127–131.
- Jaffe NS, Clayman HM, Jaffe MS. Retinal detachment in myopic eyes after intracapsular and extracapsular cataract extraction. *Am J Ophthalmol* 1984; **97**: 48–52.

A Singh and JM Stewart

Department of Ophthalmology,
University of California, San Francisco, CA,
USA
E-mail: stewartj@vision.ucsf.edu

Eye (2007) **21**, 1124; doi:10.1038/sj.eye.6702893;
published online 13 July 2007

Sir,
Use of the operating microscope for scleral buckling
We thank Drs Singh and Stewart for their interest in our paper.

We agree that time is definitely the essence for operating on an unstable super-temporal macula on rhegmatogenous retinal detachment. The objective of our paper¹ was to advocate the routine use of the operating microscope when it is available for buckling surgery. Not having access to the operating microscope in a scenario as described will definitely make it difficult to achieve the objective as suggested in our paper. In this situation, a loupe would be a superior alternative to surgery performed with the naked eye.

We assume that the authors probably mean to say that, with the universal adoption of phacoemulsification, cataract surgery trainee surgeons do not get adequate practice in suturing techniques.

This is quite true and makes it a compelling argument for suturing under the operating microscope, as there is greater stereopsis and magnification and less likelihood