

# Increased heterozygosity and allele variants are seen in Texel compared to Suffolk sheep

GA Walling, AD Wilson, BL Mctear and SC Bishop

Department of Genetics and Biometry, Roslin Institute (Edinburgh), Roslin, Midlothian EH25 9PS, UK

In this study, the Suffolk and Texel sheep breeds were compared for microsatellite marker heterozygosity throughout seven chromosomal regions in the sheep genome. A total of 623 Texel animals and 489 Suffolk animals in five and three half-sib families, respectively, were genotyped for microsatellite markers across the seven different chromosomes. Using the observed allele frequencies, the expected levels of heterozygosity were calculated for each family. The expected levels of heterozygosity did not significantly differ between the breeds across all regions studied. However, levels of expected heterozygosity were 32% higher in Texel animals on chromosome 4 due to a region of increased heterozygosity between BMS648 and BM3212. The number of allelic

variants significantly differed between the breeds, solely due to a region of increased number of alleles on chromosome 20. This region of higher numbers of allele variants in the Texel breed extended from the MHC to c. 15 cM distal to the MHC region incorporating markers OMHC1, CSRD226, TGLA387 and BM1818, which had 3.30, 7.02, 3.09 and 6.75 more alleles in Texel than in Suffolk animals, respectively. No difference was observed in the variance of allele frequency between the two breeds. It is proposed that previous selective sweeps may have reduced numbers of alleles and levels of heterozygosity in the Suffolk breed.

*Heredity* (2004) **92**, 102–109, advance online publication, 10 December 2003; doi:10.1038/sj.hdy.6800389

**Keywords:** sheep; microsatellite; MHC; nematode resistance; genetic variability

## Introduction

Selection and adaptation processes in animal populations succeed by means of genetic change. However, most of the changes that have occurred through domestication and subsequent selective breeding have been achieved without the understanding or monitoring of changes at the genetic level, for example, the establishment of the Booroola gene in Australian sheep flocks selected for a high incidence of multiple births (Piper and Bindon, 1982). During the past decade, with the application of genomic technologies, a substantial amount of genotype data for domestic livestock have accumulated in laboratories around the world, by scientists attempting to discover regions of the genome associated with commercially important traits (eg Walling *et al*, 2000). Arguably, however, these data have been under-utilised. For example, the large amounts of genotypic data collected for experiments to identify quantitative trait loci (QTLs) can be used to address additional questions regarding the genetic structure of the populations used in the studies.

Suffolk and Texel are the two main terminal-sire breeds of sheep used in the UK. The origins of the two breeds are distinct. Texel sheep originate from the Island of Texel in The Netherlands and were brought to the UK in the 1970s, with separate importations from France and The Netherlands. In contrast, the Suffolk sheep evolved from the mating of Norfolk horn ewes with Southdown

rams in the late 18th century, and were traditionally used around the rotational system of farming in southeast England. Breed improvement in both cases has been achieved through the subjective appraisal of breed characteristics and perceived commercial (eg carcass) characteristics, and latterly through intensive selection within structured breeding programmes known as sire reference schemes. The current selection uses a selection index designed to improve the yield of lean meat (Simm and Dingwall, 1989). Therefore, genetic change has been achieved without knowledge of the underlying genetic architecture of the breeds.

Genetic diversity differs between domesticated breeds of sheep, including the breeds in this study, with evidence suggesting lower levels of heterozygosity in Suffolk sheep when compared to other breeds including Texel sheep (Farid *et al*, 2000). More specifically, within some extensively investigated areas of the sheep genome, for example, prion protein gene, Suffolk sheep have lower levels of genetic variability in comparison to other breeds (O'Doherty *et al*, 2000, 2001), with Suffolk animals possessing only three different PrP alleles in comparison to the five present in Texel and many other breeds.

The UK sheep genome-mapping project has collected DNA from large numbers of Texel and Suffolk sheep, comprising several large half-sib families, in commercial flocks in the UK. Families from both breeds have been genotyped for many markers within selected chromosomal regions, enabling a more definitive comparison of the heterozygosity of the Texel and Suffolk breeds, than have previously been possible. The primary aim of this study is to use this large and comprehensive data set to investigate whether heterozygosity of microsatellite

Correspondence: SC Bishop, Department of Genetics and Biometry, Roslin Institute (Edinburgh), Roslin, Midlothian EH25 9PS, UK.

E-mail: Stephen.Bishop@bbsrc.ac.uk

Received 20 November 2002; accepted 10 August 2003

markers differs between Texel and Suffolk sheep at various locations across the genome. The size and thoroughness of this data set ensures that this may serve as a case study for expected heterozygosity differences in domestic livestock populations with distinct breed histories.

## Materials and methods

### DNA samples and genotyping

Blood samples were collected from c. 6-month-old lambs over a 2-year period from five half-sib families (T1–T5) of Texel sheep ( $n = 623$ ) and three half-sib families (S1–S3) from Suffolk sheep ( $n = 489$ ). The family size varied from 75 to 276 offspring per sire. All animals were born and reared in commercial flocks across the UK. DNA was extracted from the blood using a standard salt-extraction method on fresh samples, and a phenol–chloroform extraction on blood samples that had been frozen. Sires of these eight families were genotyped for microsatellite markers in up to seven regions of the genome across chromosomes 2, 3, 4, 5, 11, 18 and 20 (Table 1). All offspring were subsequently genotyped for all markers that were heterozygous in their sire, with an average of 6.18 informative typed markers per chromosome per family. Dams of progeny were not genotyped. The dams used had a mean coefficient of coancestry (Falconer, 1989) of 0.86 and 0.83% for the Suffolk and Texel breeds, respectively, and hence, due to the low level of relatedness, allele frequencies subsequently calculated were assumed to be indicative of the breeds. Alleles were determined on a within-family basis because individual families were run on separate genotyping gels; hence, common reference points were not available to distinguish between gel variations. Sires were run on all gels containing their offspring to remove the within-family variation across gels.

### Data analysis

Allele frequencies were estimated for each family for each marker that was genotyped in at least one family from each breed. This comprised 52 markers across the seven chromosomes, with the number of markers on any specific chromosome varying from four to 10. Markers and informative families are summarised in Appendix A. Allele frequencies were estimated using an expectation maximisation (EM) algorithm (Dempster *et al*, 1977), based on the frequencies of the alleles transmitted by the dam. Sire-transmitted alleles were not used in the

estimation of allele frequency. The dam's allelic contribution for each marker to each of their progeny was determined using Mendelian laws of inheritance. When the allelic contribution from the dam could not be determined, that is, when the progeny genotype was equal to the sire genotype, alleles were given a probability that they had originated from the dam based on their frequencies from the previous iteration. Both alleles were initially given equal probability that they had originated from the dam. The algorithm was iterated until convergence of allele frequencies (to six decimal places) was achieved for each marker.

The expected heterozygosity for an individual marker ( $k$ ) within a family can be calculated as

$$H_k = 1 - \sum_{a=1}^{n_k} p_a^2 \quad (1)$$

where  $H_k$  is the expected heterozygosity for marker  $k$ ,  $p_a$  is the allele frequency of allele  $a$  and  $n_k$  is the total number of alleles for marker  $k$ . The expected heterozygosity may be shown to be a function of the number of alleles and the variance of allele frequency.

Equation (1) may be rewritten as

$$H_k = 1 - n_k E(p^2) \quad (2)$$

Since  $E(p^2) = V(p) + (E(p))^2$

We may write

$$H_k = 1 - n[V(p) + (E(p))^2] \quad (3)$$

As allele frequencies for every marker in any family sum to one, the numerator of  $(E(p))^2$  also sums to one. Therefore, equation (3) can be rewritten as

$$H_k = 1 - n \left[ V(p) + \frac{1}{n^2} \right] \quad (4)$$

The expected heterozygosity is therefore a function of the number of alleles for a particular marker and the variance of allele frequency. Hence, from the numbers of alleles observed and the allele frequencies calculated in the Texel and Suffolk families, the following three statistics were derived for each informative marker within each family:

- (i) expected heterozygosity for marker  $k$  ( $H_k$ );
- (ii) total number of alleles for marker  $k$  ( $n_k$ );
- (iii) variance of allele frequency for marker  $k$  ( $V_k$ ).

This provided 190 measurements of each of the three statistics across all markers and families (36 informative markers for family S1, 12 informative markers for family S2, etc).

Each of the three measurements were analysed using Genstat REML using the model:

$$y_{ijkl} = \mu + b_i + bc_{ij} + m_k + e_{ijkl}$$

where  $y$  is the heterozygosity, number of alleles or variance of allele frequency of marker  $k$  on chromosome  $j$  in family  $l$  within breed  $i$ ,  $\mu$  is the mean,  $b_i$  is the fixed effect of breed  $i$ ,  $c_j$  is the fixed effect of chromosome  $j$ ,  $m_k$  is the random effect of marker  $k$ ,  $e_{ijkl}$  is the random error term associated with marker  $k$  on chromosome  $j$ , in family  $l$  within breed  $i$ .

In addition, the linear and quadratic covariates of family size were included in the analysis of  $n_k$ . Including

**Table 1** Summary of genomic regions investigated and families used within each region

Chromosome	Region	Families used	
		Suffolk	Texel
2	BMS2196-OarFCB11	S1, S3	T1, T2, T3, T4, T5
3	BMC1009-BM6433	S1, S2, S3	T1, T2, T3, T4
4	ILSTS62-TCRB	S1, S3	T1, T2, T4
5	TGLA176-BMS1247	S1, S3	T1, T2, T4
11	SRCRSP6-CSSM65	S1, S3	T1, T2, T4
18	BM3413-OarTMR01	S1, S2, S3	T1, T2, T3, T4, T5
20	OarCP73-McMA23	S1, S2, S3	T1, T2, T3, T4

the covariates of family size for  $H_k$  and  $V_k$  did not significantly improve the statistical model, and hence were omitted for these analyses.

For each of the three REML analyses, the predicted breed and chromosomal means were extracted with the standard errors of the differences between breeds and between the same chromosomes across breeds. Differences between chromosomal means across breeds were tested using a *t*-test, to ascertain whether differences were greater than could be explained by chance alone. To account for the number of tests performed on the seven chromosomes, a Bonferroni correction was used for each statistic by adjusting the significance level to  $P = 0.05/7 = 0.0071$ .

If a significant difference was detected between the two breeds within a chromosomal region, the data were further analysed using the model:

$$y_{ikl} = \mu + b_i + bm_{ik} + e_{ikl}$$

where all symbols are as previous described, except  $bm_{ik}$  being the fixed effect of marker  $k$  within breed  $i$ .

The predicted means were extracted from the REML analyses for markers on chromosomes that previously had significant differences between breeds, along with the standard errors of the differences between breeds. Differences of marker means between breeds were tested using a *t*-test to determine which marker(s) is responsible for the differences between breeds. To account for the number of tests performed on each chromosome, and hence reduce the risk of declaring chance results significant, a Bonferroni correction was used for each statistic by adjusting the significance level to  $P = 0.05/r$ , where  $r$  is the number of markers tested on an individual chromosome.

## Results

Results for all the three statistics across all markers are presented in Appendix B.

### Heterozygosity

The estimated REML breed means and chromosomal means within the breed for  $H_k$  are presented in Table 2. The mean  $H_k$  across all genomic regions studied, although marginally higher in the Texel population, was not significantly different between the two breeds ( $P = 0.19$ ). However, significant differences were observed on chromosome 4, with Texels having significantly higher expected heterozygosity. While moderately large differences between the breeds were present on chromosomes 2 and 20, after incorporating the Bonferroni correction for multiple testing, no significant differences of  $H_k$  were observed between the two breeds on any other chromosomes. The results from the analyses with individual markers (Table 3) indicate that the lower level of  $H_k$  in the Suffolk breed on chromosome 4 is primarily due to two adjacent markers (BMS648 and BM3212).  $H_k$  for these markers was very highly significantly different between the two breeds (0.704 and 0.642 in Texels versus 0.351 and 0.137 in Suffolks). After the incorporation of the Bonferroni correction for multiple testing, no other markers exhibited significantly different levels of expected heterozygosity.

**Table 2** REML estimated means of heterozygosity grouped by chromosome in each breed, the differences between breed and *P*-values between the two groups

Chromosome	Mean $H_k$			
	Texel	Suffolk	Difference	$P^a$
All	0.664	0.639	0.024	0.187
2	0.598	0.709	-0.111	0.012
3	0.611	0.608	0.004	0.932
4	0.682	0.515	0.167	0.005*
5	0.670	0.640	0.030	0.569
11	0.744	0.692	0.052	0.386
18	0.614	0.668	-0.054	0.093
20	0.726	0.644	0.082	0.043

<sup>a</sup>5, 1 and 0.1% significance levels incorporating the Bonferroni correction are represented by one, two and three asterisks, respectively.

**Table 3** REML estimated means of heterozygosity per marker located on chromosome 4

Chromosome	Marker	Mean $H_k$			
		Texel	Suffolk	Difference	$P^a$
4	ILSTS062	0.795	0.731	0.064	0.411
	LSCV15	0.387	0.612	-0.225	0.012
	OarHH35	0.769	0.619	0.150	0.039
	BMS648	0.704	0.351	0.353	<0.001***
	BM3212	0.642	0.137	0.505	<0.001***
	TCRB	0.734	0.639	0.096	0.284

The differences between breeds and *P*-values between the two groups are also shown. Markers are presented in linkage map order (Maddox *et al*, 2000).

<sup>a</sup>5, 1 and 0.1% significance levels incorporating the Bonferroni correction are represented by one, two and three asterisks, respectively.

### Number of alleles observed per marker

Markers had more alleles in the Texel breed, with on average 0.62 more alleles per marker in comparison to the Suffolk breed (Table 4). This was wholly due to the significantly larger number of alleles per marker on chromosome 20. Texel animals had, on average, 2.28 additional alleles per marker on chromosome 20 than Suffolk animals. No significant differences between the breeds for the number of alleles observed per marker were detected on any other chromosomes. The additional number of alleles on chromosome 20 in the Texel breed was due to significant differences at four markers in an interval incorporating the distal region of the major histocompatibility complex (MHC) to ~15 cM distal to the MHC locus (Table 5). OMHC1, CSR0226, TGLA387 and BM1818 had significantly more alleles in the Texel breed, and of all markers tested on chromosome 20 only OLADRBps and McMA23 had a higher estimated mean  $n_k$  in the Suffolk, although the differences for these other markers were not significant.

The linear and quadratic covariates for family size were 0.053 and  $-1.37 \times 10^{-4}$ , respectively, indicating increasing numbers of allele variants identified in larger families. The quadratic covariate highlights the finite nature of the number of alleles within a breed, and

**Table 4** REML estimated means of numbers of alleles per marker, grouped by chromosome in each breed, the differences between breed and *P*-values between the two groups

Chromosome	Mean $n_k$			<i>P</i> <sup>a</sup>
	Texel	Suffolk	Difference	
All	6.22	5.60	0.62	0.029*
2	6.45	6.95	-0.50	0.843
3	4.37	4.37	0.01	0.987
4	5.79	5.32	0.47	0.548
5	5.68	4.55	1.13	0.109
11	7.23	6.56	0.68	0.382
18	5.85	5.61	0.24	0.543
20	8.14	5.86	2.28	<0.001***

<sup>a</sup>5, 1 and 0.1% significance levels incorporating the Bonferroni correction are represented by one, two and three asterisks, respectively.

**Table 5** REML estimated means of numbers of alleles per chromosome 20 marker, the differences between breeds and *P*-values between the two groups

Marker	Mean $n_k$			<i>P</i> <sup>a</sup>
	Texel	Suffolk	Difference	
McMA36	3.00	2.98	0.02	0.987
BM1258	4.46	2.48	1.98	0.202
OLADRBps	6.96	9.19	-2.23	0.090
OMHC1	9.89	6.59	3.30	<0.001***
CSRD226	15.21	8.19	7.02	<0.001***
TGLA387	14.48	11.38	3.09	0.002*
BM1818	11.08	4.33	6.75	<0.001***
OarHH56	4.88	2.55	2.33	0.063
McMA23	2.68	4.46	-1.78	0.188

Markers are presented in linkage map order (Maddox *et al*, 2000). <sup>a</sup>5, 1 and 0.1% significance levels incorporating the Bonferroni correction are represented by one, two and three asterisks, respectively.

demonstrates an optimal family size past which additional allele variants are less likely to be identified.

### Allele frequency variance

The overall estimated  $V_k$  means were not significantly different between the two breeds (Table 6). The largest difference was observed on chromosome 4, where  $V_k$  means were higher in Suffolk animals. However, the standard errors of  $V_k$  estimates are high and thus this result was not statistically significant.

The analysis was repeated, including linear and quadratic covariates for the number of alleles; these were -0.017 and  $7.37 \times 10^{-4}$  respectively, both of which were significantly different from zero. While illustrating the negative correlation between  $n_k$  and  $V_k$ , this alternative model produced larger standard errors and hence did not change the overall conclusions (results not shown).

## Discussion

This study has detected a significant difference in the levels of expected heterozygosity between Suffolk and Texel sheep on chromosome 4, with Texels having increased heterozygosity. Further, a cluster of markers

**Table 6** REML estimated means of allele frequency variance per marker, grouped by chromosome in each breed, the differences between breed and *P*-values between the two groups

Chromosome	Mean $V_k (\times 100)$			<i>P</i> <sup>a</sup>
	Texel	Suffolk	Difference ( $\times 100$ )	
All	2.851	3.431	-0.580	0.310
2	4.000	2.238	1.762	0.201
3	3.787	4.955	-1.168	0.365
4	2.506	5.657	-3.151	0.091
5	2.514	2.893	-0.379	0.820
11	1.655	2.352	-0.697	0.713
18	3.449	2.631	0.818	0.415
20	2.043	3.289	-1.246	0.320

<sup>a</sup>5, 1 and 0.1% significance levels incorporating the Bonferroni correction are represented by one, two and three asterisks, respectively.

with increased numbers of alleles was also detected in the Texel breed along a region of chromosome 20 when compared to the Suffolk breed. Other than these two ~15 cM segments, no significant evidence for consistent differences between the two breeds was detected elsewhere in the regions of the genome covered by this study. This is in contrast to the previously published but smaller study of Farid *et al* (2000), in which Suffolks were apparently less heterogeneous than Texels. Our study has both a greater number of animals and a greater number of markers than the Farid *et al* (2000) study.

Consider the observed differences on chromosome 20. It is not possible to tell from our data set whether the breed differences in the number of alleles are due to the Texel breed having an enhanced number of alleles, or the Suffolks fewer alleles, compared to some common ancestor breed. Paterson *et al* (1998) genotyped three of the same microsatellite markers used in this study, in a feral population of Soay sheep. These markers were OLADRBps, OMHC1 and BM1818 and the total number of alleles at these markers for the Soay sheep were 6, 5 and 7, respectively, indicating greater similarity with Suffolks than Texels. While the number animals genotyped for each marker differed, the minimum number was 887; with such a large population, numbers of alleles can be considered indicative of the breed. However, the Soay breed often undergoes population crashes (Clutton-Brock *et al*, 1992), that is, minor genetic bottlenecks, perhaps accounting for the relatively low numbers of alleles at these markers. In contrast, in a population of 200 Scottish Blackface lambs, Schwaiger *et al* (1995) observed 19 alleles at an MHC region marker, albeit one not genotyped in this study (OLADRB).

Population bottleneck effects in the Suffolk may be suggested as a reason for the observed differences; however, they may be ruled out, as the numbers of alleles and heterozygosity did not differ consistently elsewhere in the genome. Some aspect of the evolutionary history of the two breeds will account for the difference, and, although a number of tests are available to distinguish between the different evolutionary processes responsible for causing decreases in heterozygosity (Tajima, 1983; Fay and Wu, 2000), these would require more detailed knowledge of the structure and history of populations used in this study, than is likely to be available.

It is possible that some form of selective sweep in the Suffolk breed can explain our results. The process of selection can lead to a reduction in genetic variation. The regions of increased numbers of alleles and heterozygosity between the breeds both extend across a distance of c. 15 cM. It is worth noting that linkage disequilibrium in sheep populations has been demonstrated to be large (McRae *et al*, 2002), extending for tens of centimorgans, that is, at least as large as the c. 15 cM regions between BMS648-BM3212 and OMHC1-BM1818 (Maddox *et al*, 2001), demonstrating differential heterozygosity and number of alleles in this study. Thus, these sized regions are consistent with selective sweeps during breed formation.

If such criteria could explain the lower heterozygosity and number of alleles in the Suffolk breed, genes underlying the traits undergoing selection would be expected to be located within these regions of the genome. Proximal to the region on chromosome 4 is the gene encoding Inhibin beta A (INHBA). The role of INHBA is to inhibit follicle-stimulating hormone (FSH) secretion with subsequent effects on fertility. A gene with a bearing on such a trait could easily become the focus of direct or indirect selection, and subsequently lower genetic variation and hence lower levels of heterozygosity.

Likewise, it should be noted that the region on chromosome 20 with increased numbers of alleles in Texel animals incorporated a segment of the MHC. The MHC of sheep has a similar structure to the HLA system of humans, with distinct class I and II regions each containing a number of expressed genes (Trowsdale, 1993, 1995). Marker OLADRBps is located within the MHC class II nonexpressed genes (Blattman and Beh, 1992) and marker OMHC1 is located in the MHC class I region (Groth and Wetherall, 1994). The well-established association between the MHC and the ability of a host's immune system to respond to parasitic infection caused O'Brien and Evermann (1988) to suggest that species or populations with low MHC diversity may be more vulnerable to infectious disease. Subsequently, the maintenance of genetic diversity at the MHC of vertebrates has become a paradigm for the manner in which genetic diversity may be maintained in natural populations (Hedrick, 1994). This paradigm is further supported by experimental evidence in human populations of heterozygous advantage of MHC genotypes for diseases such as hepatitis (Thurz *et al*, 1997) and AIDS (Carrington *et al*, 1999). In addition, evidence in other species suggests that the accumulation of homozygosity across the genome may result in increased parasitic burden (Cassinello *et al*, 2001). However, it must be realised that factors other than MHC diversity *per se* play a role in disease resistance. The actual effectiveness of an animal's specific alleles in enabling it to respond to an infectious challenge will be a more important factor than heterozygosity *per se*, at least when considering a specific disease, and genes elsewhere in the genome will also play an important role. As an empirical example, the high disease susceptibility and decline in desert bighorn sheep could not be explained by low MHC variation (Gutierrez-Espeleta *et al*, 2001).

In summary, this study has demonstrated that the expected degree of heterozygosity and numbers of alleles per marker locus throughout the regions of the genome

that were investigated are generally similar for Texel and Suffolk sheep, except for a region on chromosome 4 and another region within and distal to the MHC on chromosome 20. It is interesting to note that these differences occur in regions rather than at isolated markers, and the size of these regions is consistent with observed lengths of linkage disequilibrium in sheep. Thus, the reasons for these breed differences are unknown, but possible effects from previous selective sweeps merit further attention.

## Acknowledgements

This work is supported by the Department for the Environment, Food and Rural Affairs (DEFRA), Scottish Executive Environmental and Rural Affairs Department (SEERAD) and the Meat and Livestock Commission (MLC). Samples were provided by Elite Texels Sires (UK) Ltd and Suffolk Sire Referencing Scheme Ltd. We are grateful to Professor Mike Stear, the editor and two anonymous reviewers for comments on an earlier form of this manuscript.

## References

- Bishop MD, Kappes SM, Keele JW, Stone RT, Sunden SLF, Hawkins GA *et al* (1994). A genetic linkage map for cattle. *Genetics* **136**: 619–639.
- Blattman AN, Beh KJ (1992). Dinucleotide repeat polymorphism within the ovine major histocompatibility complex. *Anim Genet* **23**: 392.
- Buchanan FC, Galloway SM, Crawford AM (1991). Ovine microsatellites at the OarFCB5, OarFCB19, OarFCB20, OarFCB48, OarFCB129 and OarFCB226 loci. *Anim Genet* **25**: 60.
- Buitkamp J, Schwaiger W, Epplen JT (1993). Vb6 T-cell receptor elements in artiodactyls: conservation and germline polymorphisms. *Mamm Genome* **4**: 504–510.
- Burns BM, Taylor JF, Herring KL, Herring AD, Holder MT, Collins JS *et al* (1995). Bovine microsatellite dinucleotide repeat polymorphisms at the TEXAN11, TEXAN12, TEXAN13, TEXAN14, and TEXAN15 loci. *Anim Genet* **26**: 201–202.
- Carrington M, Nelson GW, Martin MP, Kissner T, Vlahov D, Goedert JJ *et al* (1999). HLA and HIV-1: heterozygote advantage and B\*35-Cw\*04 disadvantage. *Science* **283**: 1748–1752.
- Cassinello J, Gomendio M, Roldan ERS (2001). Relationship between coefficient of inbreeding and parasite burden in endangered gazelles. *Conserv Biol* **15**: 1171–1174.
- Charlier C, Segers K, Wagenaar D, Karim L, Berghmans S, Jaillon O *et al* (2001). Human-ovine comparative sequencing of a 250-kb imprinted locus and identification of six imprinted transcripts: DLK1, DAT, GTL2, PEG11, antiPEG11, and MEG8. *Genome Res* **11**: 850–862.
- Clutton-Brock TH, Price OF, Albon SD, Jewell PA (1992). Early development and population fluctuations in Soay sheep. *J Anim Ecol* **61**: 381–396.
- Davies KP, Maddox JF (1994). Ovine dinucleotide repeat polymorphism at the lysozyme (LYZ) locus. *Anim Genet* **25**: 204.
- Davies KP, Maddox JF (1997). Genetic linkage mapping of three anonymous ovine EST microsatellites KD101, KD103 and KD721. *Anim Genet* **28**: 455–456.
- Davies KP, Maddox JF, Harrison B, Drinkwater R (1995). Ovine dinucleotide repeat polymorphism at the CSRD226 and CSRD232 loci. *Anim Genet* **26**: 372.
- De Gortari M, Freking BA, Kappes SM, Leymaster KA, Crawford AM, Stone RT *et al* (1997). Extensive genomic

- conservation of cattle microsatellite heterozygosity in sheep. *Anim Genet* **28**: 274–290.
- Dempster AP, Laird NM, Rubin DB (1977). Maximum likelihood from incomplete data via the EM algorithm. *J Roy Stat Soc B* **39**: 1–38.
- Ede AJ, Pierson CA, Crawford AM (1995). Ovine microsatellites at the OarCP34, OarCP38, OarCP43, OarCP49, OarCP73, OarCP79 and OarCP99 loci. *Anim Genet* **26**: 130–131.
- Ede AJ, Pierson CA, Henry HM, Crawford AM (1994). Ovine microsatellites at the OarAE64, OarHH22, OarHH56, OarHH62 and OarVH4 loci. *Anim Genet* **25**: 51.
- Falconer DS (1989). *Introduction to Quantitative Genetics*, 3rd edn. Longman: England.
- Farid A, O'Reilly EO, Dollard C, Kelsey Jr CR (2000). Genetic analysis of ten sheep breeds using microsatellite markers. *Can J Anim Sci* **80**: 9–17.
- Fay JC, Wu CI (2000). Hitchhiking under positive Darwinian selection. *Genetics* **155**: 1405–1413.
- Freking BA, Keele JW, Beattie CW, Kappes SM, Smith TPL, Sonstegard TS *et al* (1998). Evaluation of the ovine Callipyge locus: I. Relative chromosomal position and gene action. *J Anim Sci* **76**: 2062–2071.
- Georges M, Massey J (1992). *Polymorphic DNA Markers in Bovidae*, World Intellectual Property Org.: Geneva (WO Publication No. 92/13102).
- Groth DM, Wetherall JD (1994). Dinucleotide repeat polymorphism within the ovine major histocompatibility complex class-I region. *Anim Genet* **25**: 61.
- Gutierrez-Espeleta GA, Hedrick PW, Kalinowski ST, Garrigan D, Boyce WM (2001). Is the decline of the desert bighorn sheep from infectious disease the result of low MHC variation? *Heredity* **86**: 439–450.
- Hedrick PW (1994). Evolutionary genetics of the major histocompatibility complex. *Am Nat* **143**: 945–964.
- Henry HM, Penty JM, Pierson CA, Crawford AM (1993). Ovine microsatellites at the OarHH35, OarHH41, OarHH44, OarHH47 and OarHH64 loci. *Anim Genet* **24**: 222.
- Hulme DJ, Silk JP, Redwin JM, Barendse W, Beh KJ (1994). Ten polymorphic ovine microsatellites. *Anim Genet* **25**: 434–435.
- Kossarek LM, Grosse WM, Finlay O, McGraw RA (1993). Bovine dinucleotide repeat polymorphism RM006. *J Anim Sci* **71**: 3176.
- Maddox JF, Davies KP, Crawford AM, Hulme DJ, Vaiman D, Cribeu EP *et al* (2001). An enhanced linkage map of the sheep genome comprising more than 1000 loci. *Genome Res* **11**: 1275–1289.
- Maddox JF, Schibler L, Cribeu EP, Kang N, Davies KP, Vaiman D (2000). Linkage mapping of goat ChirUCO, LSCV and SR-CRSP microsatellites in sheep. *Anim Genet* **31**: 145 (correction 292–293).
- McRae AF, McEwan JC, Dodds KG, Wilson T, Crawford AM, Slate J (2002). Linkage disequilibrium in domestic sheep. *Genetics* **160**: 1113–1122.
- O'Brien SJ, Evermann JF (1988). Interactive influence of infectious disease and genetic diversity in natural populations. *Trends Ecol Evol* **3**: 254–259.
- O'Doherty E, Aherne M, Ennis S, Weavers E, Hunter N, Roche JF *et al* (2000). Detection of polymorphisms in the prion protein gene in a population of Irish Suffolk sheep. *Vet Rec* **146**: 335–338.
- O'Doherty E, Aherne M, Ennis S, Weavers E, Roche JF, Sweeney T (2001). Prion protein gene polymorphisms in pedigree sheep in Ireland. *Res Vet Sci* **70**: 51–56.
- Paterson S, Wilson K, Pemberton JM (1998). Major histocompatibility complex variation associated with juvenile survival and parasite resistance in a large unmanaged ungulate population (*Ovis aries*). *Proc Natl Acad Sci USA* **95**: 3714–3719.
- Penty JM, Henry HM, Ede AJ, Crawford AM (1993). Ovine microsatellites at the OarAE16, OarAE54, OarAE57, OarAE119 and OarAE129 loci. *Anim Genet* **24**: 219.
- Pierson CA, Ede AJ, Crawford AM (1994). Ovine microsatellites at the OarHH30, OarHH51, OarVH54, OarCP88, OarCP93 and OarCP134 loci. *Anim Genet* **25**: 294–295.
- Piper LR, Bindon BM (1982). The Booroola Merino and the performance of medium non-Peppin crosses at Armidale. In: Piper LR, Bindon BM, Nethery RD (eds). *The Booroola Merino*, CSIRO: Melbourne pp 9–20.
- Robertson TM, Lord EA, Glass BC, Dodds KG, Broad TE (2001). Microsatellites isolated from BAC clones containing v-akt1 murine thymoma viral oncogene homolog 1 and bradykinin receptor B2 assigned to sheep chromosome 18 by linkage analysis. *J Anim Sci* **79**: 550–551.
- Simm G, Dingwall WS (1989). Selection indexes for lean meat production in sheep. *Livestock Prod Sci* **21**: 223–233.
- Schwaiger F-W, Gostomski D, Stear MJ, Duncan JL, McKellar QA, Epplen JT *et al* (1995). An ovine major histocompatibility complex DRB1 allele is associated with low faecal egg counts following natural, predominantly *Ostertagia circumcincta* infection. *Int J Parasitol* **25**: 815–822.
- Smith AJ, Hulme DJ, Silk JP, Redwin JM, Beh KJ (1995). Thirteen polymorphic ovine microsatellites. *Anim Genet* **26**: 277–278.
- Swarbrick PA, Buchanan FC, Crawford AM (1990). Ovine dinucleotide repeat polymorphism at the MAF23 locus. *Anim Genet* **21**: 191.
- Tabet-Aoul K, Schibler L, Vaiman D, Oustry-Vaiman A, Lantier I, Saisi-Mehtar N *et al* (2000). Regional characterisation of a hamster-sheep somatic cell hybrid panel. *Mamm Genome* **11**: 37–40.
- Tajima F (1983). Evolutionary relationship of DNA sequences in finite populations. *Genetics* **134**: 1289–1303.
- Thurz MR, Thomas HC, Greenwood BM, Hill AVS (1997). Heterozygote advantage for HLA class-II type in hepatitis B virus infection. *Nat Genet* **17**: 11–12.
- Trowsdale J (1993). Genomic structure and function in the MHC. *Trends Genet* **9**: 117–122.
- Trowsdale J (1995). Both man and bird and beast – comparative organisation of MHC genes. *Immunogenetics* **41**: 1–17.
- Vaiman D, Brunialti A, Bensaado M, Derbois C, Vaiman A, Crawford AM *et al* (2000). Isolation of subtelomeric DNA sequences labelling sheep and goat chromosome ends. *Genet Sel Evol* **32**: 599–619.
- Walling GA, Visscher PM, Andersson L, Rothschild MF, Wang L, Moser G *et al* (2000). Combined analyses of data from QTL mapping studies: chromosome 4 effects on porcine growth and fatness. *Genetics* **155**: 1369–1378.

## Appendix A

The summary of markers and informative families is given in Table A1.

## Appendix B

Results for all statistics for all markers are shown in Table B1.

Table A1

Marker	Chr	Informative families		Reference
		Suffolk	Texel	
TGLA10	2	S3	T2, T4	Georges and Massey (1992)
BM81124		S1, S3	T4	De Gortari <i>et al</i> (1997)
INRA40		S1	T2, T4, T5	Tabet-Aoul <i>et al</i> (2000)
TEXAN2		S1, S3	T1	De Gortari <i>et al</i> (1997)
OarHH30		S1	T1, T2	Pierson <i>et al</i> (1994)
ILSTS030		S1, S3	T1, T3, T5	De Gortari <i>et al</i> (1997)
OarFCB20		S1, S3	T1, T2, T4, T5	Buchanan <i>et al</i> (1991)
BMC1009	3	S1, S2, S3	T2, T3, T4	Bishop <i>et al</i> (1994)
KD103		S3	T1, T2	Davies and Maddox (1997)
BL4		S2	T3	De Gortari <i>et al</i> (1997)
LYZ		S1, S3	T3, T4	Davies and Maddox (1994)
MAF23		S2, S3	T1	Swarbrick <i>et al</i> (1990)
OarCP43		S2	T3	Ede <i>et al</i> (1995)
TEXAN15		S3	T1	Burns <i>et al</i> (1995)
BM8230		S1	T2, T4	De Gortari <i>et al</i> (1997)
BM6433	S1, S3	T1, T3, T4	De Gortari <i>et al</i> (1997)	
ILSTS062	4	S1	T1, T2	De Gortari <i>et al</i> (1997)
LSCV15		S3	T4	Maddox <i>et al</i> (2000)
OarHH35		S3	T1, T2, T4	Henry <i>et al</i> (1993)
BMS648		S1	T2	De Gortari <i>et al</i> (1997)
BM3212		S1	T4	De Gortari <i>et al</i> (1997)
TCRB		S3	T2	Buitkamp <i>et al</i> (1993)
RM006	5	S3	T2	Kossarek <i>et al</i> (1993)
BMS792		S1, S3	T1	De Gortari <i>et al</i> (1997)
BM1853		S1	T2	Bishop <i>et al</i> (1994)
OarAE129		S1, S3	T2	Penty <i>et al</i> (1993)
McM527		S3	T2	Hulme <i>et al</i> (1994)
McM108		S3	T1, T4	Smith <i>et al</i> (1995)
BMS1247		S1, S3	T1, T4	De Gortari <i>et al</i> (1997)
SRCRSP6	11	S1	T1, T2	Maddox <i>et al</i> (2000)
LSCV36		S1	T1, T2, T4	Maddox <i>et al</i> (2000)
CHIRUC04		S3	T1, T2, T4	Maddox <i>et al</i> (2000)
EPCDV023		S1, S3	T1, T2, T4	Vaiman <i>et al</i> (2000)
OarHH47	18	S1, S2, S3	T1, T3, T4, T5	Henry <i>et al</i> (1993)
McM38		S1	T3, T4	Hulme <i>et al</i> (1994)
McMA26		S1, S2, S3	T1, T2, T3, T4	Unpublished
OB2		S1, S2, S3	T1, T3	Robertson <i>et al</i> (2001)
CSSM18		S1, S3	T3, T5	De Gortari <i>et al</i> (1997)
DLK		S1, S2	T3, T5	Charlier <i>et al</i> (2001)
OY3		S1, S2	T1, T2, T3, T4	Freking <i>et al</i> (1998)
OY15		S1, S3	T3, T4, T5	Freking <i>et al</i> (1998)
OY5		S1, S2, S3	T1, T2, T4	Freking <i>et al</i> (1998)
OarTMR01		S1	T1, T2	Robertson <i>et al</i> (2001)
McMA36	20	S3	T1, T2	Unpublished
BM1258		S3	T3	Bishop <i>et al</i> (1994)
OLADRBps		S1, S2	T1	Blattman and Beh (1992)
OMHC1		S1, S2, S3	T1, T3, T4	Groth and Wetherall (1994)
CSRD226		S1, S3	T4	Davies <i>et al</i> (1995)
TGLA387		S1, S3	T1, T2, T4	Georges and Massey (1992)
BM1818		S1	T1, T2, T4	De Gortari <i>et al</i> (1997)
OarHH56		S1	T1, T3, T4	Ede <i>et al</i> (1994)
McMA23		S3	T1, T3	Unpublished

**Table B1**

Marker	Chr	$H_k^a$		$n_k^b$		$V_k^c (\times 100)$	
		Suffolk	Texel	Suffolk	Texel	Suffolk	Texel
TGLA10	2	0.692	0.718	7.47	7.25	2.284	1.977
BM81124		0.779	0.071	9.41	5.20	1.142	12.714
INRA40		0.718	0.350	5.31	4.40	1.636	8.948
TEXAN2		0.820	0.652	8.92	6.90	0.765	2.934
OarHH30		0.554	0.648	4.46	6.00	4.924	3.091
ILSTS030		0.559	0.727	4.90	6.41	4.830	1.697
OarFCB20		0.808	0.742	6.89	7.11	0.654	1.579
BMC1009	3	0.771	0.669	7.25	5.90	1.246	2.744
KD103		0.739	0.621	3.44	5.51	0.272	3.509
BL4		0.759	0.631	6.00	4.37	1.245	2.965
LYZ		0.283	0.674	3.38	3.80	22.492	1.897
MAF23		0.524	0.341	2.72	2.90	4.762	10.865
OarCP43		0.645	0.641	3.00	4.37	0.732	2.738
TEXAN15		0.683	0.696	5.45	5.90	2.505	2.291
BM8230		0.628	0.723	4.31	4.65	3.052	1.546
BM6433		0.511	0.470	2.38	2.85	2.469	6.548
ILSTS062		4	0.731	0.795	7.35	7.01	1.805
LSCV15	0.612		0.387	2.47	4.31	1.814	9.070
OarHH35	0.619		0.769	6.47	7.44	3.402	1.310
BMS648	0.351		0.704	3.33	4.09	10.531	1.158
BM3212	0.137		0.642	5.31	5.29	13.271	3.170
TCRB	0.639		0.734	6.47	6.33	3.121	1.649
RM006	5		0.706	0.619	3.47	4.07	1.096
BMS792		0.642	0.841	5.89	9.93	3.195	0.588
BM1853		0.630	0.650	4.33	3.09	3.003	0.546
OarAE129		0.649	0.476	4.33	3.15	2.521	6.372
McM527		0.707	0.583	4.45	5.11	1.856	4.336
McM108		0.619	0.753	5.46	8.74	3.570	4.564
BMS1247		0.599	0.691	3.89	5.11	3.785	2.177
SRCRSP6		11	0.639	0.685	5.37	6.01	3.230
LSCV36	0.748		0.802	8.33	8.11	1.585	0.906
CHIRUC04	0.651		0.787	4.45	8.79	2.988	1.122
EPCDV023	0.701		0.695	6.90	6.14	2.035	2.306
OarHH47	18	0.740	0.680	8.28	7.85	1.691	2.464
McM38		0.747	0.619	5.33	5.39	1.060	3.583
McMA26		0.822	0.719	9.26	7.92	0.765	1.951
OB2		0.675	0.633	5.93	4.14	2.641	2.230
CSSM18		0.642	0.418	3.39	5.18	1.836	8.186
DLK		0.503	0.603	4.71	4.71	6.059	3.688
OY3		0.619	0.551	3.65	6.91	2.589	4.529
OY15		0.758	0.655	5.39	4.18	1.087	1.946
OY5		0.701	0.693	4.95	7.74	1.971	2.335
OarTMR01		0.363	0.559	4.33	4.06	9.864	4.777
McMA36		20	0.537	0.641	2.98	3.00	4.310
BM1258	0.446		0.658	2.48	4.46	7.356	2.303
OLADRBps	0.810		0.730	9.19	6.96	0.846	1.812
OMHC1	0.761		0.813	6.59	9.89	1.365	0.912
CSR226	0.754		0.833	8.19	15.21	1.547	0.656
TGLA387	0.718		0.856	11.39	14.48	1.852	0.486
BM1818	0.687		0.826	4.33	11.08	1.588	0.750
OarHH56	0.204		0.714	2.55	4.88	14.793	1.718
McMA23	0.682		0.362	4.46	2.68	2.351	10.182

<sup>a</sup>Standard error of differences range 0.089–0.042 with mean 0.073.

<sup>b</sup>Standard error of differences range 1.57–0.72 with mean 1.25.

<sup>c</sup>Standard error of differences range 0.0387–0.0184 with mean 0.0316.