

Population genetic differentiation in taxa of *Lotus* (Fabaceae: Loteae) endemic to the Gran Canarian pine forest

F Oliva-Tejera¹, J Caujapé-Castells¹, J Naranjo-Suárez¹, J Navarro-Déniz¹, JR Acebes-Ginovés² and D Bramwell¹

¹Jardín Botánico Canario Viera y Clavijo, Ap. de correos 14 de Tafira Alta, 35017 Las Palmas de Gran Canaria, Spain; ²Departamento de Biología Vegetal (Botánica), Facultad de Farmacia, Universidad de La Laguna, 38071 La Laguna, Tenerife, Spain

A survey of allozyme variation at 17 loci in 14 populations representing four taxonomically problematic Gran Canarian pine forest endemic taxa of *Lotus* (*L. genistoides*, *L. holosericeus*, *L. spartioides* and some taxonomically uncertain populations collected under the designation *Lotus* sp.) was conducted to examine their diversification and systematic relationships. All groups exhibited high values of genetic variation, although inbreeding was common within populations. Considerable among-population genetic homogeneity was detected, as inferred from low values of *Gst* within each of the groups. The high population sizes of these taxa and a lack of evidence for isolation by distance or genetic bottlenecks indicate that diversity has accumulated

over a long period of environmental stability. The association of high genetic distances with low linear distances, and the substantial increase in the values of *Gst* when the taxa considered were merged in different combinations hint at an incipient (yet probably taxonomically insufficient) reproductive isolation. The genetic similarity between *L. genistoides*, *L. holosericeus* and *L. spartioides*, together with the different behaviour of the populations collected under the designation *Lotus* sp., may have important implications for the restructuring of the taxonomy of this group when the ongoing morphological studies are completed.

Heredity (2005) 94, 199–206. doi:10.1038/sj.hdy.6800598
Published online 10 November 2004

Keywords: *Lotus*; Canary Islands; endemics; isozymes; diversification; taxonomy

Introduction

Lotus L. (Fabaceae: Loteae) is a cosmopolitan genus whose Canarian circumscription consists of ca. 24 species, of which 18 are endemic to this oceanic archipelago according to Bramwell (2002), and Acebes Ginovés *et al* (2001). The Canarian species of *Lotus* can be ascribed to one of two sections: *Heinekenia* and *Pedrosia*. While section *Heinekenia* is exclusively Canarian, the geographic distribution of section *Pedrosia* encompasses all Macaronesian islands and the South of Morocco (Sunding, 1979), with *L. arenarius* Brot. also distributed in the Iberian Peninsula (Talavera *et al*, 2000).

Gran Canaria hosts 12 species of *Lotus*. Six of these are endemic to the island, one is endemic to Macaronesia, two are native and three are introduced (Hansen and Sunding, 1993). All six of the exclusively Gran Canarian endemics belong to section *Pedrosia*, and are listed in the Red List of Spanish Vascular Flora (VV AA, 2000) as critically endangered (*L. arinagensis* Bramwell, *L. kunkelii* (Esteve) Bramwell & Davis and *L. genistoides* Webb ex Pit & Proust), endangered (*L. callis-viridis* Bramwell and Davis), vulnerable (*L. spartioides* Webb & Berthel) and

minor risk (*L. holosericeus* Webb & Berthel.). The former five taxa are also listed as priority in the Atlas of endangered Spanish Vascular Flora (Bañares *et al*, 2003).

Although some of these six species are easily recognizable, the taxonomic distinction among *L. holosericeus*, *L. genistoides* and *L. spartioides* is not based on morphologically sound characters and raises considerable controversy. The lack of consensus to define operative biological units in these three Gran Canarian *Lotus* hinders the study of their diversification, and thus makes it difficult to develop a conservation strategy.

Some of the difficulties stem from the fact that the ecological distributions of these taxa overlap in the pine forests of the Gran Canarian summits (Figure 1). Although the name *L. genistoides* is used locally to refer to the populations of *Lotus* occurring in Tirajana (Webb ex Pitard and Proust, 1908), there is no formal description of this taxon, and the material used to obtain its chromosome number was collected in Guayadeque (Aldridge and Ortega, 1976), a 5 km long ravine running from the centre of the island to its eastern coast. However, recent surveys of this ravine have failed to find a single specimen of the taxon. *L. spartioides* is considered as typical from the Pine forest of Tamadaba (in the North West of Gran Canaria) (Sunding, 1972), but the material of the original description (Webb and Berthelot, 1836–50) was collected in the area of the Caldera de Tirajana (in the South of Gran Canaria), without any further specification. There are also a

Correspondence: J Caujapé-Castells, Jardín Botánico Canario Viera y Clavijo, Ap. de correos 14 de Tafira Alta, 35017 Las Palmas de Gran Canaria, Spain. E-mail: julcaujape@grancanaria.com
Received 1 October 2003; accepted 31 August 2004; published online 10 November 2004



Figure 1 Locations of the 14 populations of Gran Canarian *Lotus* surveyed. Numerical codes are those presented in Table 1.

considerable number of cases where material collected in the same area by the same collector has received different names, confusing *L. spartioides* and *L. genistoides* (Ortega, 1976, 1979, respectively) and *L. spartioides* and *L. holosericeus* (Kunkel, 1969, 1977, respectively). The taxonomic uncertainty in these Gran Canarian endemic species of *Lotus* indirectly suggests that a survey with molecular markers might help improve understanding of their diversification and systematic relationships.

Allozyme polymorphisms have been frequently used as neutral molecular markers to study levels and structuring of population genetic variation within species (see Hamrick and Godt (1989) for a comprehensive review), including several species of *Lotus* (Reason and Grant, 1988, 1989; Gauthier *et al*, 1998). Here, we use them to clarify the relationships among taxa within the complex of Gran Canarian *Lotus*.

Materials and methods

Plant material

L. spartioides is locally common in the Tamadaba pine forest, North West of Gran Canaria (Bramwell and Bramwell, 2001). It is a small subshrub with yellow flowers, small procumbent or suberect branches and linear leaflets of 5–15 mm in length that are covered with short whitish hairs. It is very frequent between 1000 and 1400 m and shows a high degree of variation in flower size and width, leaf shape and degree of hairiness.

L. holosericeus is widely distributed in the central part of the island of Gran Canaria. It is similar to *L. spartioides*, but it possesses long silky hairs that cover all of its surface. It inhabits the clearings and the shrubs in the area of Canarian Pine forests in several ravines.

L. genistoides is one of the less known species of Gran Canarian *Lotus*, with a single locality in the Caldera de Tirajana, a huge volcanic crater in the south of Gran Canaria (Bramwell and Bramwell, 2001). *L. genistoides* is woody and more robust than the other two taxa, with a smaller flower size.

Available cytogenetic data indicate that all *Lotus* taxa from Gran Canaria are $2n = 14$ (Aldridge and Ortega, 1976; Ortega, 1976).

Table 1 Details of the sampling for this survey

Taxon/population	Code	N	U.T.M.
<i>L. genistoides</i>			
1. Cañadón del Jierro	LGCJ	50	28RDR 433 923
<i>L. holosericeus</i>			
2. Barranco del Arco	LHAR	75	28RDR 394 863
3. Cruz de la Umbría	LHCU	15	28RDR 406 856
4. Pílancones	LHPI	30	28RDR 415 874
5. Pino Pílancones	LHPP	25	28RDR 408 856
6. Degollada de Rosiana	LHRO	25	28RDR 420 876
7. Las Tederas	LHTE	10	28RDR 396 852
<i>L. spartioides</i>			
8. Roque Bentayga	LSBE	30	28RDR 368 963
9. Tamadaba	LSTA	100	28RDS 317 018
<i>Lotus sp.</i>			
10. Canal de la Mina	LSPCM	50	28RDR 417 955
11. Caldera de los Marteles	LSPMA	50	28RDR 470 929
12. Roque Nublo	LSPNU	50	28RDR 397 940
13. Pajonales	LSPPA	25	28RDR 351 905
14. Presa de las Niñas	LSPPN	65	28RDR 368 903

Population numbers correspond to Figure 1. N: sample size.

Sampling

We sampled a total of 600 individuals from 14 populations representing the distribution of *L. genistoides* (one population), *L. holosericeus* (six populations), *L. spartioides* (two populations) and five taxonomically undetermined populations that we will refer to as *Lotus sp.* (Figure 1, Table 1). Since most target populations occupy a large area, each sampling was preceded by a visual survey to provide an idea of population size and area. Based on these surveys, sampling was carried out along transects that subdivided the area of a population into several equally spaced points. Five to 10 individuals (depending on abundance) were sampled from each point. This procedure aimed to provide a representative sample of individuals throughout a population. Sample sizes are strictly related to the size of target populations and ranged from 10 in population LHTE to 100 in population LSTA (Table 1). Leaf samples from individual plants were deposited in numbered zip-lock plastic bags that were kept refrigerated in a portable cooler until they were stored in -80°C freezers at the Jardín Botánico Canario Viera y Clavijo (JBCVC). Samples remained there until used for extract preparation.

Electrophoretic analyses: For each sample, about 0.1 g of frozen leaves per individual were ground in a mortar using an extraction buffer following Shields *et al* (1983). The resulting protein extract was adsorbed on to a 4 mm \times 6 mm Whatman No. 3 chromatography paper wick and stored at -80°C until electrophoresed.

Horizontal starch-gel electrophoresis was carried out in 12% w/v gels using two different gel/electrode buffer systems (Shields *et al*, 1983) that allowed examination of eight enzyme systems. Histidine pH 7.0 (System E in Shields *et al*, 1983) was used to examine phosphoglucutase (PGM, E.C. 5.4.2.2), phosphoglucose isomerase (PGI, E.C. 5.3.1.9), isocitrate dehydrogenase (IDH, E.C.1.1.1.42) and esterase (EST, 3.1.1.1). Morpholine-Citrate pH 6.1 (Clayton and Tretiak, 1972) was used to examine phosphogluconate dehydrogenase

(6PGD, E.C.1.1.1.44), shikimate dehydrogenase (SKD, E.C.1.1.1.25), malate dehydrogenase (MDH, E.C.1.1.1.37) and malic enzyme (ME, E.C.1.1.1.40).

For each enzyme, gene loci and alleles were inferred and labelled following numerical and alphabetical sequence, respectively. Intrapopulation, interpopulation and interspecific verifications of enzyme mobilities were determined through side-by-side comparisons of allelic variants on the same gel.

Data analysis: Elementary descriptors of isozyme variation (number of alleles per locus, percentage of polymorphic loci, observed and expected heterozygosity), and genetic distances and identities (Nei, 1978) were calculated using BIOSYS-1 Version 1.7 (Swofford and Selander, 1989). All calculations were made at the species and population levels from genotype data corresponding to each locus. The effective number of alleles (A_e), estimates of interpopulation gene flow and Ewens–Watterson neutrality tests were calculated using Popgene version 1.32 (1997).

Nei's (1973) and Wright's (1951) population-structure statistics were calculated over all loci for *L. holosericeus*, *L. spartioides*, *Lotus* sp. and for all populations (regardless of taxonomic differences) using GeneStat-PC 3.31 (Lewis and Whitkus, 1993) and BIOSYS-1 Version 1.7 (Swofford and Selander, 1989), respectively.

We applied a sign test for heterozygosity excess (Cornuet and Luikart, 1996) to detect whether populations had experienced recent historical bottlenecks. This test compares expected heterozygosity (H_e) under Hardy–Weinberg equilibrium expectations to the heterozygosity expected at mutation-drift equilibrium (H_{eq}) in a sample that has the same size and the same number of alleles as the sample used to measure H_e (Luikart and Cornuet, 1998). The rationale for this test is that, since low frequency alleles are lost at a much faster rate than heterozygosity in a bottleneck situation, bottlenecked populations are expected to have a heterozygote excess. Calculations were made based on allele frequency data under the Stepwise Mutation Model (SMM) and the Independent Allele Model (IAM) using the program Bottleneck-PC (Piry et al, 1998).

The relationship between geographic location and genetic makeup of all populations sampled was evaluated by Mantel (1967) tests carried out using the computer program NTSYS-pc version 2.02j (Rohlf, 1998).

Results

Genetic interpretation of the eight enzymes examined allowed the resolution of 17 putative loci, none of which was monomorphic over all populations surveyed. Of the 70 alleles detected, populations LGCJ (*L. genistoides*), LSTA (*L. spartioides*) and LSPCM and LSPPN (*Lotus* sp.) possessed one exclusive allele each (*Mdh-4c*, *6Pgd-4d*, *Pgm-1g* and *Mdh-1e*, respectively). There were also exclusive allele sharings between the population pairs LGCJ-LSPPN (*Est-3e*), LSPCM-LSPPA (*Idh-1d*), LHAR-LHCU (*6Pgd-4a*), LHAR-LSTA (*Pgi-1b*), LSPCM-LSPMA (*Pgm-2e*) and LHRO-LSPCM (*Skd-2a*). The remaining 60 alleles were shared by different combinations of the 14 populations surveyed. No taxon-diagnostic alleles were found (ie, alleles monomorphic in one taxon and not shared by the other taxa). Also, there were no alleles that

were present exclusively in all populations of one taxon (the table of allele frequencies is available upon request).

The basic indicators of polymorphism (Table 2) showed that levels of genetic variation are very similar in each taxon. *L. holosericeus* contained the maximum number of alleles per locus within a population ($A_l = 2.8$ in population LHAR) and the maximum proportion of polymorphic loci ($P = 76.5$ in population LHPI), but also the minimum values of these parameters ($A_1 = 1.6$ and $P = 41.2$ in LHCU). The values of observed heterozygosity ranged from $H_o = 0.067$ in population LHAR (*L. holosericeus*) to $H_o = 0.138$ in populations LHTE (*L. holosericeus*) and LSPMA (*Lotus* sp.). Expected heterozygosity (H_e) spanned from $H_e = 0.115$ (population LHCU) to $H_e = 0.204$ (population LHPI), both in *L. holosericeus*. The inbreeding coefficient displayed considerable variation across loci and its average values ranged from $F_{is} = 0.294$ in *L. spartioides* to $F_{is} = 0.543$ in *L. holosericeus* (Table 3). These values are all much higher than zero (the value that would correspond to a situation of random mating), and indicate a predominance of selfing in the populations surveyed. Consistent with these results, only five out of the 17 loci were in Hardy–Weinberg proportions in *L. spartioides*, two in *Lotus* sp. and one in *L. holosericeus* and *L. genistoides* (Table 3). None of these populations showed evidence of a recent bottleneck (Table 2), and all loci could be considered neutral according to Ewens–Watterson tests (Table 3).

The proportion of variation that is explained by the differentiation among populations (Table 3) was uniformly low in *L. spartioides* (average $G_{st} = 0.027$) and *L. holosericeus* (average $G_{st} = 0.089$). By contrast, *Lotus* sp. exhibited a much higher interpopulation differentiation, with an average value of $G_{st} = 0.282$.

The values of G_{st} increased sharply when the assemblages were considered in different combinations ($G_{st} = 0.290$ for all populations, $G_{st} = 0.293$ for all populations save for *L. genistoides*, $G_{st} = 0.277$ for *L. spartioides* plus *Lotus* sp. and $G_{st} = 0.159$ for *L. holosericeus* plus *Lotus* sp.).

Average Nei's (1978) genetic distances were $D = 0.014$ within *L. holosericeus*, $D = 0.008$ within *L. spartioides* and $D = 0.112$ within *Lotus* sp. *Lotus* sp. was the more genetically heterogeneous assemblage, with a maximum distance value of $D = 0.203$ (Table 4), more than five-fold the maximum genetic distance within *L. holosericeus* ($D = 0.033$) and within *L. spartioides* ($D = 0.008$). The highest genetic distance ($D = 0.245$) was detected between one population of *L. holosericeus* and one of *Lotus* sp. (LHTE-LSPMA) and the lowest ($D = 0.002$) within *L. holosericeus* (LHRO-LHPP). Average genetic distances between these assemblages were also relatively low, with a maximum value of $D = 0.144$ between *L. genistoides* and *Lotus* sp. and a minimum of $D = 0.009$ between *L. holosericeus* and *L. genistoides*.

The UPGMA dendrogram constructed from Nei's (1978) genetic distances (Figure 2) resolved three consistent groups containing: (1) *L. holosericeus* populations, one population of *Lotus* sp. (LSPPA) and the population of *L. genistoides*; (2) *L. spartioides* populations and one population of *Lotus* sp. (LSPNU) and (3) the remaining three populations that we assigned to *Lotus* sp.

Mantel tests revealed no significant relationship between genetic and geographic distances within

Table 2 Basic indicators of isozyme variation for the 14 populations of *Lotus* surveyed

Population	Bottleneck tests											
	T	A_l	A_e	P	H_o	H_e	PL	IAM		SMM		
								H_d/H_e	P	H_d/H_e	P	
<i>L. genistoides</i>												
1. LGCJ	38	2.2 (0.3)	1.2 (0.3)	52.9	0.091 (0.028)	0.135 (0.034)	11	9/2	0.021	9/2	0.005	
<i>L. holosericeus</i>												
2. LHAR	48	2.8 (0.3)	1.2 (0.2)	52.9	0.067 (0.019)	0.143 (0.030)	15	15/0	0.000	15/0	0.000	
3. LHCU	38	1.6 (0.2)	1.2 (0.3)	41.2	0.075 (0.036)	0.115 (0.039)	13	11/2	0.002	11/2	0.001	
4. LHPI	43	2.5 (0.2)	1.3 (0.4)	76.5	0.083 (0.024)	0.204 (0.037)	15	12/3	0.010	12/3	0.001	
5. LHPP	27	2.2 (0.2)	1.2 (0.3)	47.1	0.079 (0.020)	0.161 (0.040)	8	5/3	0.641	6/2	0.074	
6. LHRO	39	2.3 (0.2)	1.2 (0.3)	64.7	0.089 (0.026)	0.168 (0.034)	12	11/1	0.008	11/1	0.002	
7. LHTE	29	1.7 (0.2)	1.3 (0.3)	52.9	0.138 (0.051)	0.175 (0.048)	8	5/3	0.250	5/3	0.250	
Average <i>L. holosericeus</i>		2.2 (0.2)	1.2 (0.3)	55.9	0.089 (0.029)	0.161 (0.038)						
<i>L. spartioides</i>												
8. LSBE	31	1.8 (0.2)	1.3 (0.3)	52.9	0.118 (0.033)	0.160 (0.041)	11	8/3	0.831	8/3	0.365	
9. LSTA	39	2.3 (0.3)	1.3 (0.3)	64.7	0.105 (0.030)	0.160 (0.036)	11	8/3	0.123	11/0	0.000	
Average <i>L. spartioides</i>		2.1 (0.3)	1.3 (0.3)	58.8	0.112 (0.032)	0.160 (0.039)						
<i>Lotus sp.</i>												
10. LSPCM	45	2.6 (0.2)	1.3 (0.4)	58.8	0.104 (0.033)	0.187 (0.046)	15	12/3	0.055	13/2	0.010	
11. LSPMA	34	2.0 (0.3)	1.3 (0.4)	52.9	0.138 (0.041)	0.200 (0.052)	10	5/5	0.846	6/4	0.193	
12. LSPNU	34	2.0 (0.2)	1.2 (0.3)	52.9	0.093 (0.037)	0.159 (0.042)	11	7/4	0.520	7/4	0.175	
13. LSPPA	30	1.8 (0.2)	1.2 (0.3)	52.9	0.100 (0.032)	0.156 (0.040)	10	7/3	0.375	7/3	0.032	
14. LSPPN	46	2.7 (0.2)	1.3 (0.4)	58.8	0.119 (0.035)	0.201 (0.044)	16	11/5	0.211	12/4	0.025	
Average <i>Lotus sp.</i>		2.2 (0.2)	1.3 (0.4)	55.3	0.111 (0.036)	0.181 (0.045)						
Total averages		2.2 (0.2)	1.3 (0.3)	55.9	0.100 (0.032)	0.166 (0.040)						

T: total number of alleles scored; A_l : average number of alleles per locus; A_e : average effective number of alleles per locus; P: proportion of polymorphic loci (95% criterion); H_o and H_e : observed and expected heterozygosities. PL: number of polymorphic loci used in the bottleneck tests; H_d/H_e : number of loci with heterozygote deficiency and excess (respectively) according to the Independent Allele Model (IAM) and the Stepwise Mutation Model (SMM); P: probability of the test. Values in brackets are standard deviations.

L. holosericeus ($r=0.187$, $P=0.241$), within *Lotus sp.* ($r=0.539$, $P=0.102$) or in all the populations sampled ($r=0.115$, $P=0.107$).

Discussion

General levels of genetic variation

Values for genetic polymorphism in these Gran Canarian endemic *Lotus* species are about two-fold higher than the average values reported by Hamrick and Godt (1989) for endemic plants ($A=1.39$, $P=0.26$, $H_e=0.063$). Furthermore, they are only slightly lower than those detected by Gauthier *et al.* (1998) in the mainland diploid species *L. alpinus* (Schleich ex DC) Ramond (averages $A=2.8$, $H_o=0.219$, $H_e=0.279$). The degree of genetic variation within these taxa, as measured by the average population diversity (H_s , Table 3), is in all cases much higher than the averages published for plants of Hawaii Islands ($H_s=0.064$, DeJooide and Wendel, 1992) or the Juan Fernández Islands ($H_s=0.042$, Crawford *et al.*, 2001) and slightly higher than the average value published for Canarian taxa ($H_s=0.137$, Francisco-Ortega *et al.*, 2000). It is clear that populations of Gran Canarian endemic *Lotus* maintain substantial levels of genetic variation, thereby adding to the emerging picture of more variation in Gran Canarian endemics relative to those from other oceanic archipelagos (Francisco-Ortega *et al.*, 2000). The high population sizes of these taxa and lack of evidence for genetic bottlenecks (Table 2) suggest that genetic varia-

tion has increased in a context of environmental stability. Lack of continental populations of section *Pedrosia* in our survey prevent us from knowing whether the levels of variation of these Gran Canarian populations are quantitatively and/or qualitatively different from their closely related mainland congeners.

Since high F_{is} values are estimated in most populations (Table 3), this variation appears to be maintained despite an overall predominance of inbreeding in the populations surveyed. When the seeds of *Lotus* have ripened, they detach by their own weight and stay near the parent plant (gravity dispersal or barochory). Barochory fosters small genetic neighbourhoods, where reproduction takes place between related individuals. The F_{is} values estimated for these Gran Canarian *Lotus* are much higher than those obtained in the Californian *L. scoparius* (Nutt.) Ottley, where Montalvo, Clegg and Ellstrand (unpublished data) report low inbreeding in all populations examined based on a study of 14 allozyme loci. The high F_{is} values in these Canarian *Lotus* may therefore stem from populations being divided into different mating areas leading to a deficit in heterozygotes in the pooled population due to the Wahlund effect.

However, *L. spartioides* and *L. holosericeus* exhibit a considerable genetic homogeneity over populations, which hints at an abundant interpopulation gene flow and manifests qualitatively in the fact that exclusive alleles are very infrequent and scattered throughout the populations. Quantitatively, levels of interpopulation differentiation as estimated by G_{st} and F_{st} (Table 3) are

Table 3 Multilocus (χ^2 and neutrality tests) and population structure statistics according to Nei (1973) and Wright (1951) for the polymorphic loci in *L. spartioides*, *L. holosericeus* and *Lotus sp*

Locus	Multi-locus structure statistics					Population structure statistics					
	Hardy–Weinberg equilibrium		Ewens–Watterson neutrality tests			Nei's (1973) unmodified			Wright's F statistics		
	χ^2	P	Avg.	L95	U95	H _s	D _{st}	G _{st}	F _{is}	F _{it}	F _{st}
<i>L. spartioides</i>											
EST-3	35.6	0.000	0.680	0.367	0.969	0.097	0.001	0.011	0.270	0.278	0.011
IDH-1	7.0	0.008	0.815	0.502	0.984	0.068	0.003	0.038	0.384	0.407	0.038
MDH-2	0.1	0.822	0.816	0.502	0.984	0.090	0.005	0.053	-0.111	-0.053	0.053
MDH-4	0.0	0.875	0.813	0.503	0.984	0.090	0.005	0.053	-0.111	-0.053	0.053
ME-1	0.0	0.928	0.811	0.502	0.984	0.050	0.001	0.020	0.792	0.796	0.020
6PGD-1	41.7	0.000	0.690	0.361	0.969	0.219	0.005	0.024	0.423	0.437	0.024
6PGD-2	125.0	0.000	0.825	0.505	0.984	0.047	0.001	0.026	0.649	0.658	0.026
6PGD-3	45.0	0.000	0.678	0.363	0.969	0.387	0.023	0.057	0.710	0.726	0.057
6PGD-4	2.3	0.506	0.675	0.370	0.969	0.472	0.012	0.026	0.101	0.124	0.026
PGI-1	23.1	0.010	0.498	0.265	0.850	0.252	0.003	0.012	0.199	0.209	0.012
PGM-1	13.9	0.030	0.575	0.308	0.923	0.321	0.002	0.008	0.033	0.040	0.008
PGM-2	41.9	0.000	0.683	0.358	0.969	0.330	0.003	0.009	0.359	0.364	0.008
PGM-3	29.3	0.000	0.684	0.370	0.969	0.112	0.007	0.055	0.598	0.620	0.055
SKD-2	2.8	0.096	0.820	0.502	0.984	0.153	0.004	0.023	0.171	0.190	0.023
Total averages <i>L. spartioides</i>						0.158	0.005	0.027	0.294	0.313	0.027
<i>L. holosericeus</i>											
EST-3	28.6	0.000	0.584	0.317	0.927	0.206	0.004	0.017	0.327	0.338	0.017
MDH-1	206.9	0.000	0.589	0.317	0.927	0.114	0.064	0.360	0.805	0.875	0.359
MDH-2	0.0	1.000	0.812	0.503	0.985	0.017	0.000	0.020	0.229	0.245	0.020
MDH-4	132.3	0.000	0.583	0.321	0.927	0.104	0.003	0.031	0.474	0.490	0.031
ME-1	31.3	0.000	0.589	0.317	0.927	0.130	0.004	0.032	0.449	0.466	0.032
6PGD-1	161.3	0.000	0.682	0.360	0.971	0.234	0.027	0.103	0.990	0.991	0.103
6PGD-2	221.7	0.000	0.696	0.362	0.971	0.088	0.038	0.304	1.000	1.000	0.071
6PGD-3	168.5	0.000	0.690	0.370	0.971	0.126	0.005	0.035	0.982	0.983	0.035
6PGD-4	134.0	0.000	0.685	0.371	0.971	0.454	0.027	0.056	0.306	0.345	0.056
PGI-1	133.1	0.000	0.581	0.322	0.914	0.122	0.005	0.040	0.268	0.297	0.040
PGM-1	38.8	0.000	0.502	0.271	0.859	0.273	0.009	0.031	0.184	0.209	0.031
PGM-2	55.7	0.000	0.685	0.369	0.971	0.254	0.012	0.044	0.461	0.485	0.044
PGM-3	19.7	0.000	0.690	0.376	0.971	0.327	0.049	0.131	0.200	0.305	0.131
SKD-1	133.0	0.000	0.818	0.504	0.985	0.011	0.000	0.012	0.406	0.413	0.012
SKD-2	133.2	0.000	0.680	0.372	0.956	0.136	0.009	0.061	-0.021	0.042	0.061
Total averages <i>L. holosericeus</i>						0.155	0.015	0.089	0.543	0.576	0.080
<i>Lotus sp</i>											
EST-3	57.0	0.000	0.539	0.279	0.892	0.197	0.006	0.027	0.406	0.422	0.027
IDH-1	23.5	0.001	0.612	0.326	0.958	0.120	0.010	0.079	0.064	0.138	0.079
MDH-1	132.0	0.000	0.825	0.504	0.991	0.201	0.377	0.653	0.524	0.834	0.652
MDH-2	132.0	0.000	0.714	0.385	0.983	0.109	0.008	0.070	0.940	0.944	0.070
MDH-3	132.0	0.000	0.832	0.503	0.991	0.096	0.007	0.071	1.000	1.000	0.071
MDH-4	370.0	0.000	0.621	0.329	0.958	0.109	0.005	0.046	0.710	0.723	0.046
ME-1	21.9	0.000	0.833	0.501	0.991	0.037	0.003	0.065	0.638	0.661	0.065
6PGD-1	28.9	0.000	0.700	0.370	0.974	0.174	0.196	0.530	0.156	0.604	0.531
6PGD-2	231.0	0.000	0.843	0.503	0.991	0.011	0.000	0.012	0.711	0.715	0.012
6PGD-3	44.4	0.000	0.830	0.502	0.991	0.426	0.045	0.095	0.618	0.654	0.095
6PGD-4	19.0	0.000	0.836	0.504	0.991	0.459	0.022	0.047	0.316	0.354	0.056
PGI-1	43.9	0.000	0.608	0.329	0.949	0.132	0.020	0.134	0.263	0.362	0.134
PGM-1	101.2	0.000	0.535	0.273	0.900	0.466	0.071	0.131	0.245	0.344	0.131
PGM-2	3.0	0.812	0.615	0.327	0.949	0.190	0.382	0.667	0.149	0.717	0.667
PGM-3	0.4	0.515	0.834	0.504	0.991	0.176	0.038	0.178	-0.076	0.116	0.178
SKD-1	76.3	0.000	0.835	0.504	0.991	0.016	0.000	0.016	0.722	0.727	0.016
SKD-2	264.0	0.000	0.613	0.319	0.958	0.120	0.005	0.041	0.373	0.399	0.041
Total averages <i>Lotus sp</i>						0.179	0.070	0.282	0.380	0.556	0.283

SE: standard error; L95 and U95: lower and upper bound of the 95% confidence interval for the average value of Ewens–Watterson statistic. A locus is considered neutral when its average E–W value lies within this interval.

quite low in these taxa. Likewise, Gauthier *et al* (1998) detect a remarkable genetic cohesion in diploid populations of *L. alpinus* ($G_{st} = 0.030$).

By contrast, the average value of $G_{st} = 0.282$ in *Lotus sp.* suggests a high degree of genetic heterogeneity in this assemblage. Mantel tests for isolation by distance between populations were not significant. In fact, some

of the highest genetic distance values (Table 4) were detected between closely located populations. Therefore, sharp geographic discontinuities (eg, deep ravines and cliffs) may be important causes of genetic differentiation between some populations, especially in *Lotus sp.*

Except for some populations of *Lotus sp.*, the average estimated number of migrants per generation is high

within the three assemblages (data not shown), ranking far above the theoretical threshold of one migrant per generation assumed to prevent independent population evolution according to Slatkin (1985, 1987, 1994). These population assemblages could therefore be regarded as single reproductive units that have maintained high levels of variation through abundant genetic interchange. However, the greater genetic distances detected within *Lotus* sp. (see Table 4) suggest that this assemblage may either contain taxonomically distinct entities or is in a more advanced stage of differentiation.

Taxonomic implications

Since the relationship between phenotype and genotype is in general simpler for isozyme evidence than for morphological characters (Gottlieb, 1977), the genetic differences detected might be useful in assessing the taxonomic congruence in these Gran Canarian *Lotus*.

The existence of different taxa is in the first place a question of reproductive isolation. Therefore, if the population assemblages considered for these Gran Canarian endemic *Lotus* are indeed different taxonomic entities, then we should expect reduced levels of gene flow among them. In agreement with this prediction, the low values of G_{st} within *L. holosericeus* and *L. spartioides* are substantially increased when assemblages are compared. This finding indicates incipient reproductive isolation. Thus, the pertinent question to address is whether this degree of genetic heterogeneity is enough to sustain the ascription of these population groups to different taxonomic categories.

The average genetic identities within *L. holosericeus* ($I=0.986$), *L. spartioides* ($I=0.992$) and *Lotus* sp.

($I=0.896$) are within the range reported by Gottlieb (1977) for con-specific populations ($I=0.95 \pm 0.2$), as are those between pairwise combinations of *L. holosericeus*, *L. genistoides* and *L. spartioides* (Table 5). Remarkably, populations LSPPA and LSPNU have higher genetic identities with populations of *L. spartioides* or *L. holosericeus* than with other populations of *Lotus* sp. Removing these two populations from the calculations results in a further decrease of the values (Table 5) to a region between Gottlieb's (1977) averages for con-specific




Figure 2 UPGMA dendrogram based on Nei's (1978) distance values between pairwise combinations of the 14 populations surveyed. Symbols correspond to Figure 1.

Table 4 Values of Nei's (1978) genetic distance between the populations of *Lotus* sampled

Population	<i>L. holosericeus</i>						<i>L. spartioides</i>		<i>Lotus</i> sp.					
	2	3	4	5	6	7	8	9	10	11	12	13	14	
1. LGCJ	0.005	0.003	0.023	0.004	0.006	0.012	0.062	0.059	0.213	0.226	0.079	0.003	0.200	
2. LHAR		0.007	0.023	0.004	0.005	0.019	0.064	0.059	0.208	0.221	0.081	0.007	0.191	
3. LHCU			0.029	0.005	0.011	0.011	0.079	0.071	0.225	0.242	0.098	0.005	0.200	
4. LHPI				0.020	0.019	0.033	0.054	0.056	0.130	0.134	0.066	0.023	0.132	
5. LHPP					0.002	0.009	0.058	0.059	0.181	0.199	0.076	0.006	0.175	
6. LHRO						0.013	0.058	0.059	0.197	0.210	0.081	0.011	0.194	
7. LHTE							0.073	0.064	0.231	0.245	0.091	0.016	0.212	
8. LSBE								0.008	0.121	0.116	0.009	0.047	0.137	
9. LSTA									0.146	0.135	0.020	0.048	0.135	
10. LSPCM										0.012	0.125	0.193	0.027	
11. LSPMA											0.121	0.203	0.045	
12. LSPNU												0.059	0.149	
13. LSPPA													0.189	
14. LSPPN														

Population codes correspond to Table 1.

Table 5 Identities of Nei (1978) between pairs of taxa

	<i>L. spartioides</i>	<i>L. genistoides</i>	<i>Lotus</i> sp	<i>Lotus</i> sp*
<i>L. holosericeus</i>	0.939 (0.007)	0.991 (0.007)	0.876 (0.070)	0.822 (0.030)
<i>L. spartioides</i>		0.941 (0.003)	0.914 (0.050)	0.877 (0.010)
<i>L. genistoides</i>			0.870 (0.088)	0.808 (0.010)
<i>Lotus</i> sp.				0.849 (0.028)

Lotus sp* stands for *Lotus* sp. without the populations LSPPA and LSPNU (see Discussion); Values in brackets are standard deviations.

populations ($I=0.95\pm 0.2$) and congeneric species ($I=0.67\pm 0.07$).

Overall, our results suggest that *L. genistoides*, *L. holosericeus* and *L. spartioides* are very close genetically, and that *Lotus* sp. is a heterogeneous assemblage that contains two populations (LSPNU and LSPPA) much closer to *L. holosericeus* and *L. spartioides* than to other populations designated as *Lotus* sp. A joint consideration of these data with ongoing morphological analyses (Oliva et al, in prep.) will allow us to determine whether the triad *L. holosericeus*, *L. spartioides* and *L. genistoides* should be considered a single taxonomic entity and whether populations LSPCM, LSPMA, LSPPN within the assemblage that we designated *Lotus* sp. are taxonomically distinct.

Acknowledgements

We thank the Cabildo Insular de Gran Canaria (CIGC) for a pre-doctoral research grant to Felicia Oliva and for supporting this investigation. The biologists at the Jardín Botánico Canario 'Viera y Clavijo' provided helpful suggestions and field assistance, especially Bernardo Navarro, Isabel Santana, Blas Vilches, Olga Fernández-Palacios and Carolina Suárez. Roque M López, environment official at the Gobierno de Canarias, helped us locate some of the populations. This paper was written while Juli Caujapé-Castells (JCC) was contracted by the Ministerio de Ciencia y Tecnología (MCYT) and the CIGC. JCC thanks the MCYT and the CIGC for the cofunding of his Ramón y Cajal Project.

References

- Acebes Ginovés JR, Del Arco Aguilar M, García Gallo A, León Arencibia MC, Pérez De Paz PL, Rodríguez Delgado O et al (2001). Pteridophyta & spermatophyta. In: Izquierdo I, Martín JL, Zurita N, Arechavaleta M (eds) *Lista de especies silvestres de Canarias (hongos, plantas y animales terrestres)*. Consejería de Política Territorial y Medio Ambiente: Gobierno de Canarias, pp 98–140.
- Aldridge A, Ortega J (1976). Estudios en la flora macaronésica. *Bot Macaronésica* 2: 9–18.
- Bañares A, Blanca G, Güemes J, Moreno JC, Ortiz S (Eds.) (2003). *Atlas y Libro Rojo De La Flora Vasculare Amenazada De España*. Dirección General de Conservación de la Naturaleza: Madrid.
- Bramwell D (2002). *Lotus*: pico de paloma, corazoncillos. *Aguayro* 228: 42–45.
- Bramwell D, Bramwell ZI (2001). *Flores Silvestres de las Islas Canarias*, 4th edn. Editorial Rueda: Madrid.
- Clayton JW, Tretiak DN (1972). Amine citrate buffer for pH control in starch gel electrophoresis. *J Fish Res Board Can* 29: 1169–1172.
- Cornuet JM, Luikart G (1996). Description and evaluation of two tests for detecting recent bottlenecks. *Genetics* 144: 2001–2014.
- Crawford DJ, Ruiz E, Stuessy TF, Tepe E, Aqueveque P, Gonzalez F et al (2001). Allozyme diversity in endemic flowering plant species of the Juan Fernandez Archipelago, Chile: ecological and historical factors with implications for conservation. *Am J Bot* 88: 2195–2203.
- DeJode DR, Wendel JF (1992). Genetic diversity and origin of the Hawaiian Island cotton, *Gossypium tomentosum*. *Am J Bot* 79: 1311–1319.
- Francisco-Ortega J, Santos-Guerra A, Kim S-C, Crawford DJ (2000). Plant genetic diversity in Canary Islands: a conservation perspective. *Am J Bot* 87: 909–919.
- Gauthier P, Lumaret R, Bédécarrats A (1998). Genetic variation and gene flow in alpine diploid and tetraploid populations of *Lotus* (*L. alpinus* (D.C.) Schleicher/*L. corniculatus* L.): I. Insights from morphological and allozyme markers. *Heredity* 80: 683–693.
- Gottlieb LD (1977). Electrophoresis evidence and plant systematics. *Ann Missouri Bot Gard* 64: 161–180.
- Hamrick JL, Godt MJW (1989). Allozyme diversity in plant species. In: Brown AHD, Clegg MT, Kahler AL, Weir BS (eds) *Plant Population Genetics, Breeding and Germplasm Resources*. Sinauer Associates Inc.: Sunderland, MA. pp 43–63.
- Hansen A, Sunding P (1993). Flora of Macaronesia. Checklist of vascular plants. 4th rev. edn. *Sommerfeltia* 17: 1–295.
- Kunkel G (1969). Aliens to the Canary Flora. *Cuadernos Bot Canaria* 5: 27–44.
- Kunkel G (1977). Endemismos canarios. Inventario de las plantas vasculares endémicas en la provincia de Las Palmas. *ICONA Monografías* 15, 436pp.
- Lewis PO, Whitkus R (1993). *GeneStat-PC version 3.3*. North Carolina State University: Raleigh, North Carolina.
- Luikart G, Cornuet JM (1998). Empirical evaluation of a test for identifying recently bottlenecked populations from allele frequency data. *Conserv Biol* 12: 228–237.
- Mantel N (1967). The detection of disease clustering and a generalized regression approach. *Cancer Res* 27: 209–220.
- Nei M (1973). Analysis of gene diversity in subdivided populations. *Proc Natl Acad Sci USA* 70: 3321–3323.
- Nei M (1978). Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89: 583–590.
- Ortega J (1976). Citogenética del género *Lotus* en Macaronesia. *Bot Macaronésica* 1: 17–24.
- Ortega J (1979). Citogenética del Género *Lotus* en Macaronesia III. Variación en el contenido de glucósidos cianogénicos en *Lotus* de las Islas Canarias y en Macaronesia. *Bot Macaronésica* 5: 9–19.
- Piry S, Luikart G, Cornuet JM (1998). Bottleneck, a program for detecting recent effective population size reductions from allele frequency data. INRA, URLB, Laboratoire de Modélisation et Biologie Evolutive. Montpellier, France.
- Pitard J, Proust L (1908). *Les Iles Canaries*. Flore de L'Archipel: Paris.
- Poggene Version 1.32 (1997). Ag/For Molecular Biology and Biotechnology Centre (University of Alberta) and Center for International Forestry Research. Alberta, Canada.
- Realson J, Grant WF (1988). Evaluation of hypotheses concerning the origin of *Lotus corniculatus* (Fabaceae) using isozyme data. *Theor Appl Genet* 76: 267–276.
- Realson J, Grant WF (1989). An isozyme study in the genus *Lotus* (Fabaceae). Experimental protocols and genetic basis of the electrophoretic phenotype. *Theor Appl Genet* 77: 595–607.
- Rohlf FJ (1998). NTSYS-pc. Numerical Taxonomy and Multivariate Analysis Version 2.02j. Applied Biostatistics Inc.: Setauket, NY.
- Shields CR, Orton TJ, Stuber CW (1983). An outline of general resource needs and procedures for the electrophoretic separation of active enzymes from plant tissue. In: Tanksley SD, Orton TJ (eds) *Isozymes in Plant Genetics Breeding, Part A*. Elsevier Science Publishing Company Inc.: New York, USA. pp 443–468.
- Slatkin M (1985). Gene flow in natural populations. *Annu Rev Ecol Systemat* 16: 393–430.
- Slatkin M (1987). Gene flow and the geographic structure of natural populations. *Science* 236: 787–792.
- Slatkin M (1994). Gene flow and population structure. In: Real LA (ed) *Ecological Genetics*. Princeton University Press: Princeton, NJ. pp 3–17.
- Sunding P (1972). The Vegetation of Gran Canaria – Oslo. I. *Mat - Naturlo Klasse*. Serie. No. 29, 186pp.

- Sunding P (1979). Origins of the Macaronesian flora. In: Bramwell D (ed) *Plants and Islands*. Academic Press: London. pp 13–40.
- Swofford DL, Selander RB (1989). BYOSYS-1: a fortran program for the comprehensive analysis of electrophoretic data in population genetics and systematics. *J Hered* **72**: 281–283.
- Talavera S, Aedo C, Castroviejo S, Herrero A, Romero-Zarco C, Salgueiro FJ et al (2000). *Flora Peninsula Iberica VII (II). Leguminosae (Partim)*. Real Jardín Botánico: Madrid.
- VV AA (2000). Lista Roja de Flora Vascular Española (valoración según categorías UICN). *Conservación Vegetal* **6**(Extra): 11–38.
- Webb PB, Berthelot S (1836–50). Histoire Naturelle des Iles Canaries 3. *Phytographia Canariensis* (2/2):1–496. lam 37–136 B.
- Wright S (1951). The genetical structure of populations. *Ann Eugen* **15**: 323–354.