***Wolbachia* endosymbiont responsible for cytoplasmic incompatibility in a terrestrial crustacean: effects in natural and foreign hosts**

Y. MORET†, P. JUCHAULT & T. RIGAUD*

Université de Poitiers, Laboratoire de Génétique et Biologie des Populations de Crustacés, UMR CNRS 6556, 40 avenue du Recteur Pineau, F-86022 Poitiers Cedex, France

Wolbachia bacteria are vertically transmitted endosymbionts that disturb the reproduction of many arthropods thereby enhancing their spread in host populations. *Wolbachia* are often responsible for changes of sex ratios in terrestrial isopods, a result of the feminization of genotypic males. Here we found that the *Wolbachia* hosted by *Cylisticus convexus* (wCc) caused unidirectional cytoplasmic incompatibility (CI), an effect commonly found in insects. To understand the diversity of *Wolbachia*-induced effects in isopods, wCc were experimentally transferred in a novel isopod host, *Armadillidium vulgare*. wCc conserved the ability to induce CI. However, *Wolbachia* were not transmitted to the eggs, so the capacity to restore the compatibility in crosses involving two transinfected individuals was lost. The feminizing *Wolbachia* hosted by *A. vulgare* was unable to rescue CI induced by wCc. These results showed that *Wolbachia* in isopods did not evolved broadly to induce feminization, and that CI and the feminizing effect are probably due to different mechanisms. In addition, wCc reduces the mating capacity of infected *C. convexus* males, suggesting that the bacteria might alter reproductive behaviour. The maintenance of wCc in host populations is discussed.

Keywords: crustacean, cytoplasmic incompatibility, endosymbiont, isopod, *Wolbachia*.

Introduction

Many invertebrates harbour intracellular bacteria of the genus *Wolbachia* (Werren & O'Neill, 1997; Bandi *et al.*, 1998). These maternally inherited symbionts cause various alterations in the reproduction of their hosts, which can enhance the spread of the infection in arthropod populations (review in Bourtzis & O'Neill, 1998). The most common alteration is cytoplasmic incompatibility (CI). In diploids, CI results in a high rate of embryo mortality when the sperm of infected males fertilizes oocytes that are uninfected, or infected by another bacterial variant (review in Hoffmann & Turelli, 1997). The compatibility is restored in crosses between two individuals infected with the same *Wolbachia* variant. The embryo mortality is caused by abnormalities of mitosis during the development (e.g. Callaini

et al., 1996), probably caused by a *Wolbachia*-induced modification of sperm chromosomes. In compatible crosses, modified paternal chromosomes are rescued by the *Wolbachia* present in the eggs. *Wolbachia* bacteria also induce thelytokous parthenogenesis in a number of parasitoid wasps (Stouthamer *et al.*, 1993), male-killing in some insects (Hurst *et al.*, 1999), and convert genotypic males into phenotypic females in some terrestrial isopods (woodlice) (Martin *et al.*, 1973; review in Rigaud, 1997).

The overall evolution of *Wolbachia* is difficult to understand, considering the diversity of the effects expressed by *Wolbachia* in their different hosts, the discrepancy between host and symbiont phylogenies, and the absence of correlation between *Wolbachia* phylogenetic position and their effects (Rousset *et al.*, 1992; Werren *et al.*, 1995; Van Meer *et al.*, 1999). Horizontal transfers between species explain the distribution of *Wolbachia* (Werren *et al.*, 1995; Vavre *et al.*, 1999), but not the diversity of the effects between hosts. Recent studies argue that both *Wolbachia* strains and host characteristics explain the diversity in effects

*Correspondence. E-mail: thierry.rigaud@univ-poitiers.fr

†Present address: Eidgenössische Technische Hochschule (ETH) Zürich, Experimental Ecology, ETH-Zentrum, NW, CH-8092 Zürich, Switzerland.

observed in insects (Poinot *et al.*, 1998; Hurst *et al.*, 1999). However, data on *Wolbachia* plasticity, i.e. the ability (or not) of a given *Wolbachia* strain to express different effects in different hosts, are still scanty. Most available data are on interspecific transfers between hosts where *Wolbachia* express the same phenotype, and showed that the effects were conserved in the new host (e.g. Rousset & De Strodeur, 1994; Braig *et al.*, 1994; Rigaud & Juchault, 1995). The only transfer made between hosts where *Wolbachia* have two different effects revealed that the transinfected symbionts did not induce reproductive alteration in their new host (Van Meer & Stouthamer, 1999). However, in this case, the new infection was not stably maintained, suggesting that *Wolbachia* were not adapted to their new host.

In terrestrial isopods, *Wolbachia* infection is widespread. Most of the symbionts are closely related and belong to a monophyletic group (Bouchon *et al.*, 1998). Nevertheless, they cause at least two phenotypes: CI in *Porcellio dilatatus* (Legrand *et al.*, 1978; Rousset *et al.*, 1992) and feminization in many other species (Martin *et al.*, 1973; Juchault & Legrand, 1979; Juchault *et al.*, 1994; Rigaud *et al.*, 1999). However, the effect expressed by some *Wolbachia* belonging to the 'isopod clade' is unknown. A better knowledge of the variable effects within this clade and the analysis of phenotype following experimental transinfection between woodlice species would help in understanding of *Wolbachia* evolution in this group. This study was undertaken to elucidate the effect of the *Wolbachia* strain harboured by the woodlouse *Cylisticus convexus* (thereafter called wCc strain, following the nomenclature proposed by Zhou *et al.*, 1998). *Cylisticus convexus* is closely related to woodlice species where feminization is expressed and wCc belong to the clade of feminizing symbionts. However, a previous study suggested that wCc do not induce feminization (Bouchon *et al.*, 1998). Here, we shown that wCc cause CI in *C. convexus*. Experimental trans-specific transfers were done to test whether this CI was conserved in the isopod *Armadillidium vulgare*, which is sensitive to *Wolbachia*-induced feminization by a different strain of *Wolbachia*.

Materials and methods

Wolbachia infection in *Cylisticus convexus*

Nine males and 13 females of *Cylisticus convexus* were found in 1995 at Avanton (Vienne, France) and one male and two females were found in 1998 at Villedaigne (Aude, France). Animals were reared (each sample separated) until gravid females were observed. These gravid females were isolated, allowed to release their offspring, and were then tested for the presence of

Wolbachia. Some offspring of the Avanton sample were used to assess the *Wolbachia* vertical transmission rate; others were kept to maintain the strain. Offspring from Villedaigne were first crossed, and then tested. Presence/absence of *Wolbachia* was tested by PCR assays. Total DNA was extracted from the gonads and the nervous system, according to Bouchon *et al.* (1998). Specific primers for *Wolbachia* (99f-994r, O'Neill *et al.*, 1992) were used to amplify the bacterial 16S rDNA gene and the amplification conditions were as previously described (Rousset *et al.*, 1992). The presence/absence of PCR product determine the presence/absence of *Wolbachia* in the animal tested. In negative samples, a mitochondrial primer set was used to ensure the quality of the host's DNA, as described in Bouchon *et al.* (1998).

Pair crosses (male $\times$ female) of each of the four crossing types were made: u $\times$ u, u $\times$ wCc, wCc $\times$ u, wCc $\times$ wCc, where u denotes uninfected individuals (offspring of females collected at Villedaigne) and wCc infected individuals (from the fourth generation of the Avanton strains). Average weights of females were not significantly different between the u and wCc strains ($68.23 \text{ mg} \pm \text{SE } 1.83$ and $69.79 \text{ mg} \pm 1.36$, respectively; $t = 0.68$, $P = 0.50$), allowing a comparison of their fertility. Mating occurred in small circular plastic boxes (8 cm diameter) at 20°C and under a LD 18:6 photoperiod to promote the onset of reproduction. Woodlice lay their eggs into a ventral incubating pouch (marsupium), where they are incubated for four weeks. The transparency of the marsupium allowed us to estimate the proportion of developing embryos during the third week of development. A total counting of the embryos was nevertheless impossible, as eggs are deposited in several layers in the marsupium and are not transparent. The offspring were isolated from their parents immediately after birth, and completed their development in larger rectangular boxes (26 $\times$ 13 cm). Offspring were sexed and counted eight weeks after birth. The infection status of both sires and mothers was determined by PCR. The inseminate status of the females was verified by checking for sperm in the genital tracts. Statistical analyses were made using JMP 3.2 software (SAS Institute, 1997).

Experimental horizontal transfers

To test the effect of wCc in another genetic context, these symbionts were injected into 30 males and 30 females of uninfected *Armadillidium vulgare*, a woodlouse, which is feminized by its endemic *Wolbachia*. Uninfected *A. vulgare* were from lines reared in the laboratory for 30 years (Nice, Alpes maritimes, France). Injections were made using homogenates of 13

C. convexus females, each from a single iso-female infected line. Ovaries and nerve chords from these females were homogenized in 700 μL Ringer solution and the resulting homogenate was passed through a 1.2- μm pore membrane, which retains cell fragments but allows passage of *Wolbachia* endosymbionts (Rigaud *et al.*, 1991). The cuticle of recipient animals was pierced using a fine needle, and 1 μL of extract was injected using a thin glass needle adapted to a Hamilton syringe. The recipient individuals were reared separately for 24 weeks before mating. The uninfected brothers and sisters of inoculated individuals were used as reference. A control was made, consisting of seven uninfected males injected with Ringer solution only. Such a control was not made in females since previous experiments showed that such an injection had no effect on female reproduction (Rigaud *et al.*, 1991). The same breeding procedure as described above for *C. convexus* was used for *A. vulgare*, but we did not count the number of developing embryos in the marsupium, and we sexed the offspring 12 weeks after birth.

Ten *A. vulgare* females naturally infected with *wAv* (their native feminizing *Wolbachia*) were crossed with males injected with *wCc* (the 10 males were some of those used in the preceding experiment). The females came from a strain collected at Celles-sur-Belle (Deux-Sèvres, France) in 1992 and maintained in the laboratory.

Cytological investigations of embryonic development

The first steps of the embryonic development of *C. convexus* and *A. vulgare* were examined in embryos stained with DAPI (4'-6 diamidine-2-phenylindole dihydrochloride). DAPI forms fluorescent complexes with double-stranded DNA, thereby staining chromosomes, which allows monitoring of mitosis. The staining was adapted to woodlice embryos according to Bressac & Rousset's protocol (1993). Embryos from male *wCc* $\times$ female *u* crosses were removed from the incubating pouch, air dried, and fixed for 5 min in absolute ethanol. The embryos were dried, incubated in DAPI (1 $\mu\text{g}/\text{mL}$) for 15 min, and rinsed with Ringer. The embryos were then placed on slides in 10%

gelatine, and examined under a Zeiss Axoplan microscope equipped for epifluorescence. Embryos from male *u* $\times$ female *u* crosses were checked as controls (from 10 to 20 embryos at each development stage mentioned in Table 3).

Results

Natural *Wolbachia* infection in *Cylisticus convexus*

All females found in Avanton were infected (Table 1). Among the 119 offspring tested, 94.1% harboured *Wolbachia*, showing a high, but not complete, vertical transmission rate of the bacteria (Fisher exact test comparing the infection between males and females showed no significant difference, $P=0.11$). The two females from Villedaigne were not infected by *Wolbachia*. Their offspring tested afterwards confirmed the uninfected status of these lines (Table 1). There was no significant sex ratio deviation from 1:1 in the broods, whatever the infection status of the lineages (Kruskal-Wallis test: $\chi^2_1=1.22$; $P=0.27$, for comparison between populations).

Despite the long duration of breeding experiments (pairs were maintained for at least one month before females laid their eggs), some females were not inseminated, or inseminated in only a single genital tractus (half-inseminated) (Table 2). A logistic regression testing for the effects of male and female infection on the insemination status of females revealed an effect of male infection (Likelihood-Ratio $\chi^2_2=21.43$; $P<0.0001$), but no effect of female infection (L-R $\chi^2_2=0.13$; $P>0.90$), nor of the interaction (L-R $\chi^2_2=0.13$; $P>0.90$). Infected males were therefore less successful at mating, irrespective of the infection status of their mate. A dissection of the males revealed neither abnormalities nor macroparasites in those that did not mate. The broods of non-inseminated or half inseminated females were excluded from further analysis (Table 2).

Among the fully inseminated females, there was a significant difference in the proportion of non-developing embryos in the marsupium according to the crossing type

Table 1 Sex ratios and *Wolbachia* infection in broods of *Cylisticus convexus* females caught gravid at two different locations

Location	Mothers		Offspring						
	<i>N</i>	<i>i</i>	<i>N</i> male	<i>N</i> female	% Male $\pm$ SE	Male <i>i</i>	Male <i>u</i>	Female <i>i</i>	Female <i>u</i>
Avanton	13	13	254	235	52.5 $\pm$ 2.0	55	6	57	1
Villedaigne	2	0	39	42	47.7 $\pm$ 1.8	0	19	0	40

i, infected by *Wolbachia* (positive PCR test); *u*, uninfected by *Wolbachia* (negative PCR test).

Table 2 Crosses between *Cylisticus convexus* and *Armadillidium vulgare* lineages infected or not by *Wolbachia* endosymbionts

Species	Cross type (male × female)	N	N0	N1	N2	Mean proportion of dead embryos per brood ± SE† (embryos counted)‡	Mean number of offspring per brood ± SE† (total offspring)	Mean proportion of males per brood ± SE† (total males)
<i>C. convexus</i>	wCc × wCc	18	6	2	10	0.02 ± 0.01 (193)	23.5 ± 1.6 (235)	0.51 ± 0.03 (118)
	wCc × u	20	8	1	11	0.51 ± 0.05 (230)	15.7 ± 1.7 (173)	0.40 ± 0.05 (78)
	u × wCc	19	0	2	17	0.05 ± 0.02 (299)	22.6 ± 1.3 (385)	0.48 ± 0.02 (188)
	u × u	21	0	2	19	0.02 ± 0.01 (309)	23.8 ± 1.8 (452)	0.49 ± 0.02 (220)
<i>A. vulgare</i>	wCc × wCc	14	0	0	14	—	7.6 ± 3.3 (107)	0.35 ± 0.01 (49)
	wCc × u	14	0	0	14	—	8.7 ± 3.1 (122)	0.45 ± 0.01 (62)
	u × wCc	9	1	0	8	—	108.6 ± 8.1 (869)	0.50 ± 0.00 (432)
	u × u	10	0	0	10	—	92.3 ± 8.8 (923)	0.49 ± 0.00 (446)
	R × u	7	0	1	6	—	88.5 ± 6.4 (531)	0.50 ± 0.01 (266)
	wCc × wAv	10	0	0	10	—	0	—

wAv, infected by *Wolbachia* of *A. vulgare*; wCc, infected by *Wolbachia* of *C. convexus*; u, uninfected; R, males injected with Ringer solution; N, number of crosses; N0, number of females not inseminated; N1, number of females half inseminated; N2, number of females fully inseminated.

† Measures made on fully inseminated females only.

‡ The number of embryos counted was often smaller than the total offspring because not all embryos could be counted in the marsupium (see methods).

(Table 2, Kruskal–Wallis test: $\chi^2_{33} = 31.53$; $P < 0.0001$). Crossing type male wCc × female u showed higher rates of non-developing eggs when compared to all other crossing types (Tukey test, $P < 0.0001$ in all cases), while other crossing types did not differ significantly from each other ($P > 0.80$ for each other comparisons). The result is a lower number of adult offspring from the male wCc × female u crossing type (Table 2, ANOVA: $F_{3,53} = 4.19$; $P = 0.01$; Fisher PLSD post-hoc test: $P < 0.03$ for comparisons between male wCc × female u group with all others; $P > 0.50$ for other comparisons). There was no differences in the sex ratio of broods produced in the different crossing types (Table 2, Kruskal–Wallis test: $\chi^2_3 = 4.67$; $P = 0.20$).

Experimental horizontal transfers of wCc into *Armadillidium vulgare*

Twelve inoculated males and 11 inoculated females were tested by PCR, and were shown to harbour *Wolbachia*. None of the five control males tested was infected. None of the *A. vulgare* recipient males were feminized

20 weeks after inoculation with wCc, and they were therefore used for crosses.

Unlike in *C. convexus*, nearly all females were inseminated (Table 2). The broods of females mated with inoculated males were much smaller than broods sired by uninfected males, whatever the inoculation status of the females (Table 2; Kruskal–Wallis test: $\chi^2_3 = 33.06$; $P < 0.0001$; Tukey post-hoc pairwise test significant [$P < 0.0001$] for the differences between broods from inoculated and uninfected males). No reproductive alteration was observed in crosses involving males injected with Ringer, when compared with crosses with no injection (Table 2; ANOVA testing a difference in the number of offspring: $F_{1,14} = 0.09$; $P > 0.70$). The reproductive alteration was therefore not due to the injection itself, but was correlated with male infection. Some females produced a second brood, and the results were the same as in the first (results not shown).

PCR tests on one male and one female offspring from broods of inoculated females revealed no *Wolbachia* (44 individuals tested). The maximum probability of any

Wolbachia infection in this negative sample is 0.06 according to the estimation method of Post & Millest (1991), and a comparison with the transmission rate of wCc in *C. convexus* showed a highly significant difference (Fisher exact test, $P < 0.0001$). The wCc transmission from transinfected mothers to their offspring was therefore unlikely or at least very low compared to that in the native host. This result is consistent with those obtained previously on experimental transfers in isopods: the *Wolbachia* transmission is inefficient after transfers between phylogenetically distant species. This contrasts with high transmission efficiency when transfers are made within the same species or between closely related species (between 86 and 100% as computed from data in Legrand & Juchault, 1970; Rigaud & Juchault, 1995). *Wolbachia* infection in mothers had no effect on the sex ratio of the broods (Kruskal–Wallis test: $\chi^2_3 = 3.68$; $P > 0.25$), and the slight excess of females in the crosses male wCc $\times$ female wCc (Table 3) was probably a bias due to the small number of offspring.

Ten *A. vulgare* females naturally infected with their feminizing wAv *Wolbachia* were crossed with males injected with wCc (Table 2). None of these naturally infected *A. vulgare* females produced viable embryos, while dissection showed that the males had inseminated the females. Here the incompatibility was therefore total.

Cytological investigations of embryonic development

A total of 211 and 120 embryos from male wCc $\times$ female u crosses were examined in eight crosses of *C. convexus* and four crosses of *A. vulgare*, respectively. In both species, these embryos showed a similar increasing disturbance of mitosis (Table 3). The alterations did not occur during the first mitosis, as it is the case in some insects (Reed & Werren, 1995), but appeared gradually after two or three cycles of cell divisions. The chromosomes did not segregate normally in anaphase, and bridges of DNA were visible between the two sets of dividing chromosomes, often generating parachute-like figures (Fig. 1b). Then, this DNA condensed and dispersed into the cytoplasm (Fig. 1c). A similar CI pattern occurred in embryos of *Drosophila simulans* reported by Callaini *et al.* (1996), where some of the chromosomes did not attach properly to the kinetochore microtubules. These abnormalities were also similar to those observed in *Nasonia* (Reed & Werren, 1995), but at later mitotic stages. No mitotic divisions could be observed in abnormal embryos of *A. vulgare* 50 h after fertilization (stage > 64 cells). In *C. convexus*, death of embryos occurred as late as the segmentation stage (Table 3). In most dead embryos, nuclei were very diffuse, and the cytoplasm became stained with DAPI,

Table 3 Chronology of mitotic disorders during the development of *Cylisticus convexus* and *Armadillidium vulgare* embryos from male wCc $\times$ female u crosses

Species	<i>t</i>	Development in male u $\times$ female u crosses	Development in male wCc $\times$ female u crosses			
			Normal embryos <i>n</i>	Embryos with abnormal divisions (AD) or dead embryos		
				<i>n</i>	Stage of development	% cells with AD (<i>n</i>)*
<i>Cylisticus convexus</i>	12 h	1–2 cells	35	0	—	—
	24 h	2–4 cells	10	5	4 cells	25.0% (20)
	50 h	32–64 cells	8	1	64 cells	4.7% (64)
				11	32 $\leq$ cells $\leq$ 64	38.7% (88)
				2	Cells $<$ 32	54.0% (28)
				24	8 $\leq$ cells $\leq$ 250	n.d.
				33	Cells $>$ 250	n.d.
				2	Cells $>$ 1000	n.d.
				5	Segmentation	n.d.
				3	n.d.	n.d.
<i>Armadillidium vulgare</i>	12 h	1–2 cells	30	0	—	—
	24 h	2–4 cells	2	28	4 cells	35.7% (112)
	60 h	32–64 cells	0	30	20 $<$ cells $\leq$ 64	100% (100)
	25 d	Full embryo	1	29	Cells $\leq$ 64	n.d.

wCc, infected by *Wolbachia* of *C. convexus*; u, uninfected; *t*, time after laying (in hours or days); n.d., not possible to determine due to DNA lysis in the embryos.

*(*n*), number of cells where abnormal divisions were examined.

Fig. 1 Chromosomes (stained with DAPI) during the first steps in the development of woodlice embryos: (a) mitosis (anaphase) in an *Armadillidium vulgare* embryo from the control cross: male u $\times$ female u, 38 h after fertilization ($\times 400$); (b) abnormal mitosis (anaphase) in an *A. vulgare* embryo from the cross: male wCc $\times$ female u, 40 h after fertilization ($\times 400$); (c) abnormal mitosis (anaphase) in a *C. convexus* embryo from the cross: male wCc $\times$ female u, 40 h after fertilization ($\times 400$).

as if the DNA had gradually diffused into the cytoplasm. In control crosses, none of these abnormalities was observed. Furthermore, unfertilized eggs do not develop in isopods (Rigaud, personal observation). Therefore, abnormal mitosis observed in male wCc $\times$ female u crosses cannot be explained by unfertilized eggs in these crosses.

Discussion

The experimental crosses showed that *Wolbachia* of *C. convexus* (wCc) did not cause feminization or male-killing in their natural host. They nevertheless caused embryo mortality in crosses between infected males and uninfected females, due to abnormal mitosis during embryo development. This pattern is similar to the unidirectional cytoplasmic incompatibility (CI) that occurs in several insect species (Hoffmann & Turelli, 1997). This wCc-induced CI was moderate, since only half of the embryos died during their development. The wCc transmission was imperfect but high, and, in compatible crosses, no difference in fertility between infected and uninfected females was detected (our experimental procedure did not allow fecundity estimations). These parameters, showing the infection success of this bacterial strain in its common host, may indicate that the *Wolbachia* infection in *C. convexus* is not recent (Turelli, 1994).

CI-inducing *Wolbachia* found in isopods (both wCc and *Wolbachia* harboured by *Porcellio dilatatus*) belong to the monophyletic group of feminizing *Wolbachia*, a group well separated from those including CI-inducing *Wolbachia* hosted by insects (Bouchon *et al.*, 1998). The presence of a second CI-inducing symbiont in the isopod *Wolbachia* clade indicates that *Wolbachia* infecting isopods have not evolved broadly to induce feminization. Trans-infection experiments in the novel host *A. vulgare* showed that (i) wCc did not induce feminization, (ii) wCc maintained CI expression in a different

genetic context, and (iii) feminizing *Wolbachia* of *A. vulgare* were unable to rescue the CI induced by wCc. This means that different selective pressures have led isopod *Wolbachia* to evolve toward feminization or toward CI, and that the two effects induced by closely related symbionts are due to different mechanisms. Little is known about differences on sex differentiation between species in terrestrial isopods. However, differences exist in the androgenic hormones (a potential target of the *Wolbachia* feminizing effect) between different woodlice species (Martin & Juchault, 1999). This could explain why some symbionts are unable to induce feminization in some species. But differences could also exist in life history traits, e.g. differences in the constraint due to the lack of males in the context of feminization (Hatcher *et al.*, 1999).

In the new *A. vulgare* host, compatibility was not restored in crosses between transinfected males and females. This result may be explained by the very low vertical transmission rate to offspring by transinfected females, i.e. *Wolbachia* may not have reached the eggs, where rescue of compatibility occurs (Hoffmann & Turelli, 1997; Bourtzis & O'Neill, 1998). Therefore, wCc seems to be unadapted to the host *A. vulgare*.

Our results also suggest that wCc lowers mating capacity of infected *C. convexus* males. As the males were kept for a long time with females for mating, and as dissection of these males did not show any evidence of physiological alteration, it is likely that *Wolbachia* cause an alteration in reproductive behaviour. However, a recent study showed that males of *Drosophila simulans* produce less sperm and are less fertile than uninfected ones (Snook *et al.*, 2000). Our study did not allowed sperm counting, but such a screening could be a promising way to extent our observations.

Mating with infected males is required to induce CI in uninfected females. A low mating rate of these males would therefore potentially decrease the spreading

capacity of *Wolbachia* in host populations, especially when the infection is rare. The moderate CI expression and the imperfect *Wolbachia* transmission to offspring would not help the symbiont spreading either (Hoffmann & Turelli, 1997). By using our parameters estimates in Hoffman & Turelli's model (μ , the fraction of uninfected offspring produced by infected females = 0.059; H , the relative hatch rate from incompatible vs. compatible crosses = 0.5; F , the relative fecundity of infected females = 1), we showed that $F(1 - \mu) < 1$, so that drift is required for the infection to be maintained locally. Several aspects of the host population dynamics support maintenance of *C. convexus* infection by drift. Populations are very scattered in France, and animals are generally found in low numbers in the wild (Vandel, 1962) as illustrated by our very small sample sizes. As discussed by Hoffmann & Turelli (1997), drift could favour the fixation of CI-inducing symbionts in small populations, by pushing their frequency above the threshold required for a deterministic spreading to occur. Thus *Wolbachia* maintenance may be favoured in small populations of *C. convexus*, despite moderate CI expression, incomplete vertical transmission to offspring and *Wolbachia* effect on male reproductive success. On the other hand, drift could also easily induce local losses of *Wolbachia* in small populations. Owing to the rare data on wild populations of *C. convexus*, further studies are needed to understand the dynamics and distribution of *Wolbachia* in this isopod.

Acknowledgements

Animals from Villedaigne were collected by Gilbert Martin. Robert Comte provided efficient help with the rearing. Many thanks to Sylvain Charlat and Hervé Merçot for helpful discussions and comments on the manuscript. Cytological observations were made at the SIMIS (Service Interdisciplinaire de Microscopie et d'Imagerie Scientifique), University of Poitiers.

References

- BANDI, C., ANDERSON, T. J. C., GENSHI, G. AND BLAXTER, M. L. 1998. Phylogeny of *Wolbachia* in filarial nematodes. *Proc. R. Soc. B*, **265**, 2407–2413.
- BOUCHON, D., RIGAUD, T. AND JUHAULT, P. 1998. Evidence for widespread *Wolbachia* infection in isopod crustaceans: molecular identification and host feminization. *Proc. R. Soc. B*, **265**, 1081–1090.
- BOURTZIS, K. AND O'NEILL, S. L. 1998. *Wolbachia* infection and arthropod reproduction. *BioScience*, **48**, 287–293.
- BRAIG, H. R., GUZMAN, H., TESH, R. R. AND O'NEILL, S. L. 1994. Replacement of the natural *Wolbachia* symbiont of *Drosophila simulans* with a mosquito counterpart. *Nature*, **367**, 453–455.
- BRESSAC, C. AND ROUSSET, F. 1993. The reproductive incompatibility system in *Drosophila simulans*: DAPI-staining analysis of the *Wolbachia* symbionts in sperm cysts. *J. Invert. Pathol.*, **63**, 226–230.
- CALLAINI, G., RIPARBELLI, M. G., GIORDANO, R. AND DALLAI, R. 1996. Mitotic defects associated with cytoplasmic incompatibility in *Drosophila simulans*. *J. Invert. Pathol.*, **67**, 55–64.
- HATCHER, M. J., TANEYHILL, D. E., DUNN, A. M. AND TOFTS, C. 1999. Population dynamics under parasitic sex ratio distortion. *Theor. Pop. Biol.*, **56**, 11–28.
- HOFFMANN, A. A. AND TURELLI, M. 1997. Cytoplasmic incompatibility in insects. In: O'Neill, S. L., Hoffmann, A. A. and Werren, J. H. (eds) *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*, pp. 42–80. Oxford University Press.
- HURST, G. D. D., JIGGINS, F. M., SCHULENBURG, J. H. G., BERTRAND, D. ET AL. 1999. Male-killing *Wolbachia* in two species of insects. *Proc. R. Soc. B*, **266**, 735–740.
- JUCHAULT, P. AND LEGRAND, J. J. 1979. Analyse génétique et physiologique de la détermination du sexe dans une population du crustacé isopode onisco *Armadillidium nasatum* (Budde-lund). *Arch. Zool. Exp. Gén.*, **120**, 24–43.
- JUCHAULT, P., FRELON, M., BOUCHON, D. AND RIGAUD, T. 1994. New evidence for feminizing bacteria in terrestrial isopods: evolutionary implications. *C. R. Acad. Sci. Paris Life Sciences*, **317**, 225–230.
- LEGRAND, J. J. AND JUHAULT, P. 1970. Modification expérimentale de la proportion des sexes chez les crustacés isopodes terrestres: induction de la thélygène chez *Armadillidium vulgare* Latr. *C. R. Acad. Sci. Paris Series D*, **270**, 706–708.
- LEGRAND, J. J., MARTIN, G. AND ARTAULT, J. C. 1978. Corrélation entre la présence d'un symbiote bactérien dans les ovocytes de *Porcellio dilatatus petiti*, et la stérilité du croisement *P. d. petiti* mâle $\times$ *P. d. dilatatus* femelle. *Arch. Inst. Pasteur Tunis*, **55**, 507–514.
- MARTIN, G. AND JUHAULT, P. 1999. Androgenic hormone specificity in terrestrial isopods (Oniscidea): systematic involvements. *J. Crust. Biol.*, **19**, 684–689.
- MARTIN, G., JUHAULT, P. AND LEGRAND, J. J. 1973. Mise en évidence d'un micro-organisme intracytoplasmique symbiote de l'Onisco *Armadillidium vulgare* L., dont la présence accompagne l'intersexualité ou la féminisation totale des mâles génétiques de la lignée thélygène. *C. R. Acad. Sci. Paris III*, **276**, 2313–2316.
- O'NEILL, S. L., GIORDANO, R., COLBERT, A. M. E., KARR, T. L. ET AL. 1992. 16S rDNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects. *Proc. Natl. Acad. Sci. U.S.A.*, **89**, 2699–2702.
- POINSOT, D., BOURTZIS, K., MARKAKIS, G., SAVAKIS, C. ET AL. 1998. *Wolbachia* transfer from *Drosophila melanogaster* into *D. simulans*: host effect and cytoplasmic incompatibility relationships. *Genetics*, **150**, 227–237.
- POST, R. J. AND MILLEST, A. L. 1991. Sample size in parasitological and vector surveys. *Parasitol. Today*, **7**, 141.

- REED, K. M. AND WERREN, J. H. 1995. Induction of paternal genome loss by the paternal sex ratio chromosome and cytoplasmic incompatibility bacteria (*Wolbachia*): a comparative study of early embryonic events. *Mol. Reprod. Devel.*, **40**, 408–418.
- RIGAUD, T. 1997. Inherited microorganisms and sex determination of arthropod hosts. In: O'Neill, S. L., Hoffmann, A. A. and Werren, J. H. (eds) *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*, pp. 81–101. Oxford University Press.
- RIGAUD, T. AND JUCHAULT, P. 1995. Success and failure of horizontal transfers of feminizing *Wolbachia* endosymbionts in woodlice. *J. Evol. Biol.*, **8**, 249–255.
- RIGAUD, T., SOUTY-GROSSET, C., RAIMOND, R., MOCQUARD, J. P. ET AL. 1991. Feminizing endocytobiosis in the terrestrial crustacean *Armadillidium vulgare* Latr. (Isopoda): recent acquisitions. *Endocytobiosis Cell Res.*, **7**, 259–273.
- RIGAUD, T., MOREAU, J. AND JUCHAULT, P. 1999. *Wolbachia* infection in the terrestrial isopod *Oniscus asellus*: sex ratio distortion and effect on fecundity. *Heredity*, **83**, 469–475.
- ROUSSET, F. AND DE STRODEUR, E. 1994. Properties of *Drosophila simulans* strains experimentally infected by different clones of the bacterium *Wolbachia*. *Heredity*, **71**, 325–331.
- ROUSSET, F., BOUCHON, D., PINTUREAU, B., JUCHAULT, P. ET AL. 1992. *Wolbachia* endosymbionts responsible for various alterations of sexuality in arthropods. *Proc. R. Soc. B*, **250**, 91–98.
- SNOOK, R. S., CLELAND, S. Y., WOLFNER, M. F. AND KARR, T. L. 2000. Offsetting effects of *Wolbachia* infection and heat shock on sperm production in *Drosophila simulans*: analyses of fecundity, fertility and accessory gland proteins. *Genetics*, **155**, 167–178.
- STOUTHAMER, R., BREEUWER, J. A. J., LUCK, R. F. AND WERREN, J. H. 1993. Molecular identification of microorganisms associated with parthenogenesis. *Nature*, **361**, 66–68.
- TURELLI, M. 1994. Evolution of incompatibility-inducing microbes and their hosts. *Evolution*, **48**, 1500–1513.
- VANDEL, A. 1962. *Faune de France*. 64, isopodes terrestres. P. Chevalier, Paris.
- VAN MEER, M. M. M. AND STOUTHAMER, R. 1999. Cross-order transfer of *Wolbachia* from *Muscidifurax uniraptor* (Hymenoptera: Pteromalidae) to *Drosophila simulans* (Diptera: Drosophilidae). *Heredity*, **82**, 163–169.
- VAN MEER, M. M. M., WITTEVELDT, J. AND STOUTHAMER, R. 1999. Phylogeny of the arthropod endosymbiont *Wolbachia* based on *wsp* gene. *Insect Mol. Biol.*, **8**, 399–408.
- VAVRE, F., FLEURY, F., LEPETIT, D., FOUILLET, P. ET AL. 1999. Phylogenetic evidence for horizontal transmission of *Wolbachia* in host–parasitoid associations. *Mol. Biol. Evol.*, **16**, 1711–1723.
- WERREN, J. H. AND O'NEILL, S. L. 1997. The evolution of heritable symbionts. In: O'Neill, S. L., Hoffmann, A. A. and Werren, J. H. (eds) *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*, pp. 1–41. Oxford University Press.
- WERREN, J. H., ZHANG, W. AND GUO, L. R. 1995. Evolution and phylogeny of *Wolbachia*: reproductive parasites of arthropods. *Proc. R. Soc. B*, **261**, 55–71.
- ZHOU, W. F., ROUSSET, F. AND O'NEILL, S. L. 1998. Phylogeny and PCR based classification of *Wolbachia* strains using *wsp* gene sequences. *Proc. R. Soc. B*, **265**, 509–515.