

CORRIGENDUM

Pharmacokinetics of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in Seveso adults and veterans of operation Ranch Hand

JOEL E. MICHALEK, JAMES L. PIRKLE, LARRY L. NEEDHAM, DONALD G. PATTERSON Jr.,
 SAMUEL P. CAUDILL, RAM C. TRIPATHI AND PAOLO MOCARELLI

Journal of Exposure Analysis and Environmental Epidemiology (2002) 12, 44–53 DOI: 10.1038/sj/jea/7500201

Since the publication of the above paper, the authors have identified an error in Table 3. A revised version of the table is given below.

Table 3. TCDD concentrations in the Seveso cohort.

	TCDD concentration/years since exposure			
	0	0.27	3	16.35
(a) Males				
	(<i>n</i> =20)	(<i>n</i> =11)	(<i>n</i> =17)	(<i>n</i> =24)
Mean	2833.3	2243.7	1311.4	195.4
Standard deviation	2404.3	1574.7	1057.8	160.1
Percentiles				
0	77	358	96	30.7
5	103.5	358	96	33.7
50	2500	2010	1170	119.5
95	7235	5910	3810	488
100	9140	5910	3810	531
(b) Females				
	(<i>n</i> =21)	(<i>n</i> =12)	(<i>n</i> =15)	(<i>n</i> =25)
Mean	2266.7	1622.2	1009.7	354.4
Standard deviation	1909.0	1241.2	670.1	297.7
Percentiles				
0	103	375	97.2	35.4
5	145	375	97.2	46.1
50	1770	1515	933	271
95	5730	4410	2400	871
100	6320	4410	2400	878
(c) All				
	(<i>n</i> =41)	(<i>n</i> =23)	(<i>n</i> =32)	(<i>n</i> =49)
Mean	2543.1	1919.4	1170	276.5
Standard deviation	2156.4	1413.5	896.5	240.5
Percentiles				
0	77	358	96	30.7
5	130	375	97.2	35.4
50	1940	1720	1045	218
95	5730	4410	3130	833
100	9140	5910	3810	878