

- 26 Carritt B, Litt M: Report of the committee on the genetic constitution of chromosomes 20 and 21. *Cytogenet Cell Genet* 1989;51:358-371.
- 27 Korenberg JR, Kawashima H, Pulst SM, Allen L, Magenis E, Epstein CJ: Down syndrome: Toward a molecular definition of the phenotype. *Am J Med Genet* 1990;suppl 7:91-97.
- 28 Rao VN, Papas TS, Reddy SP: ERG, a human ETS-related gene on the chromosome 21: Alternative splicing, polyadenylation and translation. *Science* 1987;237:635-639.
- 29 Créte N, Gosset P, Théophile D, Duterque-Coquillaud M, Blouin JL, Vayssettes C, Sinet PM, Créau-Goldberg N: Mapping the Down syndrome chromosome region. Establishment of a YAC contig spanning 1.2 megabases. *Eur J Hum Genet* 1993;1:51-63.
- 30 Boulukos KE, Pognonec P, Begue A, Galibert F, Gesquière JC, Stéhelin D, Ghysdael J: Identification in chickens of an evolutionarily conserved cellular ets-2 (c-ets2) encoding nuclear proteins related to the products of the c-ets2 protooncogene. *EMBO J* 1988;7:697-705.
- 31 Landsman D, McBride OW, Soares N, Crippa MP, Srikantha T, Bustin M: Chromosomal protein HMG-14. *J Biol Chem* 1989;264:3421-3427.
- 32 Horisberger MA, Wathelet M, Szpirer J, Szpirer C, Islam Q, Levan G, Huez G, Content J: cDNA cloning and assignment to chromosome 21 of IFI-78K gene, the human equivalent of murine Mx gene. *Somat Cell Mol Genet* 1988;14:123-131.
- 33 Patterson D: Report of the second international workshop on human chromosome 21. *Cytogenet Cell Genet* 1991;57:167-174.
- 34 Petersen MB, Tranebjaerg L, McCormick MK, Michelsen N, Mikkelsen M, Antonarakis SE: Clinical, cytogenetic, and molecular genetic characterization of two unrelated patients with different duplications of 21q. *Am J Med Genet* 1990;suppl 7:104-109.
- 35 Williams CA, Frias JL, McCormick MK, Antonarakis SE, Cantu ES: Clinical, cytogenetic, and molecular evaluation of a patient with partial trisomy 21 (21q11-q22) lacking the classical Down syndrome phenotype. *Am J Med Genet* 1990;suppl 7:110-114.
- 36 Daniel A: Normal phenotype and partial trisomy for the G positive region of chromosome 21. *J Med Genet* 1979;16:227-229.
- 37 Shabtai FS, Schwartz A, Klar D, Hart J, Dar H, Kessler E, Halbrecht I: Free proximal trisomy 21 in the mother and malformation syndrome in the son. *Am J Med Genet* 1990;suppl 7:182-185.
- 38 Hagemeyer A, Smit MEE: Partial trisomy 21. Further evidence that trisomy of band 21q22 is essential for Down's phenotype. *Hum Genet* 1977;38:15-23.
- 39 Leschot N, Slater RM, Joenje H, Becker-Bloemkolk MJ, de Nef JJ: SOD-A and chromosome 21. Conflicting findings in a familial translocation (9p24;21q21.4). *Hum Genet* 1981;57:220-223.
- 40 Poissonnier M, Saint-Paul B, Dutrillaux B, Chassaingne M, Gruyer P, de Blignières-Strouk G: Trisomie 21 partielle (21q21→21q22.2). *Ann Génét* 1976;19:69-73.
- 41 Lejeune J, Berger R, Vidal OR, Réthore MO: Un cas de translocation G ~ G en tandem. *Ann Génét* 1965; 8:9-11.

Delabar/Theophile/Rahmani/
Chettouh/Blouin/Prieur/Noel/Sinet

Down Syndrome Map

Erratum

The authors of the paper entitled 'A MERRF/MELAS Overlap Syndrome Associated with a New Point Mutation in the Mitochondrial DNA tRNA^{Lys} Gene' (1993;1:80-87) ask us to correct the following error in the legend of figure 3b (p. 85): the mutation should be at position 8356 (*not* at position 3260).