

Full-length article

Gene expression profile induced by oral administration of baicalin and gardenin after focal brain ischemia in rats¹

Zhan-jun ZHANG², Zhong WANG³, Xiao-yan ZHANG⁴, Kang YING⁵, Jian-xun LIU³, Yong-yan WANG⁶

²The Clinical School of Beijing University of Traditional Chinese Medicine, Beijing 100029, China; ³Xi-yuan Hospital of the Academy of Traditional Chinese Medicine, Beijing 100089, China; ⁴Beijing Normal University, Beijing 100036, China; ⁵BioStar Genechip Incorporated, Shanghai 200012, China; ⁶The Academy of Traditional Chinese Medicine, Beijing 100059, China

Key words

microarray; gene expression; baicalin; gardenin; brain ischemia

¹ Project supported by the Major State Basic Research Development Program of the People's Republic of China (No G19990554).

² Correspondence to Dr Zhan-jun ZHANG.
Phn 86-10-6287-5599, ext 6114.
Fax 86-10-6287-4049.
E-mail zhangzhanjun1972@yahoo.com.cn

Received 2004-06-29

Accepted 2004-11-19

doi: 10.1111/j.1745-7254.2005.00051.x

Abstract

Aim: To investigate differential gene expression and the pharmacological mechanism of baicalin and gardenin in focal cerebral ischemia in rats with high-density cDNA microarray. **Methods:** Rat left middle cerebral arteries were occluded and treated with either baicalin or gardenin. The pharmacological effects were investigated using the difference in infarction areas before and after treatment, which were determined by 2,3,5-triphenyltetrazolium chloride (TTC) staining. Gene expression was demonstrated using a "Biostar40S" gene microarray. Semiquantitative reverse transcription-polymerase chain reaction (RT-PCR) was used to verify the result of the selected genes. **Results:** Both baicalin and gardenin reduced the infarction areas in focal cerebral ischemia rats ($P < 0.05$). The differential genes were 211, 177, and 70 (upregulated or downregulated) in the model group, baicalin, and gardenin treatment groups compared with the sham-operated group, respectively. Gene expression of Rpl19 and Csnk2 underwent an approximately 1.9 and 2.1-fold increase, respectively, verified by semiquantitative RT-PCR, which was the same trend as the cDNA microarray. **Conclusion:** Differential gene expression with respect to the pharmacological effects of baicalin and gardenin on focal cerebral ischemia by cDNA microarray revealed a number of clues with respect to the therapeutic mechanisms of Chinese traditional medicine. In addition, the present study provided theoretical and experimental evidence that will aid future studies examining cerebral ischemia.

Introduction

Stroke, mostly caused by cerebral ischemia, is a multifactorial disease involving the activation of myriads of death inducers, which leads to injury of neurons by eliciting cascades of signal transduction pathways^[1]. The Qing Kai Ling injection (QKLI) is a modified preparation of the "An Gong Niu Huang" pill, a famous traditional Chinese medicament. In recent studies, QKLI showed strong therapeutic effects on stroke in clinical usage. However, this preparation is a compound comprising of many components. Its therapeutic mechanism is very complex, with regard to both individual and integrative effects. Baicalin and gardenin are two effective compounds of QKLI. Baicalin, known as an antioxidant flavonoid *in vitro*, functions as a biological response modi-

fier^[2,3], and has been investigated for its neuroprotective effects against glutamate/*N*-methyl-*D*-aspartate (Glu/NMDA) stimulation and glucose deprivation in primary cultured rat brain neurons. Baicalin was found to significantly reduce Glu/NMDA-increased lactate dehydrogenase (LDH) release^[4]. Baicalin is also a potent inhibitor of endothelial cell proliferation, migration, and differentiation^[5]. Gardenin can significantly decrease the content of transcription factor monocyte chemoattractant protein-1 (MCP-1) in rat brains suffering focal ischemia^[6]. cDNA microarray possesses the ability to analyze the expression changes of hundreds, thousands, or even tens of thousands of genes simultaneously. Several waves of gene expression have been observed after focal cerebral ischemia^[7]; however, a comparative study examin-

ing the pharmacological mechanisms of different drugs has not been reported in the literature. Results from the present study suggest that baicalin and gardenin play a pharmacological role in focal cerebral ischemia by regulating gene expression.

Materials and methods

Cerebral ischemia Forty-four Sprague-Dawley rats (250–280 g, supplied by Beijing Weitong-Lihua Experimental Animal Center, Beijing, China) were randomly divided into four groups: model, baicalin-treatment, gardenin-treatment, and sham-operated group, respectively. Under isofurane anesthesia, rats were subjected to occlusion of the left middle cerebral artery (MCAO) using an intraluminal filament as described by Haruo *et al*^[8]. After 24 h of focal cerebral ischemia, the rats were deeply re-anesthetized using halothane and the left half of the brain was removed for total RNA isolation. In the sham-operated rats, all procedures except for MCAO were carried out. Baicalin or gardenin (National Institute for the Control of Pharmaceutical and Biological Products, Beijing, China) was given orally (40 mg/kg) 2 h after MCAO.

Measurement of cerebral infarction areas Eight rats from each group were used to calculate the infarction ratio. In brief, the cerebrum was removed and cut into five slices on the coronal section at the location from which the distance to the prefrontal is 1, 3, 5, and 7 mm, respectively. The slices were stained in 4% 2,3,5-triphenyltetrazolium chloride (TTC) solution for 10 min^[9]. Images of the slices were captured using a digital camera (Color CCD camera TP-6001A, Topica Inc, Japan). We calculated the areas of the infarction region using a Pathology Image Analysis System (Topica Inc, Japan). The ratio of the infarction area to the total slice area was calculated.

Total RNA isolation The left half of the brain from the remaining three rats in each group was carefully dissected out from the re-anesthetized rats under RNAase-free conditions. The total RNA in each sample was extracted with Trizol reagent (Invitrogen, San Diego, CA, USA) using the procedure outlined in the manufacturer's protocol.

Microarrays Hybridization was carried out on the rat "Biostar40S" gene microarray (BioStar Genechip Incorporated, Shanghai, China), which contained 4096 elements. The array provided a broad overview of the rat genome, which was added to the high-quality array cDNA libraries, and the sequence information, maps, and expression data were placed into the public domain (NCBI, Nucleotide). All clones used for production of the microarrays

were sequence verified. Three pieces of total RNA from the same group were pooled following transcription into cDNA and were labeled with Cy5 and Cy3, respectively, and hybridized on the array. Image files were processed using the Axon GenePix 4000B scanner (Axon, USA) and datasets were prepared according to the routine procedures using Genepix 4.0 software (Axon, USA).

Data analysis Data sheets derived from the results of the Genepix 4.0 analysis were further evaluated using Excel software (Microsoft, USA). The microarrays were hybridized in triplicate and each measurement containing the extracted total RNA from each group was used to analyze each gene, which had three data points of relative changes. Thus, a balance coefficient was calculated to correct variations resulting from unequal amounts of fluorescent dye fade; the average signal from all elements in the Cy3 channel was divided by the average signal from all elements in the Cy5 channel, which resulted in the balance coefficient. The Cy5 signal for each element was then multiplied by the balance coefficient, prior to calculating the expression ratio (Cy3/Cy5). A fold value of =2.0 or =-2.0 indicated that differences in the Cy3: Cy5 ratio were detected at a 99% confidence level (information provided by Incyte, USA). All probes fulfilling the criterion (=2.0 or =-2.0) were compiled^[10–13].

Semiquantitative RT-PCR The procedures for isolating total RNA using Trizol reagent (Invitrogen, Life Technology, Incorporated, New York, NY, USA) and synthesizing first strand cDNA were exactly as previously described^[14]. The optimal polymerase chain reaction (PCR) amplification conditions and cycle number were determined experimentally to ensure specific signal generation. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was considered to be stably expressed after focal cerebral ischemia acted as an interference gene. All amplifications were carried out on a Gradient PCR System (Biometra, USA). cDNA synthesis was carried out using 200 ng of mRNA with 5×10^6 U/L reverse transcriptase (TaKaRa, Japan) in a total 100 μ L reaction mixture and incubated for 60 min at 48 °C. The primer pairs were for ribosomal protein L19 (Rp19): AACAGATCAAGGAGC TGA TCA AGA; AGT CTT GAT GAT CTC CTC CTT CTT (NM 031103); casein kinase II beta subunit (Csnk2b): CGG ACATAAAGATGAGTAGCTCTGA; GTGGTG CCTAGA GGA CTT GGG TGT G (NM 031021); GAPDH: ACC ACA GTC CAT GCC ATC AC; TCC ACC ACC CTG TTG CTG TA (NM 008084.1). After heating to 95 °C for 5 min, each RT reaction mixture was used for PCR amplification. The PCR mixture started with in an initial step of 5 min at 95 °C followed by 35 cycles of 30 s at 95 °C and an annealing temperature of 1 min at 64.5 °C for Csnk2b mRNA, at 57.4 °C for

Rpl19 mRNA and GAPDH, followed by 72 °C for 10 min. Products of 5 μL from each PCR mixture were dyed with 1 μL SYBR Green I Nucleic Acid Gel stain (Cambrex Bio Science Rockland Incorporated, USA) and loaded on 2.5% agarose gels, which were evaluated according to the intensity differences of image electrophoresis using Gel Analyzing Imager (FuRi, Shanghai, China).

Results

TTC staining Both baicalin and gardenin reduced the infarction areas in cerebral ischemia rats by 6%–7% compared with the model group. The ratio of differences was tested using the Student *t*-test (Figure 1, Table 1).



Figure 1. Cross-sections showing cerebral infarction areas (in white) of a number of samples stained with 2,3,5-triphenyltetrazolium chloride.

Table 1. Cerebral infarction areas from a number of samples stained with 2,3,5-triphenyltetrazolium chloride. *n*=8. Mean±SD. ^b*P*<0.05 vs the model group.

Group	Percentage of infarcted area to cerebral area/%
Model	24.7±5.6
Baicalin	14.7±3.0 ^b
Gardenin	15.4±3.2 ^b

Changes in gene expression after MCAO occlusion

Genes were classified into 12 categories by the International General Principle (www.geneontology.org). One hundred and ninety-nine genes were significantly upregulated which are involved in metabolism, signal transduction, cell organization, response to stress, cell adhesion, transport, apoptosis, and a variety of other processes. Another 12 genes showed downregulated expression under the same conditions (Table 2).

Metabolism The largest increases in gene expression were shown by genes involved in protein synthesis: the ribosomal proteins L10, S5, S3a, S24, S6, S11, and L19, glyco-

protein 38, eukaryotic translation elongation factor 1 alpha and actin-related protein complex 1b. The expression of genes related to cell metabolism changed with increased expression of coenzyme A dehydrogenase, procollagen-lysine, lactate dehydrogenase and ubiquinone oxidoreductase subunit B13.

Cell organization and adhesion Cell organization and adhesion genes also showed prominent changes in expression. These included Fibronectin 1, Integrin, H2A histone family member and apolipoprotein M.

Signal transduction Increases in expression were shown by genes involved in cellular signal transduction: CDK5, pyruvate kinase 3, phosphatidylinositol 4-kinase, and casein kinase II. In addition, the G protein pathway suppressor showed decreased expression over this interval.

Response to stress Several “stress” genes involved in cellular responses to inflammation and injury were induced, including Cd63 antigen, MRC OX-45 surface antigen and immunoglobulin superfamily member. Another induced gene, glutathione peroxidase 1, was also upregulated after focal cerebral ischemia.

Apoptotic genes The expression of several apoptotic effector genes was altered. These included programmed-cell-death-8, Urmodulin, and tumor protein translationally controlled genes.

Effect of baicalin on gene expression Increases in gene expression were observed in 89 genes. These included genes involved in metabolism, signal transduction, cell organization, responses to stress, and transcription regulators. In addition to the induced genes, 88 genes simultaneously showed decreased expression (Table 3).

Metabolism Metabolism-related genes showed prominent changes in expression. These included ADP-ribosylation factor, enolase, adenine phosphoribosyltransferase, and palmitoyl-protein thioesterase. In addition, histamine *N*-methyltransferase simultaneously showed decreased expression.

Signal transduction The largest increases in expression were shown by genes involved in protein tyrosine phosphatase receptor type D and receptor type A. S100 calcium-binding protein A9 also showed prominent changes. In addition, the expression of protein kinase C-binding protein Zeta and surfactant-associated protein decreased. Protein kinase is controlled by specific binding proteins, which are believed to sequester each type of kinase to the region of a neuron, such as the postsynaptic specialization or cell nucleus, that requires its function^[15]. The protein kinase C-binding protein Zeta is one of these proteins. Arachidonic acid epoxygenase, an adapter protein of the prostaglandin

Table 2. Partial differential genes in the model compared with the sham-operated group.

Accession number	Average microarray ratio	Definition	Classification
NM_024151	2.369	ADP-ribosylation factor 4	Metabolism
NM_012898	2.293	alpha-2- <i>HS</i> -glycoprotein	Metabolism
AW918082	3.201	Deoxyribose-phosphate aldolase	Metabolism
NM_012576	2.313	Nuclear receptor subfamily 3	Signal transduction
BE113312	2.127	Adapter-related protein complex 3 beta 1 subunit	Signal transduction
X79328	2.153	CaBP1 mRNA	Signal transduction
AI407932	3.301	Retinoid-inducible serine carbopeptidase precursor	Signal transduction
AF335277	2.371	Centrosomal protein centrin 3 mRNA	Cell cycle
NM_017082	2.327	Urmodulin (Tamm-Horsfall protein)	Apoptosis
AW918198	2.295	Myosin heavy chain mRNA	Cell mobility
L24897	3.528	TROPOMYOSIN BETA CHAIN	Cell mobility
NM_031509	3.132	Glutathione-S-transferase (Mu2)	Response to stress
NM_053440	2.908	Superiorcervical ganglia, neural specific 10	Others
AW915152	4.633	Hypothetical protein C32E8.9	Others
BF388747	2.172	ESTs	Others
AA893226	2.198	ESTs	Others
BI276980	2.485	ESTs	Others
BF391240	2.248	ESTs	Others
AI639155	2.512	ESTs	Others
BF556197	3.577	ESTs	Others
NM_031797	2.591	Suppression of tumorigenicity	Others
AW918368	3.391	Hypothetical protein	

and leukotriene family of intracellular messengers, also appears to play an important role in the regulation of signal transduction in the brain and elsewhere^[14].

Cell organization The expression of nucleolar phosphoprotein p130, an adapter protein that participates in nucleolar disassembly and cell cycle, decreased after MCAO. In addition, peroxisomal membrane protein showed increased expression over this interval.

Transcription regulator Two genes related to the transcription regulator were altered. These were spliceosome-associated protein and DNA primase.

Effect of gardenin on gene expression Increases in gene expression were observed for 68 genes. These included genes involved in metabolism, signal transduction, cell organization, response to stress, cell cycle, and cell mobility. In addition, two genes showed decreased expression over this interval (Table 4).

Metabolism The expression of several genes involved in protein synthesis and cell metabolism was altered. These included ADP-ribosylation factor, HS-glycoprotein and deoxyribose-phosphate aldolase.

Signal transduction Changes in the expression of genes indirectly related to signal transduction were observed, and consisted of increased expression in nuclear receptor,

adapter-related protein complex and retinoid-inducible serine carbopeptidase precursor.

Response to stress Several "stress" genes involved in cellular responses to oxidation were induced. The most striking was the increased expression of glutathione-S-transferase.

Cell mobility Myosin heavy chain and tropomyosin beta chain genes that participate in cell mobility and cell cycle were also induced.

Semiquantitative RT-PCR The outcome of this study showed that the expression of ribosomal protein RpL19 mRNA and Csnk2 mRNA were consistently upregulated in the individual samples from MCAO rats relative to sham-operated rats. Expression increased by 1.9- and 2.1-fold, respectively, and showed the same trend as the results of the microarray. The outcome of RT-PCR indicated that broad, array-based gene expression measurements were reliable for determining gene expression patterns in the brain (Figure 2).

Discussion

Messenger RNA is only an intermediate on the way to the production of the eventual protein products. In the present study we explored the potential role of cDNA

Table 3. Partial differential genes in the model compared with the baicalin-treatment group.

Accession number	Average microarray ratio	Definition	Classification
NM_019156	0.222	Vitronectin	Others
NM_022399	0.383	Calreticulin	Others
NM_023986	0.472	TEMO	Others
NM_031140	2.317	Vimentin	Others
NM_053372	3.058	Secretory leukocyte protease inhibitor	Others
NM_012559	3.359	Fibrinogen, gamma polypeptide	Others
NM_017131	2.848	Calsequestrin 2	Others
BI285007	0.271	Spliceosome-associated protein	Transcription regulator
U67994	0.441	DNA primase small subunit mRNA	Transcription regulator
NM_022869	0.327	Nucleolar phosphoprotein p130	Cell organization
NM_017234	3.408	Peroxisomal membrane protein 3	Cell organization
NM_031559	0.365	Carnitine palmitoyltransferase 1 alpha	Metabolism
NM_024152	0.402	ADP-ribosylation factor 6	Metabolism
NM_012554	0.405	Enolase 1, alpha	Metabolism
L04970	0.408	Adenine phosphoribosyltransferase	Metabolism
NM_022502	0.491	Palmitoyl-protein thioesterase	Metabolism
NM_031044	2.921	Histamine N-methyltransferase	Metabolism
[1609]	0.264	Protein kinase C-binding protein Zeta	Signal transduction
NM_017329	0.436	Surfactant-associated protein 1	Signal transduction
NM_053587	3.054	S100 calcium-binding protein A9	Signal transduction
NM_031839	4.286	Arachidonic acid epoxygenase	Signal transduction
NM_019140	7.844	Protein tyrosine phosphatase receptor type D	Signal transduction
NM_012763	6.988	Protein tyrosine phosphatase receptor type A	Signal transduction
NM_031140	2.317	Vimentin	Response to stress
NM_053372	3.058	Secretory leukocyte protease inhibitor	Response to stress
NM_012559	3.359	Fibrinogen, gamma polypeptide	Response to stress

microarrays for gene expression analysis after focal brain ischemia and examined the differences in gene expression after the action of different compounds of QKLI. An ischemia period of 24 h was used because this is the maximum period of cerebral ischemia that is compatible with neuroprotection^[16-18]. To select results from microarray ex-

periments with reliably altered ratios, we filtered the results using two criteria^[10-13]: minimal fold-change values, and ratios reproducibly different from unity. The hybridization ratio had to be at least two-fold higher or lower than the control groups. Of these filters, the two-fold ratio filter was by far the most restrictive and there was no uniform standard across the different platforms of the arrays. We have listed the information of differential gene expression according to the functional classification of genes. In brief, baicalin and gardenin appear to have a profound influence on the model by regulating the expression of different genes or by acting on different metabolic pathways. The genes related to cell metabolism presented striking changes, and showed increased expression in both model and baicalin-treatment groups in contrast to the gardenin-treatment group. Baicalin appears to have a more prominent action on signal transduction in cells than gardenin, including activation of ion channels, regulation of kinase and phosphorylation of receptor proteins. In contrast, the effects of gardenin in stress responses and anti-oxidation appear to be more significant




Figure 2. Some strongly regulated Rpl19 and Csnk2 genes are confirmed with RT-PCR. Size of the internal control is 550 bp, and that of the target genes are 350 and 320 bp. Ratios indicate the signal intensity of examined gene vs that of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) at different groups. The bottom panel in each graph is a representative gel of the PCR products of three independent repeats.

Table 4. Partial differential genes in the model compared with the gardenin-treatment group.

Accession number	Average microarray ratio	Definition	Classification
NM_019156	0.222	Vitronectin	Others
NM_022399	0.383	Calreticulin	Others
NM_023986	0.472	TEMO	Others
NM_031140	2.317	Vimentin	Others
NM_053372	3.058	Secretory leukocyte protease inhibitor	Others
NM_012559	3.359	Fibrinogen, gamma polypeptide	Others
NM_017131	2.848	Calsequestrin 2	Others
BI285007	0.271	Spliceosome-associated protein	Transcription regulator
U67994	0.441	DNA primase small subunit mRNA	Transcription regulator
NM_022869	0.327	Nucleolar phosphoprotein p130	Cell organization
NM_017234	3.408	Peroxisomal membrane protein 3	Cell organization
NM_031559	0.365	Carnitine palmitoyltransferase 1 alpha	Metabolism
NM_024152	0.402	ADP-ribosylation factor 6	Metabolism
NM_012554	0.405	Enolase 1, alpha	Metabolism
L04970	0.408	Adenine phosphoribosyltransferase	Metabolism
NM_022502	0.491	Palmitoyl-protein thioesterase	Metabolism
NM_031044	2.921	Histamine <i>N</i> -methyltransferase	Metabolism
[1609]	0.264	Protein kinase C-binding protein Zeta	Signal transduction
NM_017329	0.436	Surfactant-associated protein 1	Signal transduction
NM_053587	3.054	S100 calcium-binding protein A9	Signal transduction
NM_031839	4.286	Arachidonic acid epoxygenase	Signal transduction
NM_019140	7.844	Protein tyrosine phosphatase receptor type D	Signal transduction
NM_012763	6.988	Protein tyrosine phosphatase receptor type A	Signal transduction
NM_031140	2.317	Vimentin	Response to stress
NM_053372	3.058	Secretory leukocyte protease inhibitor	Response to stress
NM_012559	3.359	Fibrinogen, gamma polypeptide	Response to stress

than those of baicalin.

Changes in gene expression after MCAO Prominent changes were recorded in the ribosomal proteins S6, L6, S3a, S24, L5, L10, L19, and S11 on the microarray^[19]. In general, this finding supports previous studies^[20], perhaps reflecting recovery from an early postischemic transcriptional defect. Integrin is a transmembrane protein, and the adhesion effects between leucocytes and cerebral microvessel cells mediated by integrin participated in the injury and destruction of inflammation factors to tissues. The increased expression of integrin after MCAO is, in general, in line with a preceding report^[21]. The increased expression of protein kinase (PK) is mostly caused by the large amount of PK released as a result of metabolic dysfunction, necrosis, ischemia and hypoxia of cells, which occurs in acute cerebrovascular diseases, and this finding was in accordance with the clinical diagnosis. The increased expression of genes in the G protein pathway suppressor 1 may have contributed to the adaptive modulation of the body. In general, G protein plays a role in signal amplification and in switching the molecule on and off in the

course of signal transduction. G protein pathway suppressor was upregulated after focal cerebral ischemia, which might occur in compensation for stress and signal transduction regulation in the body^[22,23]. Casein kinase II is a necessary substance for cell survival, and increased expression of casein kinase II is correlated with hyperplasia and the proliferation of cells after cerebral ischemia. Cyclin-dependent kinase 5 (CDK5) mRNA was found to be downregulated after focal brain ischemia, whereas variable changes were noted at the protein level in focal ischemia^[24]. Downregulation of phosphatases was noted, which would alter the balance of protein phosphorylation in several cellular signaling pathways; much more information is needed before any suggestions regarding functional effects can be made. As the oxidation of glutathione peroxidase might offer a new modulating mechanism of cellular signal transduction^[25], it was suggested that its increased expression could provide a sign in response to stress in the body.

Effect of baicalin on gene expression Programmed-cell-death-8 was upregulated after MCAO with baicalin treatment.

Furthermore, novel mediators of cerebral ischemia inducing neuronal death have also been found using cDNA microarrays. All these studies reinforce the idea that common cell death pathways are activated in response to neuron death inducers, which promise to serve as potential therapeutic targets for modulation to achieve neuroprotection^[26]. The expression of the protein kinase C-binding protein Zeta was downregulated in rats treated with baicalin, which resulted in a decrease in the activity of protein kinase C (PKC). Aronowski *et al*^[27] observed that the activity of PKC in the brain cortex and hippocampus decreased significantly after ischemia and this decrease could be alleviated by the pre-treatment of the NMDA receptor antagonist. Further studies need to examine whether baicalin acts as an antagonist to the NMDA receptor. After all, the relationship between the changing activity of PKC and neuron injury has not been elucidated completely in many key pathophysiological procedures of ischemia, such as increment of fermentation, acidosis, and deficiency of ATP production. The prominent differential genes involved in protein tyrosine phosphatase receptor type D and A are central to the course of signal transduction, which has been implicated both in the regulation of cell growth and the rearrangement of actin that is mediated by several receptor tyrosine kinases. Differential gene expression showed that baicalin played an important role in cell signal transduction and protein phosphorylation after MCAO, and might act as a neuroprotectant.

Effect of gardenin on gene expression The differential genes in the gardenin-treatment group showed extreme variation compared with the baicalin-treatment group. Several genes encoding anti-oxidation were upregulated. For example, an increase in Mu2 suggested that Mu2 played an activation role in anti-oxidation responses, and was regarded as one of the markers of internal anti-injury because of its anti-oxidative and antidotal effects^[25]. According to the actions of Mu2, the anti-oxidative and antidotal effects of gardenin might be one of the mechanisms of cerebral protection after MCAO.

The most interesting phenomenon observed was that there was no overlap in the genes showing differential expression in the three groups despite the similar trends in expression. The results suggest that there are considerable differences in the pharmacological effects of baicalin and gardenin at the molecular level after MCAO. Compared with the role of gardenin and bacalin, which are components of QKLI, it appears that QKLI plays an integral role and has important therapeutic effects requiring further investigation.

In conclusion, the cDNA microarray study not only confirmed that changes in many genes contributed to cerebral

ischemia, but also suggested several potential targets for further investigation. Of course, global expression of genes measured at the levels of mRNA transcription obtained using microarray analysis should be viewed from two different standpoints. Looking from mRNA towards protein, one would ultimately like to test the increased production of the proteins encoded by the upregulated mRNA or demonstrate a loss of protein encoded by downregulated mRNA. Thus, it may be rewarding to reconstruct the networks of regulation in response to ischemia and, by inference, to hypoxia, using bioinformatic strategies^[28]. Understanding such regulatory networks and the therapeutic mechanisms of QKLI for ischemia-hypoxia responsive gene expression in neurons may extend the relevance of these studies.

References

- Hou ST, MacManus JP. Molecular mechanisms of cerebral ischemia-induced neuronal death. *Int Rev Cytol* 2002; 221: 93–148.
- Chen X, Nishida H, Konishi T. Baicalin promoted the repair of DNA single strand breakage caused by H₂O₂ in cultured NIH3T3 fibroblasts. *Biol Pharm Bull* 2003; 26: 282–4.
- Huang Y, Wong CM, Lau CW, Yao X, Tsang SY, Su YL, *et al*. Inhibition of nitric oxide/cyclic GMP-mediated relaxation by purified flavonoids, baicalin and baicalein in rat aortic rings. *Biochem Pharmacol* 2004; 67: 787–94.
- Lee HH, Yang LL, Wang CC, Hu SY, Chang SF, Lee YH. Differential effects of natural polyphenols on neuronal survival in primary cultured central neurons against glutamate- and glucose deprivation-induced neuronal death. *Brain Res* 2003; 986: 103–15.
- Liu JJ, Huang TS, Cheng WF, Lu FJ. Baicalein and baicalin are potent inhibitors of angiogenesis: inhibition of endothelial cell proliferation, migration and differentiation. *Int J Cancer* 2003; 106: 559–65.
- Zhou JW. Gardenin and bacalin show the effect on MCP-1 of rat brain suffered focal ischemia. *Chin Arch Tradit Chin Med* 2004; 22:1016–7.
- Sharp FR, Lu A, Tang Y, Millhorn DE. Multiple molecular penumbras after focal cerebral ischemia. *J Cereb Blood Flow Metab* 2000; 20: 1011–32.
- Haruo N, Agasawa K, Yuya K. Correlation between cerebral blood flow and histologic changes in a new rat model of middle cerebral artery occlusion. *Stroke* 1989; 20: 1037–42.
- Bederson JB. Evaluation of 2,3,5-triphenyltetrazolium chloride as a stain for detection and quantification of experimental cerebral infarction in rats. *Stroke* 1986; 17: 1304–8.
- Jiang CH, Tsien JZ, Schultz PG, Hu Y. The effects of aging on gene expression in the hypothalamus and cortex of mice. *Proc Natl Acad Sci USA* 2001; 98: 1930–4.
- Jin K, Mao XO, Eshoo MW, Nagayama T, Minami M, Simon RP, *et al*. Microarray analysis of hippocampal gene expression in global cerebral ischemia. *Ann Neurol* 2001; 50: 93–103.
- Lee CK, Weindruch R, Prolla TA. Gene-expression profile of

- the ageing brain in mice. *Nat Genet* 2000; 25: 294–7.
- 13 Sandberg R, Yasuda R, Pankratz DP, Barlow C. Regional and strain-specific gene expression mapping in the adult mouse brain. *Proc Natl Acad Sci USA* 2000; 97: 11038–43.
 - 14 Hou ST, Callaghan D, Fournier MC, Hill I, Kang L, Massie B, *et al*. The transcription factor E2F1 modulates apoptosis of neurons. *J Neurochem* 2000; 75: 91–100.
 - 15 Jaken S. An overlay assay for detecting protein kinase C-binding proteins and substrates. *Methods Mol Biol* 2003; 233: 359–68.
 - 16 Belayev L, Liu Y, Zhao W, Busto R, Ginsberg MD. Human albumin therapy of acute ischemic stroke: marked neuroprotective efficacy at moderate doses and with a broad therapeutic window. *Stroke* 2001; 32: 553–60.
 - 17 Huh PW, Belayev L, Zhao W, Koch S, Busto R, Ginsberg MD. Comparative neuroprotective efficacy of prolonged moderate intras ischemic and postischemic hypothermia in focal cerebral ischemia. *J Neurosurg* 2000; 92: 91–9.
 - 18 Belayev L, Zhao W, Busto R, Ginsberg MD. Transient middle cerebral artery occlusion by intraluminal suture: I. Three-dimensional autoradiographic image-analysis of local cerebral glucose metabolism-blood flow interrelationships during ischemia and early recirculation. *J Cereb Blood Flow Metab* 1997; 17: 1266–80.
 - 19 Guhaniyogi J, Brewer G. Regulation of mRNA stability in mammalian cells. *Gene* 2001; 265: 11–23.
 - 20 Schmidt-Kastner R, Zhang B, Belayev L, Khoutorova L, Amin R, Busto R, *et al*. DNA microarray analysis of cortical gene expression during early recirculation after focal brain ischemia in rat. *Brain Res Mol Brain Res* 2002; 108: 81–93.
 - 21 Zhang ZG, Chopp M, Tang WX, Jiang N, Zhang RL. Postischemic treatment (2-4) with anti-CD11b and anti-CD18 monoclonal antibodies is neuroprotective after transient (2 h) focal cerebral ischemia in the rat. *Brain Res* 1995; 698: 79–85.
 - 22 Bogousslavsky J, Paciaroni M, Gallai V. Glycoprotein (GP) IIb/IIIa inhibitors for acute stroke treatment. *Clin Exp Hypertens* 2002; 24: 603–10.
 - 23 Wannier-Morino P, Rager G, Sonderegger P, Grabs D. Expression of neuroserpin in the visual cortex of the mouse during the developmental critical period. *Eur J Neurosci* 2003; 17: 1853–60.
 - 24 Hayashi T, Warita H, Abe K, Itoyama Y. Expression of cyclin dependent kinase 5 and its activator p35 in rat brain after middle cerebral artery occlusion. *Neurosci Lett* 1999; 265: 37–40.
 - 25 Paolicchi A, Dominici S, Pieri L, Maellaro E, Pompella A. Glutathione catabolism as a signaling mechanism. *Biochem Pharmacol* 2002; 64: 1027–35.
 - 26 Read SJ, Parsons AA, Harrison DC, Philpott K, Kabnick K, O'Brien S, *et al*. Stroke genomics: approaches to identify, validate, and understand ischemic stroke gene expression. *J Cereb Blood Flow Metab* 2001; 21: 755–78.
 - 27 Aronowski J, Waxham MN, Grotta JC. Neuronal protection and preservation of calcium/calmodulin-dependent PKII and PKC activity by dextrorphan treatment in lobal ischemia. *J Cereb Blood Flow Metab* 1993;13: 550–7.
 - 28 Thieffry D. From global expression data to gene networks. *Bioessays* 1999; 21: 895–9.
-

The 15th World Congress of Pharmacology (IUPHAR-2006)

2006, July 2–7 Beijing International Convention Center, China

Info: Ms Xiao-dan ZHAO
Chinese Pharmacological Society
1, Xian Nong Tan St
Beijing 100050, China
Phn/Fax 86-10-6316-5211
E-mail: zhxd@imm.ac.cn
<http://www.cnphars.org>