

Full-length article

Expression of PLK1 and survivin in non-Hodgkin's lymphoma treated with CHOPLin LIU, Min ZHANG, Ping ZOU¹*Department of Hematology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China***Key words**

Polo-like kinase 1; survivin; non-Hodgkin's lymphoma

¹ Correspondence to Prof Ping ZOU.

Phn/Fax 86-27-8572-6387.

E-mail xys387@126.com

Received 2007-07-19

Accepted 2007-10-15

doi: 10.1111/j.1745-7254.2008.00750.x

Abstract

Aim: The present study was designed to investigate the expression of Polo-like kinase 1 (PLK1) and survivin in non-Hodgkin's lymphoma (NHL). **Methods:** The expression of PLK1 and survivin were detected with immunohistochemical techniques. **Results:** The expression rate of PLK1 and survivin were 63.6% (56/88) and 79.5% (70/88) in NHL, respectively. PLK1 expression correlated with systemic symptoms, lactate dehydrogenase levels, and international prognostic index scores in B-NHL and T-NHL, while survivin did not. **Conclusion:** PLK1 and survivin are both overexpressed in NHL. There is a significant relationship between the overexpression of PLK1 and clinical features.

Introduction

Non-Hodgkin's lymphoma (NHL) forms a highly heterogeneous group of diseases. Despite steady improvement in their classification, patients with the same diagnosis can have markedly different clinical courses. Since prognosis remains uncertain, several biologically-based prognostic indicators have been proposed in addition to the international prognostic index (IPI) scores. However, none have proven to be completely effective^[1], so novel independent factors are required to better characterize the malignant potential of NHL.

Accumulating evidence suggests that abnormalities in cell cycle and apoptosis are hallmarks of clonal expansion leading to cancer emergence^[2,3]. Polo-like kinase 1 (PLK1) has been implicated in various aspects of the progression of the cell cycle^[4], including centrosome maturation^[5], assembly of the mitotic spindle^[6], regulation of the anaphase-promoting complex^[7], and cytokinesis^[8]. Survivin, a new inhibitor of the apoptosis protein (IAP) family, directly inhibits caspase 3 and caspase 7 and is one of the strongest anti-apoptosis factors by far^[9]. Although PLK1 and survivin play completely different roles, they have some characteristics in common^[10-13]: they are mostly expressed in the G₂/M phase of the cell cycle and they are not expressed in most normal differentiated tissues; however, they are overexpressed in a broad spectrum of cancer types and the

expression often correlates with poor outcomes. To assess the clinical significance of PLK1 and survivin in NHL, the expression of PLK1 and survivin were examined and correlated with clinical outcome.

Materials and methods

Patients A total of 88 patients (age: 5–86 years, median 47 years) were diagnosed with NHL at Union Hospital (Wuhan, China) between January 2001 and June 2006. None of the patients underwent radiotherapy or chemotherapy prior to diagnosis. Classification was established on standard hematoxylin-eosin-stained sections according to the guidelines of the World Health Organization 2001 Classification of Lymphoid Neoplasms^[14]. Eighty-eight cases were treated with CHOP or E-CHOP or CHOP plus Rituximab (600 mg/m² CTX, iv, d 1; 50 mg/m² ADR, iv, d 1; 2 mg/m² VCR, iv, d 1; 60 mg/m² Pred, po, d 1–7; 100 mg/m² etoposide, iv, d 1; 375 mg/m² Rituximab, iv, d 1). Clinical follow-up data were available for all patients. The median follow-up was 33 months (from 2 to 64 months). As a control for NHL, 5 cases of reactive lymph node proliferation were included in the study.

Immunohistochemistry on tissue slides Immunostaining was performed with primary antibodies against PLK1 (1:100; rabbit polyclonal antibody; Biologend, San Diego, CA, USA) and survivin (1:150; rabbit polyclonal

antibody; Neomarkers, Fremont, CA, USA). In brief, 4 μ m thick formalin-fixed paraffin sections were dewaxed and rehydrated through a graded series of ethanols. After antigen retrieval with microwave heating, the sections were treated with 0.3% hydrogen peroxide followed by incubation in bovine serum albumin. The sections were then incubated with primary antibodies. After washing with phosphate-buffered saline, the sections were incubated with biotinylated anti-immunoglobulin G of the appropriate species and streptavidin peroxidase using a labeled streptavidin biotin kit (Zhongshan Biotechnology, Beijing, China). The immunoperoxidase reaction product was visualized with diaminobenzidine/hydrogen peroxide. The sections were counterstained with hematoxylin.

Evaluation of tissue staining The staining intensity of the tissue slides was evaluated independently by 2 pathologists who were blinded to the patients' characteristics and survival. To assess the differences in staining intensity, an immunoreactivity scoring system (IRS) was applied. The intensity of staining was designated as negative (0), weak (1), moderate (2), or strong (3). Additionally, the percentage of positive cells was evaluated and scored as either no cells (0), less than 10% of cells (1), 10%–50% of cells (2), 51%–80% of cells (3), or over 80% of the cells stained (4). By multiplication of these 2 parameters, the IRS for each case was calculated. Finally, the cases were grouped as an-

tigen negative (IRS 0–6) or antigen positive (IRS 7–12) for statistical analysis^[15].

Statistical analysis Patient characteristics were compared by Fisher's exact test or χ^2 -test. Generally, $P < 0.05$ was considered statistically significant. For all statistical procedures, SPSS v11.5 software (SPSS, Chicago, IL, USA) was used.

Results

Expression of 2 proteins in NHL In all 88 NHL cases, 63.6% expressed PLK1 and 79.5% expressed survivin. The detailed distribution is listed in Table 1. The expression of PLK1 and survivin were then investigated in B-NHL and T-NHL, respectively. The PLK1 expression rate had no significance difference between B-NHL and T-NHL (57.8% vs 77.3%; $P = 0.103$), as did survivin (76.6% vs 86.4%; $P = 0.544$).

PLK1 was often strongly expressed in nuclei and seldom weakly in cytoplasm, while survivin was mainly strongly expressed in nuclei. None of the 5 reactive lymph node proliferation cases showed sufficient staining of PLK1 or survivin to be designated as positive (Figures 1, 2).

Relationship between the expression of 2 proteins and clinical characteristics in B-NHL and T-NHL In B-NHL, there was a highly significant positive correlation of PLK1 expression with age, Ann Arbor stage, lactate de-

Table 1. Expression of 2 proteins in NHL.

Type	Case (n)	PLK1 positive (n/%)	Survivin positive (n/%)
B-NHL			
Follicular lymphoma (FL)	7	4/57.1	2/28.6
MALT type	8	2/25	7/87.5
Splenic marginal zone lymphoma	2	0/0	1/50
SLL	5	2/40	3/60
Plasmacytoma	1	1/100	1/100
DLBCL	39	26/66.7	32/82.1
Primary mediastinal large B-cell lymphoma	1	0/0	1/100
Precursor B-lymphoblastic lymphoma	1	1/100	1/100
T-NHL			
Intestinal T-cell lymphoma	1	1/100	1/100
Anaplastic large cell lymphoma and null cell type	2	0/0	2/100
Extranodal natural killer/T-cell lymphoma, nasal type	5	5/100	5/100
Precursor T-lymphoblastic lymphoma	1	1/100	1/100
Angioimmunoblastic lymphoma	3	3/100	3/100
PTLU	10	8/80	8/80
Uncertain type	2	2/100	2/100




Figure 1. (A) expression of PLK1 in DLBCL ($\times 400$) and (B) PTLU ($\times 400$).




Figure 2. (A) expression of survivin in DLBCL ($\times 400$) and (B) PTLU ($\times 400$).

hydrogenase (LDH) levels, systemic symptoms, IPI scores, and therapeutic effect. However, in T-NHL, the PLK1 expression was only associated with LDH levels, systemic symptoms, and IPI scores. There were no significant relationships between the survivin expression and these outcomes in either B-NHL or T-NHL (Table 2).

Discussion

In recent years, a growing spectrum of biochemical investigations has contributed to better characterize biological tumor behavior, thereby improving treatment strategies^[16]. Among these studies, PLK1 and survivin were selected as the study objects, concerning the 2 important aspects of tumor biological behavior: proliferation and apoptosis.

PLK1 is a cell cycle-regulated, cyclin-independent serine/threonine protein kinase, and its protein level changes from very low level during the G₁ phase to at least a 10-fold increase by the G₂/M phase^[17]. PLK1 is involved in mitotic spindle organization and cytokinesis and seems to play an essential role in regulating chromosomal segregation^[18]. Furthermore, overexpressed PLK1 can cause the formation of oncogenic foci in NIH 3T3 cells, which are capable of forming tumors in nude mice^[19]. The el-

evated expression of PLK1 had been found in non-small cell lung cancer^[20], squamous cell carcinoma of the head and neck^[21], esophageal cancer^[22], gastric cancer^[23], ovarian cancer^[24], endometrial cancer^[25], colon cancer^[26], breast cancer^[27], and malignant melanoma^[28], prostate^[29], papillary carcinoma^[30], and oropharyngeal carcinomas^[31]. The expression of PLK1 was correlated with an adverse clinical outcome in some tumors.

Survivin belongs to the IAP family and it suppresses apoptosis induced by Fas, Bax, caspases, and anticancer drugs^[32]. Survivin is expressed in a strict cell cycle-dependent and tissue-restricted manner. It is expressed at high levels in the G₂/M phase and in proliferating tissues^[33]. Overexpressed survivin can be detected in almost all human tumors and is correlated with poor prognosis^[34].

In the present study, the expression rate of PLK1 was lower in some less aggressive or indolent types, such as small lymphocytic lymphoma (SLL) and marginal zone B-cell lymphoma (MALT) than that in some more aggressive types, such as diffuse large B-cell lymphoma (DLBCL) and peripheral T-cell lymphoma unspecified (PTLU), while the expression rate of survivin seemed to have no such trend. This suggested that PLK1 overexpression was correlated with cell-proliferating activity, and survivin

Table 2. Relationship between the expression of 2 proteins and the clinical characteristics of B-NHL and T-NHL.

Item	Case (n)	B-NHL		Survivin		Case (n)	T-NHL		Survivin	
		PLK1 positive (n)	P	positive (n)	P		PLK1 positive (n)	P	positive (n)	P
Age (years)										
>=60	19	7	0.027	14	0.753	2	1	0.411	2	0.74
<60	45	30		35		20	16		17	
Stage										
I+II	25	10	0.021	18	0.49	9	5	0.116	7	0.544
III+IV	39	27		31		13	12		12	
LDH										
Normal	37	13	<0.001	27	0.427	9	4	0.005	8	0.642
Elevated	27	24		22		13	13		11	
Systemic symptoms										
Yes	29	24	<0.001	25	0.097	14	14	0.002	11	0.273
No	35	13		24		8	3		8	
IPI score										
0~2	43	20	0.009	30	0.114	12	7	0.04	10	0.571
3~5	21	17		19		10	10		9	
Effect										
CR+PR	45	21	0.005	32	0.195	13	8	0.054	10	0.24
NCR	19	16		17		9	9		9	

overexpression implied the common characterization of NHL: the dysregulation of apoptosis.

PLK1 overexpression was positively linked to age, Ann Arbor stage, LDH levels, systemic symptoms, IPI scores, and therapeutic effect in B-NHL, and was only linked to LDH levels, systemic symptoms, and IPI scores in T-NHL. The reason may be the highly heterogeneity and limited number of cases of T-NHL. There was no significant difference between the survivin-negative and -positive groups for these characteristics, either in B-NHL or in T-NHL. It verified again that apoptosis resistance may be the essence of NHL.

In conclusion, PLK1 and survivin were expressed at high levels in NHL cases, and PLK1 overexpression was associated with elderly patient, advanced tumor stage, systemic symptoms, elevated LDH levels, and poor therapeutic effect. The overexpression of survivin was not correlated with bad outcome.

Acknowledgements

The authors thank Dr Xiu NIE and Hua-xiong PAN for supplying the lymphoma tissues and for their skillful technical assistance.

References

- 1 Fisher RI, Shah P. Current trends in large cell lymphoma. *Leukemia* 2003; 17: 1948–60.
- 2 Corn PG, El-Deiry WS. Derangement of growth and differentiation control in oncogenesis. *Bioessays* 2002; 24: 83–90.
- 3 Evan GI, Vousden KH. Proliferation, cell cycle and apoptosis in cancer. *Nature* 2001; 411: 342–8.
- 4 Lowery DM, Lim D, Yaffe MB. Structure and function of Polo-like kinases. *Oncogene* 2005; 24: 248–59.
- 5 Dai W, Wang Q, Traganos F. Polo-like kinases and centrosome regulation. *Oncogene* 2002; 21: 6195–200.
- 6 McInnes C, Mazumdar A, Mezna M, Meades C, Midgley C, Scaerou F, *et al.* Inhibitors of Polo-like kinase reveal roles in spindle-pole maintenance. *Nat Chem Biol* 2006; 2: 608–17.
- 7 Lindon C, Pines J. Ordered proteolysis in anaphase inactivates Plk1

- to contribute to proper mitotic exit in human cells. *J Cell Biol* 2004; 164: 233–41.
- 8 Zhou T, Aumais J P, Liu X, Yu-Lee LY, Erikson RL. A role for Plk1 phosphorylation of NudC in cytokinesis. *Dev Cell* 2003; 5: 127–38.
 - 9 Shin S, Sung BJ, Cho YS, Kim HJ, Ha NC, Hwang JI, *et al*. An anti-apoptotic protein human survivin is a direct inhibitor of caspase-3 and -7. *Biochemistry* 2001; 40: 1117–23.
 - 10 Duffy MJ, O'Donovan N, Brennan DJ, Gallagher W M, Ryan BM. Survivin: a promising tumor biomarker. *Cancer Lett* 2007; 249: 49–60.
 - 11 Sah NK, Khan Z, Khan GJ, Bisen PS. Structural, functional and therapeutic biology of survivin. *Cancer Lett* 2006; 244: 164–71.
 - 12 Takai N, Hamanaka R, Yoshimatsu J, Miyakawa I. Polo-like kinases (Plks) and cancer. *Oncogene* 2005; 24: 287–91.
 - 13 Eckerdt F, Yuan J, Strebhardt K. Polo-like kinases and oncogenesis. *Oncogene* 2005; 24: 267–76.
 - 14 Jaffe ES, Harris NL, Stein H, editors. Pathology and genetics of tumours of hematopoietic and lymphoid tissues. World Health Organization Classification of Tumours, Lyon, France: IARC Press; 2001.
 - 15 Weichert W, Denkert C, Schmidt M, Gekeler V, Wolf G, Kobel M, *et al*. Polo-like kinase isoform expression is a prognostic factor in ovarian carcinoma. *Br J Cancer* 2004; 90: 815–21.
 - 16 Cheson BD. What is new in lymphoma? *CA Cancer J Clin* 2004; 54: 260–72.
 - 17 Strebhardt K, Ullrich A. Targeting polo-like kinase 1 for cancer therapy. *Nat Rev Cancer* 2006; 6: 321–30.
 - 18 van Vugt MA, van de Weerd BC, Vader G, Janssen H, Calafat J, Klompmaker R. Polo-like kinase-1 is required for bipolar spindle formation but is dispensable for anaphase promoting complex/Cdc20 activation and initiation of cytokinesis. *J Biol Chem* 2004; 279: 36 841–54.
 - 19 Smith MR, Wilson ML, Hamanaka R, Chase D, Kung H, Longo DL, *et al*. Malignant transformation of mammalian cells initiated by constitutive expression of the polo-like kinase. *Biochem Biophys Res Commun* 1997; 234: 397–405.
 - 20 Wolf G, Elez R, Doermer A, Holtrich U, Ackermann H, Stutte HJ, *et al*. Prognostic significance of polo-like kinase (PLK) expression in non-small cell lung cancer. *Oncogene* 1997; 14: 543–9.
 - 21 Knecht R, Elez R, Oechler M, Solbach C, von Ilberg C, Strebhardt K. Prognostic significance of polo-like kinase (PLK) expression in squamous cell carcinomas of the head and neck. *Cancer Res* 1999; 59: 2794–7.
 - 22 Tokumitsu Y, Mori M, Tanaka S, Akazawa K, Nakano S, Niho Y. Prognostic significance of polo-like kinase expression in esophageal carcinoma. *Int J Oncol* 1999; 15: 687–92.
 - 23 Kanaji S, Saito H, Tsujitani S, Matsumoto S, Tatebe S, Kondo A, *et al*. Expression of polo-like kinase 1 (PLK1) protein predicts the survival of patients with gastric carcinoma. *Oncology* 2006; 70: 126–33.
 - 24 Takai N, Miyazaki T, Fujisawa K, Nasu K, Hamanaka R, Miyakawa I. Expression of polo-like kinase in ovarian cancer is associated with histological grade and clinical stage. *Cancer Lett* 2001; 164: 41–9.
 - 25 Takai N, Miyazaki T, Fujisawa K, Nasu K, Hamanaka R, Miyakawa I. Polo-like kinase (PLK) expression in endometrial carcinoma. *Cancer Lett* 2001; 169: 41–9.
 - 26 Takahashi T, Sano B, Nagata T, Kato H, Sugiyama Y, Kunieda K. Polo-like kinase 1 (PLK1) is overexpressed in primary colorectal cancers. *Cancer Sci* 2003; 94: 148–52.
 - 27 Wolf G, Hildenbrand R, Schwar C, Grobholz R, Kaufmann M, Stutte HJ, *et al*. Polo-like kinase: a novel marker of proliferation: correlation with estrogen-receptor expression in human breast cancer. *Pathol Res Pract* 2000; 196: 753–9.
 - 28 Kneisel L, Strebhardt K, Bernd A, Wolter M, Binder A, Kaufmann R. Expression of polo-like kinase (PLK1) in thin melanomas: a novel marker of metastatic disease. *J Cutan Pathol* 2002; 29: 354–8.
 - 29 Weichert W, Schmidt M, Gekeler V, Denkert C, Stephan C, Jung K, *et al*. Polo-like kinase 1 is overexpressed in prostate cancer and linked to higher tumor grades. *Prostate* 2004; 60: 240–5.
 - 30 Ito Y, Miyoshi E, Sasaki N, Kakudo K, Yoshida H, Tomoda C, *et al*. Polo-like kinase 1 overexpression is an early event in the progression of papillary carcinoma. *Br J Cancer* 2004; 90: 414–8.
 - 31 Knecht R, Oberhauser C, Strebhardt K. PLK (polo-like kinase), a new prognostic marker for oropharyngeal carcinomas. *Int J Cancer* 2000; 89: 535–6.
 - 32 Wheatley SP, McNeish IA. Survivin: a protein with dual roles in mitosis and apoptosis. *Int Rev Cytol* 2005; 247: 35–88.
 - 33 Li F. Role of survivin and its splice variants in tumorigenesis. *Br J Cancer* 2005; 92: 212–6.
 - 34 Altieri DC. Survivin, versatile modulation of cell division and apoptosis in cancer. *Oncogene* 2003; 22: 8581–9.