

Review

Pharmacological tools for lysophospholipid GPCRs: development of agonists and antagonists for LPA and S1P receptors

Dong-Soon IM*

College of Pharmacy (BK21 Project) and Longevity Institute of Life Science and Technology, Pusan National University, Busan 609–735, Republic of Korea

Previous studies on lysophosphatidic acid (LPA) and sphingosine 1-phosphate (S1P) using various approaches have shown that both the molecules can act as intercellular signaling molecules. The discovery of the Edg subfamily of G-protein-coupled receptors (GPCRs) (later renamed LPA₁₋₃ and S1P₁₋₅) for these molecules has opened up a new avenue for pathophysiological research on lysophospholipids. Genetic and molecular studies on lysophospholipid GPCRs have elucidated pathophysiological impacts and roles in cellular signaling pathways. Recently, lysophospholipid GPCR genes have been used to develop receptor subtype-selective agonists and antagonists. The discovery of FTY720, a novel immune modulator, along with other chemical tools, has provided a means of elucidating the functions of each lysophospholipid GPCR on an organ and the whole body level. This communication attempts to retrospectively review the development of agonists and antagonists for lysophospholipid GPCRs, provide integrated information on pharmacological tools for lysophospholipid GPCR signaling, and speculate on future drug development.

Keywords: lysophosphatidic acid; sphingosine 1-phosphate; agonist; antagonist; G-protein-coupled receptor; lysolipid

Acta Pharmacologica Sinica (2010) 31: 1213–1222; doi: 10.1038/aps.2010.135; published online 23 Aug 2010

Discovery of GPCRs for LPA and S1P

Lysophosphatidic acid (LPA) and sphingosine 1-phosphate (S1P) are two representative lysophospholipid mediators acting on G-protein-coupled receptors (GPCRs). LPA was first identified three decades ago as a factor affecting blood pressure, platelet aggregation, and smooth muscle contraction^[1-5] and rediscovered as a mitogenic serum lipid inducer of neurite retraction^[6,7]. The name 'lyso' originates from lysis of blood cells; thus, the non-specific detergent action of LPA has been doubted. In 1996, the high potency (nmol/L range) and GPCR implications of LPA were finally connected to the discovery of the first LPA receptor (LPA₁, formerly known as Edg-2)^[8]. Subsequently, LPA₂ and LPA₃ were identified as members of the endothelial differentiation gene (Edg) subfamily of GPCRs^[9-11]. These three LPA receptors (Edg family) share a high homology with each other^[12].

Recently, the non-Edg family of LPA receptors, LPA₄ (GPR23, p2y9), LPA₅ (GPR92), and LPA₆ (p2y5), were reported as members of the purinergic GPCR cluster^[13-19]. Using the

GPCR-Gα₁₆ fusion expression system, Fujita's group reported GPR87 as another LPA receptor and P2Y₁₀ as a dual receptor for both LPA and S1P^[20,21]. These results have not yet been confirmed by other research groups^[22]. Instead, lysophosphatidylserine has been suggested to act as a ligand for P2Y₁₀ without confirmation of LPA and S1P as ligands in NIH3T3 cells^[23].

S1P was initially reported 15 years ago to be a second messenger, mediating an increase in calcium levels due to PDGF and IgE signaling^[24,25]. The molecular target of S1P in the cytosol has not yet been identified. The initial finding of an S1P-induced calcium increase stimulated research in S1P biology and was linked to its recognition as an intercellular first messenger. The involvement of trimeric G proteins in S1P-induced actions as well as pertussis toxin sensitivity strongly suggested the presence of S1P GPCRs in the plasma membrane^[26-30]. The discovery of S1P₁ (formerly known as Edg-1) in 1998, along with four other receptors (S1P₂₋₅) of the Edg subfamily GPCRs, became a milestone in S1P biology^[12,31-36].

Development of agonists and antagonists for LPA receptors

Prior to receptor cloning, various aspects of LPA receptor

* To whom correspondence should be addressed.

E-mail imds@pusan.ac.kr

Received 2010-05-25 Accepted 2010-07-16

structure-activity relationships had been studied: 1) fatty acid chain length and the presence of double bonds, 2) acyl and alkyl linkage, 3) stereo-selectivity on the *sn*-2 position, and 4) modification of phosphate groups^[37]. Oleoyl LPA (18:1) was considered the optimum ratio to increase Ca²⁺ levels in A431 cells^[38]. Alkyl LPA showed a better effect than acyl LPA in platelet aggregation assays and an equal effect in other responses^[39, 40]. Several groups reported a lack of stereo-selectivity on the *sn*-2 position of LPA. Early studies showed that modification of the phosphate group resulted in an absence of LPA responses^[38, 40].

The initial synthetic approach to LPA response was characterized using platelet aggregation assays. Replacement of the glycerol backbone with amino acids resulted in the production of *N*-palmitoyl serine phosphatidic acid (NPSPA) and *N*-palmitoyl tyrosine phosphatidic acid (NPTyrPA)^[40]. These molecules were not active on platelet aggregation in humans, but did demonstrate activity as LPA receptor antagonists in *Xenopus*^[41]. Later, NPSPA and NPTyrPA were reported as partial agonists of mammalian LPA receptors^[42, 43]. Sugiura *et al* introduced the ethanolamine-based LPA mimetic, *N*-acyl aminoethanol phosphoric acid (NAEPA), as an equipotent agonist of platelet aggregation^[40]. Later, research groups from the University of Virginia used NASPA and NAEPA as platforms to synthesize a series of VPC compounds^[37, 44]. The phosphonate analogue of NAEPA lost its platelet aggregation properties^[40], but methylene phosphonate LPA was equipotent to LPA as an inhibitor of forskolin-driven cAMP accumulation in rat C6 glioma cells^[45]. Jalink *et al* synthesized various phosphonate analogues along with fatty alcohol phosphates and the methyl ester of LPA (lysophosphatidylmethanol, LPM), but could not show a significant impact of these compounds on Ca²⁺ increase in A431 cells^[38]. Ironically, these chemicals turned out to be selective or non-selective agonists of cloned LPA receptors (see details below). In the early era of LPA biology, suramin and lysophosphatidylglycerol were used to demonstrate GPCR involvement in LPA responses^[46] and as an antagonist of LPA-induced Ca²⁺ responses in Jurkat T cells^[47], respectively.

LPA GPCR agonists

Since the discovery of the three-Edg family of LPA receptors, the development of selective receptor-subtype agonists and antagonists has accelerated. The optimal chain length and the presence of double bonds have been found to vary depending on receptor subtype. For example, LPA₃ showed a preference for unsaturated LPA similar to oleoyl LPA^[48], whereas LPA₆ showed a preference for 2-acyl LPA^[19]. Synthesis of LPA derivatives with phosphonate or thiophosphate groups instead of the phosphate group showed receptor-subtype selective activity similar to 1-oleoyl-2-*O*-methyl-*rac*-glycerophosphothionate (OMPT), 1-*O*-acyl- α -fluoromethylenephosphonate, and α -hydroxymethylenephosphonate LPA analogues as LPA₃ receptor selective agonists^[49-52]. The phosphonate derivatives also provided a path toward the development of phosphatase-resistant long-lasting LPA derivatives^[53]. A dialkyl phospho-

tidic acid (PA) 8:0 analog with a thiophosphate was reported as a potent and selective LPA₃ agonist^[54], but later, agonistic activity on the LPA₅ receptor and antagonistic activity on the LPA_{1,3} receptors was reported^[55]. Based on computational modeling, dodecyl fatty alcohol phosphate was shown to be a specific LPA₂ receptor agonist^[56], and later, oleoyl-thiophosphate was identified as a pan-agonist (LPA₁₋₃)^[43]. A methylene phosphonate LPA analogue was reported as a selective LPA₂ agonist, but it recently was shown to exert agonistic and antagonistic activities on LPA₅ and LPA₃, respectively^[52, 57]. T-15 (LPA₁ agonist) and T-13 (LPA₃ agonist) were both synthesized using a carbohydrate scaffold^[58]. Darmstoff analogues were introduced as novel scaffolds for subtype-selective LPA receptor ligands^[59]. Finally, octadecenyl phosphate was shown to be a selective agonist for LPA₄ and LPA₅ receptors^[55].

Although alkyl LPA has been shown to exert equipotent activity on each Edg-family LPA receptor compared to acyl LPA^[60], alkyl LPA was found to be more potent than acyl LPA in platelet aggregation responses^[3]. Therefore, the Edg-family LPA receptors were not able to account for the LPA response in platelets^[61]. Recently, non-Edg (purinergic) LPA receptors have been identified. The LPA₅/GPR92 receptor has demonstrated alkyl preference and a presence in platelets^[55]. Farnesyl diphosphate, *N*-arachidonyl glycine, and carba-cyclic phosphatidic acid were shown to be selective agonists for LPA₅^[55, 62, 63]. LPA₆/p2y5 showed a preference for 2-acyl LPA over 1-acyl LPA^[19]. Alkyl-OMPT, an LPA₃ agonist, and 2-linoleoyl LPA showed better agonistic effects on the LPA₆ receptor than 2-oleoyl LPA showed^[19, 64]. In contrast, the methyl ester of LPA (LPM) was shown to be a pan-agonist for LPA₁₋₅ although it was less potent than LPA^[65].

LPA GPCR antagonists

The development of LPA derivatives with a bulky group on the *sn*-2 position has demonstrated the stereo-selectivity of LPA receptors and has led to the further development of the LPA₁/LPA₃ selective antagonists VPC12249 and VPC32183^[66, 67]. 2-Pyridyl-containing phosphonate was developed as a non-hydrolyzable LPA₃ receptor antagonist^[68]. Diocetyl glyceropyrophosphate (DGPP) and Ki16425 were determined to be selective LPA₁/LPA₃ antagonists following screening of available lipid and non-lipid molecules^[69, 70]. The thiophosphate version of PA 8:0 was shown to be the most potent LPA₃ antagonist, whereas the dialkyl PA 8:0 analog with a thiophosphate was shown to be a potent and selective LPA_{3,5} agonist, as mentioned above^[54, 55]. Tetradecyl-phosphonate was identified as a pan-antagonist (LPA₁₋₃)^[43]. Furthermore, farnesyl phosphate and farnesyl diphosphate potentially and selectively blocked the LPA₃-mediated Ca²⁺ increase^[71], although farnesyl diphosphate is a LPA₅ selective agonist, as mentioned above^[55]. T-14, which used a carbohydrate scaffold, was shown to be an LPA₃ antagonist^[58].

Recently, virtual screening identified non-lipid LPA₃ antagonist NSC161613, LPA₂ antagonist H2L5186303, LPA_{4,5} antagonist 5987411, and LPA_{1,6} antagonist 5765834^[55, 72]. An LPA analogue of α -bromomethylene phosphonate was reported to act

as a pan-antagonist to LPA₁₋₄ and an autotoxin inhibitor^[52, 73].

Pharmacological tools for LPA GPCR signaling

Commercially available chemicals for studying LPA receptor subtypes are currently in development, although the effects of previously developed chemicals on recently identified non-Edg LPA receptors have not been completely verified (Figure 1). For LPA₁ or LPA₃ receptor signaling, a combined application of LPA_{1,3} antagonists such as VPC12249, VPC32193, DGPP, and Ki16425 and LPA₃ agonists such as OMPT and α -fluoromethylene phosphonate would be more favorable. For LPA₂ receptor signaling, dodecyl phosphate is an adequate LPA₂ selective agonist. For LPA₅ receptor, farnesyl diphosphate could be used as a selective agonist. For LPA₆ receptor, alkyl OMPT, an LPA_{3,6} agonist, could be used in combination with LPA_{1,3} antagonists (Table 1).

Development of agonist and antagonist for S1P receptors

In contrast to LPA, the structure-activity relationship of S1P has a very short history. Using cloned S1P receptors, sphinganine-1-phosphate (dihydro-S1P) and sphingosylphosphorylcholine (SPC) were shown to be equipotent and far less potent on each S1P receptor^[36, 74-76].

S1P GPCR agonists

Following the cloning of the S1P receptors, the development of S1P agonists and antagonists began. The importance of the *D-erythro* configuration of S1P was demonstrated using the cloned receptors^[77]. The linkage of the immune modulator FTY720 to S1P receptors, however, boosted this area of research and opened a new direction for S1P biology^[78-80]. Lymphopenia induction by inhibiting lymphocyte egress from

lymphoid organs was shown to be mediated through the S1P₁ receptor^[81]. High-throughput screening (HTS) of an available chemical library showed that SEW2871 acted as an *in vivo* active heterocyclic S1P₁ selective agonist^[81, 82] and compound 26 was synthesized as a potent 3,5-diphenyl-12,4-oxadiazole S1P₁ agonist^[83]. Later, using ultra-HTS, 3,5-diaryloxadiazole (CYM5181) and dicyclohexylamide were found to be selective agonists for S1P₁ and S1P₃, respectively^[84]. Using computational modeling, CYM-5442 was developed as an S1P₁ selective agonist that was more potent than CYM5181^[85]. AUY954, an aminocarboxylate analogue of FTY720, was also introduced as an S1P₁ selective agonist^[86]. VPC01091, a cyclized analogue of FTY720, was shown to act as an orally active S1P₁ agonist and an S1P₃ antagonist^[87]. KRP-203 is a pro-drug immune modulator similar to FTY720; the phosphorylated form of KRP-203 was shown to be a selective S1P₁ agonist^[88, 89]. Constrained azacyclic analogues of FTY720 showed selective agonist activities on S1P₄ and S1P₅ receptors^[90]. Finally, phytosphingosine-1-phosphate was shown to act as a potent and selective agonist on the S1P₄ receptor^[76].

S1P GPCR antagonists

Suramin was temporarily used as an S1P₃ antagonist^[75, 91]. Human S1P₅ was also reported to be sensitive to suramin and its analogue NF023^[92].

Following screening of an available chemical library, JTE-013, a pyrazopyridine derivative, was identified as an S1P₂ antagonist^[93, 94]. Modification of the FTY720-phosphate structure led to the development of VPC23019 and VPC25239 as selective S1P₁/S1P₃ antagonists^[95]. As mentioned above, VPC01091 is an orally active S1P₁ agonist and S1P₃ antagonist^[87]. W146, hexyl phenyl amide phosphonate, was found to




Figure 1. Structures of commercially available agonists and antagonists for LPA GPCR signaling. Sources are Avanti polar lipid, Biomol international, Echelon bioscience, Enzo Life sciences, and Sigma-Aldrich.

Table 1. Agonistic and antagonistic characters of each compound on each LPA receptor. Numbers (nmol/L) mean EC₅₀ or K_D values for agonists and IC₅₀ or K_I values for antagonists.

Name	LPA ₁	LPA ₂	LPA ₃	LPA ₄	LPA ₅	LPA ₆
Ki16425 ^[70]	Antagonist (250 nmol/L)	Antagonist (5600 nmol/L)	Antagonist (360 nmol/L)			
DGPP ^[69]	Antagonist (6600 nmol/L)		Antagonist (106 nmol/L)			
VPC32183 ^[67]	Antagonist (109 nmol/L)		Antagonist (175 nmol/L)			
VPC12249 ^[66]	Antagonist (137 nmol/L)		Antagonist (428 nmol/L)			
2-Pyridyl phosphonate ^[68]			Antagonist			
Thiophosphatidic acid 8:0 ^[54]	Antagonist (360 nmol/L)		Antagonist (5 nmol/L)			
T14 ^[58]			Antagonist			
NSC161613 ^[72]			Antagonist (24 nmol/L)			
Compound 12 ^[120]	Antagonist (48 nmol/L)		Antagonist (230 nmol/L)			
H2L5186303 ^[72]		Antagonist (7.2 nmol/L)	Antagonist (310 nmol/L)			
5987411 ^[55]				Antagonist (741 nmol/L)	Antagonist (1300 nmol/L)	
5765834 ^[55]	Antagonist (48 nmol/L)				Antagonist (292 nmol/L)	
α-bromomethylene phosphonate ^[52,55,73,117]	Antagonist (751 nmol/L)	Antagonist (304 nmol/L)	Antagonist (380 nmol/L)	Antagonist (167 nmol/L)	Weak agonist	
Tetradecyl-phosphonate ^[43]	Antagonist (10000 nmol/L)	Antagonist (5500 nmol/L)	Antagonist (3100 nmol/L)			
Farnesyl diphosphate ^[55,71]		Antagonist (2100 nmol/L)	Antagonist (155 nmol/L, 4600 nmol/L)	Antagonist (1980 nmol/L)	Agonist (40 nmol/L)	
Carba cyclic PA ^[55]					Agonist	
OMPT ^[49]			Agonist (68 nmol/L)			
Alkyl OMPT ^[19,64]	Agonist (790 nmol/L)		Agonist (62 nmol/L)			Agonist
α-fluoromethylene phosphonate ^[51]	Weak agonist	Weak agonist	Agonist (0.5 nmol/L)			
α-hydroxymethylene-phosphonate ^[52]			Agonist (393 nmol/L)			
Compound 8bo ^[121]	Agonist (9.1 nmol/L)		Agonist (123 nmol/L)			
Dialkyl thiophosphatidic acid 8:0 ^[54, 55]	Agonist (695 nmol/L)	Agonist (5720 nmol/L)	Agonist (3 nmol/L)		Agonist	
	Antagonist (382 nmol/L)		Antagonist (184 nmol/L)			
Dodecylphosphate ^[56]		Agonist (700 nmol/L)	Antagonist (90 nmol/L)			
α-methylene phosphonate ^[52,55]		Agonist (>281 nmol/L)		Weak agonist (3900 nmol/L)	Agonist	
		Antagonist (1420 nmol/L)				
T15 ^[58]	Agonist (5 nmol/L)		Agonist (50 nmol/L)			

(Continued)

Name	LPA ₁	LPA ₂	LPA ₃	LPA ₄	LPA ₅	LPA ₆
T13 ^[58]	Week agonist (500 nmol/L)		Agonist (0.5 nmol/L)			
Octadecenyl phosphate ^[55]				Agonist (608 nmol/L)	Agonist	
NPSA ^[11,43,122]	Weak agonist (1850 nmol/L)	Weak agonist	Weak agonist (1600 nmol/L)			
NPTyrPA ^[43,55,122]	Antagonist (3450 nmol/L)	Weak agonist (11000 nmol/L)	Antagonist (5570 nmol/L)			Agonist
NAEPA ^[41]	Agonist	Agonist	Weak agonist			
Oleoyl-thiophosphate ^[43]	Agonist (193 nmol/L)	Agonist (244 nmol/L)	Agonist (546 nmol/L)			
LPM ^[65]	Agonist	Agonist	Agonist	Agonist	Agonist	

be a selective S1P₁ antagonist^[96]. VPC44116, an octyl analogue of W146 and γ -aminophosphonate analogue of VPC23019, antagonized lymphopenia and lung permeability via the S1P₁ receptor^[97]. SB64146 was reported to act as an inverse agonist on the S1P₁ receptor^[98]. Ascotricins A and B were isolated from a cultured broth of a fungus identified as *Ascotricha chartarum* and shown to inhibit the S1P₁ receptor and S1P-mediated HUVEC migration^[99]. Sankyo Co synthesized compound lead 2 (CL2), 2-(4-ethoxyphenoxy)-5-(3-octadecyl-5-oxo-4,5-dihydro-1H-pyrazol-1-yl) benzenesulfonate, which antagonized the S1P₁>S1P₃>S1P₂ receptors^[100]. Human S1P₁ receptor-selective antagonist and agonist effects of a rat monoclonal antibody (4B5.2) *in vivo* were also reported^[101]. Using a 3D database search, BML-241, 2-alkylthiazolidine-4-carboxylic acid, was found to act as an S1P₃ antagonist, but its selectivity and potency were not recapitulated in CHO-K1 cells expressing the S1P₃ receptor^[102,103]. A pharmacophore-based design of an S1P₃ antagonist with a 3,4-dialkoxybenzophenone scaffold was suggested^[104].

Pharmacological tools for S1P GPCR signaling

Commercially available tools for studying S1P receptor sub-

types are highlighted in Figure 2. For S1P₁ receptor signaling, CYM-5442 or SEW2871, both potent selective S1P₁ agonists, and W146, a selective S1P₁ antagonist, should be sufficient to elucidate S1P₁ receptor involvement. S1P₂ receptor signaling could be dissected using JTE-013, an S1P₂ selective antagonist. For S1P₃ GPCR signaling, a combined application of an S1P_{1,3} antagonist (VPC23019) and S1P₁ antagonist (W146) or S1P₁ agonist (CYM-5442) could be useful. Phytosphingosine 1-phosphate, an S1P₄ selective agonist, could be used to study S1P₄-mediated signaling. S1P_{1,3} antagonist (VPC23019)-insensitive, S1P₂ antagonist (JTE-013)-insensitive, S1P₄ agonist-non-responsive, and S1P- or FTY720-phosphate-sensitive signaling might be interpreted as S1P₅ receptor or unidentified S1P receptor signaling (Table 2).

Development of drugs acting on lysophospholipid GPCRs

Prior to the molecular cloning of GPCRs for LPA and S1P, a number of studies were conducted with the primary goal of determining the functions of lipid mediators at both the cellular and the organ level. These functions included platelet aggregation, smooth muscle contraction, and cell proliferation, among others^[1-6,105]. The discovery of GPCRs allowed signal




Figure 2. Structures of commercially available agonists and antagonists for S1P GPCR signaling. Sources are Avanti polar lipid, Biomol international, Cayman chemical, Echelon bioscience, Enzo Life sciences, Sigma-Aldrich, and Tocris.

Table 2. Agonistic and antagonistic characters of each compound on each S1P receptor. Numbers (nmol/L) mean EC₅₀ or K_D values for agonists and IC₅₀ or K_I values for antagonists.

Name	SIP ₁	SIP ₂	SIP ₃	SIP ₄	SIP ₅
AUY954 ^[86]	Agonist (1.2 nmol/L)		Agonist (1210 nmol/L)		Agonist (340 nmol/L)
CYM-5442 ^[85]	Agonist (1.2 nmol/L)				
CYM-5181 ^[84]	Agonist (3.4 nmol/L)				
SEW2871 ^[81]	Agonist (13 nmol/L)				
Compound 26 ^[83]	Agonist (0.6 nmol/L)		Agonist (12000 nmol/L)	Agonist (70 nmol/L)	Agonist (1.0 nmol/L)
Compound 12 ^[90]				Agonist (7.4 nmol/L)	Agonist (10.2 nmol/L)
Compound 18 ^[90]				Agonist (16.8 nmol/L)	Agonist (5.8 nmol/L)
VPC44116 ^[97]	Antagonist (30 nmol/L)		Antagonist (300 nmol/L)		Partial agonist (33 nmol/L)
W146 ^[96,106]	Antagonist (36 nmol/L)				
SB649146 ^[98]	Antagonist (300 nmol/L)				
VPC23019 ^[95]	Antagonist (13.8 nmol/L)		Antagonist (1175 nmol/L)	Agonist (263 nmol/L)	Partial agonist (85.1 nmol/L)
VPC25239 ^[95]	Antagonist (13.4 nmol/L)		Antagonist (97.7 nmol/L)	Agonist (166 nmol/L)	Partial agonist (11.5 nmol/L)
CL2 ^[100]	Antagonist (4400 nmol/L)	Antagonist (37000 nmol/L)	Antagonist (6700 nmol/L)		
VPC01091-P ^[87]	Agonist (6.6 nmol/L)		Antagonist	Agonist	Partial agonist
Ascotricins A and B ^[99]	Antagonist (8200 nmol/L, 1800 nmol/L)				
JTE-013 ^[93]		Antagonist (17 nmol/L)			
BML-241 ^[102, 103]			Antagonist (?)		
TY-52156 ^[123]			Antagonist (110 nmol/L)		
DS-SG-44 ^[124]	Agonist	Agonist	Agonist		
KRP-203-P ^[88,89]	Agonist (0.8 nmol/L)			Agonist (9.6 nmol/L)	
PhytoS1P ^[76]				Agonist (1.6 nmol/L)	
DihydroS1P ^[36,75,76]	Agonist	Agonist	Agonist	Agonist (8.6 nmol/L)	Agonist
SPC ^[34,36,75]	Partial agonist	Partial agonist	Partial agonist	Partial agonist	Partial agonist
FTY720-P ^[78,79]	Agonist (6.3 nmol/L), functional antagonist		Agonist (4.0 nmol/L)	Agonist (6.3 nmol/L)	Agonist (6.3 nmol/L)
AFD-R ^[79]	Agonist (2.5 nmol/L)		Agonist (4.0 nmol/L)	Agonist (4.0 nmol/L)	Agonist (1.3 nmol/L)

transduction studies to proceed in cells over-expressing the receptors and in receptor knock-down transgenic mouse models^[106–109]. The discovery of the pathophysiological significance of LPA and S1P, particularly on each receptor subtype, would contribute favorably to new drug development. In developing new medications acting on LPA or S1P receptors, subtype selectivity would be a major issue along with potency and efficacy to avoid side effects and ensure drug safety.

FTY720 was initially developed as an immune modulator for organ transplant patients^[110]. At present, this compound is under clinical study for the treatment of multiple sclerosis^[111]. Amira Pharmaceuticals reported LPA₁ selective antagonist (4'-{4-[(R)-1-(2-chloro-phenyl)-ethoxycarbonylamino]-3-methyl-isoxazol-5-yl]-biphenyl-4-yl)-acetic acid, AP2966, which showed good therapeutic potential in idiopathic pulmonary fibrosis and good pharmacokinetic profiles, including oral bioavailability^[112]. Pfizer global research and development introduced the S1P₁ selective agonists PF-A and PF-B. These agonists resulted in lymphopenia in rats and monkeys similar to FTY720 and reduced collagen-induced arthritis^[113]. Using a different approach, Lpath Inc. recently introduced

monoclonal antibodies against S1P and LPA. Humanized anti-S1P monoclonal antibody (mAb) sonopizumab blocked the tumorigenic effect of S1P produced by cancer cells as well as the angiogenic effect induced during pathological angiogenesis^[114, 115]. The LPA receptor pan-antagonist (LPA_{1–4}) and autotoxin inhibitor, α -bromomethylene phosphonate LPA analogue, was shown to be an excellent anti-cancer agent^[116, 117].

Closing remarks

Following the initial discoveries of LPA activity 30 years ago and S1P activity 15 years ago, a dark dawn era in lysophospholipid biology occurred due to the lack of identifiable targets. The discovery of the Edg-family GPCRs for LPA and S1P shined light on this field. The first meeting that focused on lysophospholipid biology was a New York Academy of Science meeting conducted in 1999 at Rockefeller University^[118]. Since 2001, FASEB summer research conferences on lysophospholipids have been held biannually. Lysophospholipid receptor nomenclature has been systematically assigned^[119], and more LPA receptors (purinergic or non-Edg LPA recep-

tors) are being discovered each year. Every two years, we have had exciting findings including the linkage of FTY720 to the S1P receptor, discovery of an autotoxin as a LPA-producing lysoPLD, lysolipid-sensitive proton-sensing GPCRs (OGR1 subfamily), and finally, the development of new chemicals. This review integrates accumulated information regarding pharmacological tools for lysophospholipid GPCR signaling to compare their characteristics and provides valuable information such as available chemical sources. Using a combination of receptor expression patterns in each organ or cell, these pharmacological tools might prove useful in defining the pathophysiological impact and significance of lysophospholipids. As mentioned above, the control of lysophospholipid functions using specific agonists or antagonists will contribute toward novel drug development. At a 2009 FASEB meeting in Carefree (Arizona, USA), we began to see several of the chemicals described here, in addition to FTY720, being used in clinical applications^[112-114, 116]. We now know more about many things than ever before. It is very likely that in the near future, the agonists/antagonists for LPA or S1P receptors will be on the market commercially and that there will be a section on lysophospholipid GPCRs in every basic pharmacology textbook.

Acknowledgements

This work was supported by the National Research Foundation, 2010 Korea-Japan Joint Research Grant (2010-616-E00015).

References

- Gerrard JM, Kindom SE, Peterson DA, Peller J, Krantz KE, White JG. Lysophosphatidic acids. Influence on platelet aggregation and intracellular calcium flux. *Am J Pathol* 1979; 96: 423-38.
- Tokumura A, Fukuzawa K, Yamada S, Tsukatani H. Stimulatory effect of lysophosphatidic acids on uterine smooth muscles of non-pregnant rats. *Arch Int Pharmacodyn Ther* 1980; 245:74-83.
- Simon MF, Chap H, Douste-Blazy L. Human platelet aggregation induced by 1-alkyl-lysophosphatidic acid and its analogs: a new group of phospholipid mediators? *Biochem Biophys Res Commun* 1982; 108: 1743-50.
- Tsukatani H, Yamada S, Tokumura A, Miyamoto T, Takauchi K. Isolation of an acute hypotensive substance from bovine brain lipid fraction. *Chem Pharm Bull (Tokyo)* 1976; 24: 2294-300.
- Moolenaar WH. Lysophosphatidic acid, a multifunctional phospholipid messenger. *J Biol Chem* 1995; 270: 12949-52.
- van Corven EJ, Groenink A, Jalink K, Eichholtz T, Moolenaar WH. Lysophosphatidate-induced cell proliferation: identification and dissection of signaling pathways mediated by G proteins. *Cell* 1989; 59: 45-54.
- Tigy G, Mileddi R. Lysophosphatidates bound to serum albumin activate membrane currents in *Xenopus* oocytes and neurite retraction in PC12 pheochromocytoma cells. *J Biol Chem* 1992; 267: 21360-7.
- Hecht JH, Weiner JA, Post SR, Chun J. Ventricular zone gene-1 (vzg-1) encodes a lysophosphatidic acid receptor expressed in neurogenic regions of the developing cerebral cortex. *J Cell Biol* 1996; 135: 1071-83.
- An S, Bleu T, Hallmark OG, Goetzl EJ. Characterization of a novel subtype of human G protein-coupled receptor for lysophosphatidic acid. *J Biol Chem* 1998; 273: 7906-10.
- Bandoh K, Aoki J, Hosono H, Kobayashi S, Kobayashi T, Murakami-Murofushi K, et al. Molecular cloning and characterization of a novel human G-protein-coupled receptor, EDG7, for lysophosphatidic acid. *J Biol Chem* 1999; 274: 27776-85.
- Im DS, Heise CE, Harding MA, George SR, O'Dowd BF, Theodorescu D, et al. Molecular cloning and characterization of a lysophosphatidic acid receptor, Edg-7, expressed in prostate. *Mol Pharmacol* 2000; 57: 753-9.
- Lynch KR, Im DS. Life on the edge. *Trends Pharmacol Sci* 1999; 20: 473-5.
- Noguchi K, Ishii S, Shimizu T. Identification of p2y9/GPR23 as a novel G protein-coupled receptor for Lysophosphatidic acid, structurally distant from the Edg family. *J Biol Chem* 2003; 278: 25600-6.
- Lee CW, Rivera R, Dubin AE, Chun J. LPA₄/GPR23 is an LPA receptor utilizing G_s, G_q/G_i-mediated calcium signaling and G_{12/13}-mediated Rho activation. *J Biol Chem* 2007; 282: 4310-7.
- Lee CW, Rivera R, Gardell S, Dubin AE, Chun J. GPR92 as a New G_{12/13}- and G_q-coupled Lysophosphatidic Acid Receptor That Increases cAMP, LPA₅. *J Biol Chem* 2006; 281: 23589-97.
- Kotarsky K, Boketoft A, Bristulf J, Nilsson NE, Norberg A, Hansson S, et al. Lysophosphatidic Acid Binds to and Activates Gpr92, a G Protein-Coupled Receptor Highly Expressed in Gastro-Intestinal Lymphocytes. *J Pharmacol Exp Ther* 2006; 318: 619-628.
- Yanagida K, Ishii S, Hamano F, Noguchi K, Shimizu T. LPA₄/p2y9/GPR23 mediates Rho-dependent morphological changes in a rat neuronal cell line. *J Biol Chem* 2007; 282: 5814-24.
- Pasternack SM, von Kugelgen I, Aboud KA, Lee YA, Ruschendorf F, Voss K, et al. G protein-coupled receptor P2Y5 and its ligand LPA are involved in maintenance of human hair growth. *Nat Genet* 2008; 40: 329-34.
- Yanagida K, Masago K, Nakanishi H, Kihara Y, Hamano F, Tajima Y, et al. Identification and characterization of a novel lysophosphatidic acid receptor, p2y5/LPA₆. *J Biol Chem* 2009; 284: 17731-41.
- Tabata K, Baba K, Shiraiishi A, Ito M, Fujita N. The orphan GPCR GPR87 was orphanized and shown to be a lysophosphatidic acid receptor. *Biochem Biophys Res Commun* 2007; 363: 861-6.
- Murakami M, Shiraiishi A, Tabata K, Fujita N. Identification of the orphan GPCR, P2Y(10) receptor as the sphingosine-1-phosphate and lysophosphatidic acid receptor. *Biochem Biophys Res Commun* 2008; 371: 707-12.
- Noguchi K, Herr D, Mutoh T, Chun J. Lysophosphatidic acid (LPA) and its receptors. *Curr Opin Pharmacol* 2009; 9: 15-23.
- Kitamura H, Okutani M, Arima N, Inoue A, Makide K, Aoki J. Identification of a novel receptor for lysophosphatidylserine. *FASEB Summer Research Conference 2009: Poster #24*.
- Olivera A, Spiegel S. Sphingosine-1-phosphate as second messenger in cell proliferation induced by PDGF and FCS mitogens. *Nature* 1993; 365: 557-60.
- Choi OH, Kim JH, Kinet JP. Calcium mobilization via sphingosine kinase in signalling by the Fc epsilon RI antigen receptor. *Nature* 1996; 380: 634-6.
- Goodemote KA, Mattie ME, Berger A, Spiegel S. Involvement of a pertussis toxin-sensitive G protein in the mitogenic signaling pathways of sphingosine 1-phosphate. *J Biol Chem* 1995; 270: 10272-7.
- van Koppen C, Meyer zu Heringdorf M, Laser KT, Zhang C, Jakobs KH, Bunemann M, et al. Activation of a high affinity Gi protein-coupled plasma membrane receptor by sphingosine-1-phosphate. *J Biol Chem* 1996; 271: 2082-7.

- 28 Bunemann M, Brandts B, zu Heringdorf DM, van Koppen CJ, Jakobs KH, Pott L. Activation of muscarinic K^+ current in guinea-pig atrial myocytes by sphingosine-1-phosphate. *J Physiol* 1995; 489: 701–7.
- 29 Okajima F, Tomura H, Sho K, Kimura T, Sato K, Im DS, *et al.* Sphingosine 1-phosphate stimulates hydrogen peroxide generation through activation of phospholipase C- Ca^{2+} system in FRTL-5 thyroid cells: possible involvement of guanosine triphosphate-binding proteins in the lipid signaling. *Endocrinology* 1997; 138: 220–9.
- 30 Im DS, Fujioka T, Katada T, Kondo Y, Ui M, Okajima F. Characterization of sphingosine 1-phosphate-induced actions and its signaling pathways in rat hepatocytes. *Am J Physiol* 1997; 272: G1091–9.
- 31 Lee MJ, Van Brocklyn JR, Thangada S, Liu CH, Hand AR, Menzeleev R, *et al.* Sphingosine-1-phosphate as a ligand for the G protein-coupled receptor EDG-1. *Science* 1998; 279: 1552–5.
- 32 An S, Bleu T, Huang W, Hallmark OG, Coughlin SR, Goetzl EJ. Identification of cDNAs encoding two G protein-coupled receptors for lysosphingolipids. *FEBS Lett* 1997; 417: 279–82.
- 33 Okamoto H, Takuwa N, Yatomi Y, Gonda K, Shigematsu H, Takuwa Y. EDG3 is a functional receptor specific for sphingosine 1-phosphate and sphingosylphosphorylcholine with signaling characteristics distinct from EDG1 and AGR16. *Biochem Biophys Res Commun* 1999; 260: 203–8.
- 34 Van Brocklyn JR, Graler MH, Bernhardt G, Hobson JP, Lipp M, Spiegel S. Sphingosine-1-phosphate is a ligand for the G protein-coupled receptor EDG-6. *Blood* 2000; 95: 2624–9.
- 35 Yamazaki Y, Kon J, Sato K, Tomura H, Sato M, Yoneya T, *et al.* Edg-6 as a putative sphingosine 1-phosphate receptor coupling to Ca^{2+} signaling pathway. *Biochem Biophys Res Commun* 2000; 268: 583–9.
- 36 Im DS, Heise CE, Ancellin N, O'Dowd BF, Shei GJ, Heavens RP, *et al.* Characterization of a novel sphingosine 1-phosphate receptor, Edg-8. *J Biol Chem* 2000; 275: 14281–6.
- 37 Lynch KR, Macdonald TL. Structure-activity relationships of lysophosphatidic acid analogs. *Biochim Biophys Acta* 2002; 1582: 289–94.
- 38 Jalink K, Hengeveld T, Mulder S, Postma FR, Simon MF, Chap H, *et al.* Lysophosphatidic acid-induced Ca^{2+} mobilization in human A431 cells: structure-activity analysis. *Biochem J* 1995; 307: 609–16.
- 39 Gueguen G, Gaige B, Grevy JM, Rogalle P, Bellan J, Wilson M, *et al.* Structure-activity analysis of the effects of lysophosphatidic acid on platelet aggregation. *Biochemistry* 1999; 38: 8440–50.
- 40 Sugjura T, Tokumura A, Gregory L, Nouchi T, Weintraub ST, Hanahan DJ. Biochemical characterization of the interaction of lipid phosphoric acids with human platelets: comparison with platelet activating factor. *Arch Biochem Biophys* 1994; 311: 358–68.
- 41 Liliom K, Bittman R, Swords B, Tigyi G. N-palmitoyl-serine and N-palmitoyl-tyrosine phosphoric acids are selective competitive antagonists of the lysophosphatidic acid receptors. *Mol Pharmacol* 1996; 50: 616–23.
- 42 Hooks SB, Ragan SP, Hopper DW, Honemann CW, Durieux ME, Macdonald TL, *et al.* Characterization of a receptor subtype-selective lysophosphatidic acid mimetic. *Mol Pharmacol* 1998; 53: 188–94.
- 43 Durgam GG, Virag T, Walker MD, Tsukahara R, Yasuda S, Liliom K, *et al.* Synthesis, structure-activity relationships, and biological evaluation of fatty alcohol phosphates as lysophosphatidic acid receptor ligands, activators of PPAR γ , and inhibitors of autotaxin. *J Med Chem* 2005; 48: 4919–30.
- 44 Lynch KR, Hopper DW, Carlisle SJ, Catalano JG, Zhang M, MacDonald TL. Structure/activity relationships in lysophosphatidic acid: the 2-hydroxyl moiety. *Mol Pharmacol* 1997; 52: 75–81.
- 45 Proll MA, Clark RB. Potent G_i -mediated inhibition of adenylyl cyclase by a phosphonate analog of monooleylphosphatidate. *Mol Pharmacol* 1991; 39: 740–4.
- 46 van der Bend RL, Brunner J, Jalink K, van Corven EJ, Moolenaar WH, van Blitterswijk WJ. Identification of a putative membrane receptor for the bioactive phospholipid, lysophosphatidic acid. *EMBO J* 1992; 11: 2495–501.
- 47 Xu Y, Casey G, Mills GB. Effect of lysophospholipids on signaling in the human Jurkat T cell line. *J Cell Physiol* 1995; 163: 441–50.
- 48 Bandoh K, Aoki J, Taira A, Tsujimoto M, Arai H, Inoue K. Lysophosphatidic acid (LPA) receptors of the EDG family are differentially activated by LPA species. Structure-activity relationship of cloned LPA receptors. *FEBS Lett* 2000; 478: 159–65.
- 49 Hasegawa Y, Erickson JR, Goddard GJ, Yu S, Liu S, Cheng KW, *et al.* Identification of a phosphothionate analogue of lysophosphatidic acid (LPA) as a selective agonist of the LPA $_3$ receptor. *J Biol Chem* 2003; 278: 11962–9.
- 50 Xu Y, Qian L, Prestwich GD. Synthesis of monofluorinated analogues of lysophosphatidic acid. *J Org Chem* 2003; 68: 5320–30.
- 51 Xu Y, Aoki J, Shimizu K, Umezu-Goto M, Hama K, Takanezawa Y, *et al.* Structure-activity relationships of fluorinated lysophosphatidic acid analogues. *J Med Chem* 2005; 48: 3319–27.
- 52 Jiang G, Xu Y, Fujiwara Y, Tsukahara T, Tsukahara R, Gajewiak J, *et al.* Alpha-substituted phosphonate analogues of lysophosphatidic acid (LPA) selectively inhibit production and action of LPA. *Chem Med Chem Lett* 2007; 2: 679–90.
- 53 Santos WL, Heasley BH, Jarosz R, Carter KM, Lynch KR, Macdonald TL. Synthesis and biological evaluation of phosphonic and thiophosphoric acid derivatives of lysophosphatidic acid. *Bioorg Med Chem Lett* 2004; 14: 3473–6.
- 54 Durgam GG, Tsukahara R, Makarova N, Walker MD, Fujiwara Y, Pigg KR, *et al.* Synthesis and pharmacological evaluation of second-generation phosphatidic acid derivatives as lysophosphatidic acid receptor ligands. *Bioorg Med Chem Lett* 2006; 16: 633–40.
- 55 Williams JR, Khandoga AL, Goyal P, Fells JI, Perygin DH, Siess W, *et al.* Unique Ligand Selectivity of the GPR92/LPA $_5$ Lysophosphatidate Receptor Indicates Role in Human Platelet Activation. *J Biol Chem* 2009; 284: 17304–19.
- 56 Virag T, Elrod DB, Liliom K, Sardar VM, Parrill AL, Yokoyama K, *et al.* Fatty alcohol phosphates are subtype-selective agonists and antagonists of lysophosphatidic acid receptors. *Mol Pharmacol* 2003; 63: 1032–42.
- 57 Gajewiak J, Tsukahara R, Tsukahara T, Fujiwara Y, Yu S, Lu Y, *et al.* Alkoxymethylenephosphonate analogues of (Lyso) phosphatidic acid stimulate signaling networks coupled to the LPA $_2$ receptor. *Chem Med Chem Lett* 2007; 2: 1789–98.
- 58 Tamaruya Y, Suzuki M, Kamura G, Kanai M, Hama K, Shimizu K, *et al.* Identifying specific conformations by using a carbohydrate scaffold: discovery of subtype-selective LPA-receptor agonists and an antagonist. *Angew Chem Int Ed Engl* 2004; 43: 2834–7.
- 59 Gududuru V, Zeng K, Tsukahara R, Makarova N, Fujiwara Y, Pigg KR, *et al.* Identification of Darmstoff analogs as selective agonists and antagonists of lysophosphatidic acid receptors. *Bioorg Med Chem Lett* 2006; 16: 451–6.
- 60 Xu Y, Tanaka M, Arai H, Aoki J, Prestwich GD. Alkyl lysophosphatidic acid and fluoromethylene phosphonate analogs as metabolically-stabilized agonists for LPA receptors. *Bioorg Med Chem Lett* 2004; 14: 5323–8.
- 61 Hooks SB, Santos WL, Im DS, Heise CE, Macdonald TL, Lynch KR. Lysophosphatidic acid-induced mitogenesis is regulated by lipid phosphate phosphatases and is Edg-receptor independent. *J Biol Chem* 2001; 276: 4611–21.
- 62 Oh DY, Yoon JM, Moon MJ, Hwang JI, Choe H, Lee JY, *et al.* Identifica-

- tion of farnesyl pyrophosphate and N-arachidonylglycine as endogenous ligands for GPR92. *J Biol Chem* 2008; 283: 21054–64.
- 63 Baker DL, Fujiwara Y, Pigg KR, Tsukahara R, Kobayashi S, Murofushi H, et al. Carba analogs of cyclic phosphatidic acid are selective inhibitors of autotaxin and cancer cell invasion and metastasis. *J Biol Chem* 2006; 281: 22786–93.
- 64 Qian L, Xu Y, Simper T, Jiang G, Aoki J, Umezue-Goto M, et al. Phosphorothioate analogues of alkyl lysophosphatidic acid as LPA₃ receptor-selective agonists. *ChemMedChem* 2006; 1: 376–83.
- 65 Endo T, Kano K, Motoki R, Hama K, Okudaira S, Ishida M, et al. Lysophosphatidylmethanol is a pan lysophosphatidic acid receptor agonist and is produced by autotaxin in blood. *J Biochem* 2009; 146: 283–93.
- 66 Heise CE, Santos WL, Schreihof AM, Heasley BH, Mukhin YV, Macdonald TL, et al. Activity of 2-substituted lysophosphatidic acid (LPA) analogs at LPA receptors: discovery of a LPA₁/LPA₃ receptor antagonist. *Mol Pharmacol* 2001; 60: 1173–80.
- 67 Heasley BH, Jarosz R, Lynch KR, Macdonald TL. Initial structure-activity relationships of lysophosphatidic acid receptor antagonists: discovery of a high-affinity LPA₁/LPA₃ receptor antagonist. *Bioorg Med Chem Lett* 2004; 14: 2735–40.
- 68 Heasley BH, Jarosz R, Carter KM, Van SJ, Lynch KR, Macdonald TL. A novel series of 2-pyridyl-containing compounds as lysophosphatidic acid receptor antagonists: development of a nonhydrolyzable LPA₃ receptor-selective antagonist. *Bioorg Med Chem Lett* 2004; 14: 4069–74.
- 69 Fischer DJ, Nusser N, Virag T, Yokoyama K, Wang D, Baker DL, et al. Short-chain phosphatidates are subtype-selective antagonists of lysophosphatidic acid receptors. *Mol Pharmacol* 2001; 60: 776–84.
- 70 Ohta H, Sato K, Murata N, Damirin A, Malchinkhuu E, Kon J, et al. Ki16425, a subtype-selective antagonist for EDG-family lysophosphatidic acid receptors. *Mol Pharmacol* 2003; 64: 994–1005.
- 71 Liliom K, Tsukahara T, Tsukahara R, Zelman-Femiak M, Swiezewska E, Tigyi G. Farnesyl phosphates are endogenous ligands of lysophosphatidic acid receptors: inhibition of LPA GPCR and activation of PPARs. *Biochim Biophys Acta* 2006; 1761: 1506–14.
- 72 Fells JI, Tsukahara R, Fujiwara Y, Liu J, Perygin DH, Osborne DA, et al. Identification of non-lipid LPA₃ antagonists by virtual screening. *Bioorg Med Chem* 2008; 16: 6207–17.
- 73 Prestwich GD, Gajewiak J, Zhang H, Xu X, Yang G, Serban M. Phosphatase-resistant analogues of lysophosphatidic acid: agonists promote healing, antagonists and autotaxin inhibitors treat cancer. *Biochim Biophys Acta* 2008; 1781: 588–94.
- 74 Im DS, Clemens J, Macdonald TL, Lynch KR. Characterization of the human and mouse sphingosine 1-phosphate receptor, S1P₅ (Edg-8): structure-activity relationship of sphingosine 1-phosphate receptors. *Biochemistry* 2001; 40: 14053–60.
- 75 Ancellin N, Hla T. Differential pharmacological properties and signal transduction of the sphingosine 1-phosphate receptors EDG-1, EDG-3, and EDG-5. *J Biol Chem* 1999; 274: 18997–9002.
- 76 Candelore MR, Wright MJ, Tota LM, Milligan J, Shei GJ, Bergstrom JD, et al. Phytosphingosine 1-phosphate: a high affinity ligand for the S1P₄/Edg-6 receptor. *Biochem Biophys Res Commun* 2002; 297: 600–6.
- 77 Lim HS, Park JJ, Ko K, Lee MH, Chung SK. Syntheses of sphingosine-1-phosphate analogues and their interaction with EDG/S1P receptors. *Bioorg Med Chem Lett* 2004; 14: 2499–503.
- 78 Mandala S, Hajdu R, Bergstrom J, Quackenbush E, Xie J, Milligan J, et al. Alteration of lymphocyte trafficking by sphingosine-1-phosphate receptor agonists. *Science* 2002; 296: 346–9.
- 79 Brinkmann V, Davis MD, Heise CE, Albert R, Cottens S, Hof R, et al. The immune modulator FTY720 targets sphingosine 1-phosphate receptors. *J Biol Chem* 2002; 277: 21453–7.
- 80 Im DS. Linking Chinese medicine and G-protein-coupled receptors. *Trends Pharmacol Sci* 2003; 24: 2–4.
- 81 Sanna MG, Liao J, Jo E, Alfonso C, Ahn MY, Peterson MS, et al. Sphingosine 1-phosphate (S1P) receptor subtypes S1P₁ and S1P₃, respectively, regulate lymphocyte recirculation and heart rate. *J Biol Chem* 2004; 279: 13839–48.
- 82 Jo E, Sanna MG, Gonzalez-Cabrera PJ, Thangada S, Tigyi G, Osborne DA, et al. S1P₁-selective *in vivo* – active agonists from high-throughput screening: off-the-shelf chemical probes of receptor interactions, signaling, and fate. *Chem Biol* 2005; 12: 703–15.
- 83 Li Z, Chen W, Hale JJ, Lynch CL, Mills SG, Hajdu R, et al. Discovery of potent 3,5-diphenyl-1,2,4-oxadiazole sphingosine-1-phosphate-1 (S1P₁) receptor agonists with exceptional selectivity against S1P₂ and S1P₃. *J Med Chem* 2005; 48: 6169–73.
- 84 Schurer SC, Brown SJ, Gonzalez-Cabrera PJ, Schaeffer MT, Chapman J, Jo E, et al. Ligand-binding pocket shape differences between sphingosine 1-phosphate (S1P) receptors S1P₁ and S1P₃ determine efficiency of chemical probe identification by ultrahigh-throughput screening. *ACS Chem Biol* 2008; 3: 486–98.
- 85 Gonzalez-Cabrera PJ, Jo E, Sanna MG, Brown S, Leaf N, Marsolais D, et al. Full pharmacological efficacy of a novel S1P₁ agonist that does not require S1P-like headgroup interactions. *Mol Pharmacol* 2008; 74: 1308–18.
- 86 Pan S, Mi Y, Pally C, Beerli C, Chen A, Guerini D, et al. A monoselective sphingosine-1-phosphate receptor-1 agonist prevents allograft rejection in a stringent rat heart transplantation model. *Chem Biol* 2006; 13: 1227–34.
- 87 Zhu R, Snyder AH, Kharel Y, Schaffter L, Sun Q, Kennedy PC, et al. Asymmetric synthesis of conformationally constrained fingolimod analogues – discovery of an orally active sphingosine 1-phosphate receptor type-1 agonist and receptor type-3 antagonist. *J Med Chem* 2007; 50: 6428–35.
- 88 Fujishiro J, Kudou S, Iwai S, Takahashi M, Hakamata Y, Kinoshita M, et al. Use of sphingosine-1-phosphate 1 receptor agonist, KRP-203, in combination with a subtherapeutic dose of cyclosporine A for rat renal transplantation. *Transplantation* 2006; 82: 804–12.
- 89 Song J, Matsuda C, Kai Y, Nishida T, Nakajima K, Mizushima T, et al. A novel sphingosine 1-phosphate receptor agonist, 2-amino-2-propanediol hydrochloride (KRP-203), regulates chronic colitis in interleukin-10 gene-deficient mice. *J Pharmacol Exp Ther* 2008; 324: 276–83.
- 90 Hanessian S, Charron G, Billich A, Guerini D. Constrained azacyclic analogues of the immunomodulatory agent FTY720 as molecular probes for sphingosine 1-phosphate receptors. *Bioorg Med Chem Lett* 2007; 17: 491–4.
- 91 Himmel HM, Meyer Zu Heringdorf D, Graf E, Dobrev D, Kortner A, Schuler S, et al. Evidence for Edg-3 receptor-mediated activation of I(K.ACh) by sphingosine-1-phosphate in human atrial cardiomyocytes. *Mol Pharmacol* 2000; 58: 449–54.
- 92 Niedernberg A, Scherer CR, Busch AE, Kostenis E. Comparative analysis of human and rat S1P₅ (edg8): differential expression profiles and sensitivities to antagonists. *Biochem Pharmacol* 2002; 64: 1243–50.
- 93 Ikeda H, Satoh H, Yanase M, Inoue Y, Tomiya T, Arai M, et al. Antiproliferative property of sphingosine 1-phosphate in rat hepatocytes involves activation of Rho via Edg-5. *Gastroenterology* 2003; 124: 459–69.
- 94 Ohmori T, Yatomi Y, Osada M, Kazama F, Takafuta T, Ikeda H, et al. Sphingosine 1-phosphate induces contraction of coronary artery

- smooth muscle cells via S1P₂. *Cardiovasc Res* 2003; 58: 170–7.
- 95 Davis MD, Clemens JJ, Macdonald TL, Lynch KR. Sphingosine 1-phosphate analogs as receptor antagonists. *J Biol Chem* 2005; 280: 9833–41.
- 96 Sanna MG, Wang SK, Gonzalez-Cabrera PJ, Don A, Marsolais D, Matheu MP, et al. Enhancement of capillary leakage and restoration of lymphocyte egress by a chiral S1P₁ antagonist *in vivo*. *Nat Chem Biol* 2006; 2: 434–41.
- 97 Foss FW Jr, Snyder AH, Davis MD, Rouse M, Okusa MD, Lynch KR, et al. Synthesis and biological evaluation of gamma-aminophosphonates as potent, subtype-selective sphingosine 1-phosphate receptor agonists and antagonists. *Bioorg Med Chem* 2007; 15: 663–77.
- 98 Waters CM, Long J, Gorshkova I, Fujiwara Y, Connell M, Belmonte KE, et al. Cell migration activated by platelet-derived growth factor receptor is blocked by an inverse agonist of the sphingosine 1-phosphate receptor-1. *FASEB J* 2006; 20: 509–11.
- 99 Yonesu K, Ohnuki T, Ono Y, Takatsu T, Nara F. Ascotricins A and B, novel antagonists of sphingosine-1-phosphate receptor 1 from *Ascotricha chartarum* Berk. SANK 14186. *J Antibiot (Tokyo)* 2009; 62: 359–64.
- 100 Yonesu K, Kawase Y, Inoue T, Takagi N, Tsuchida J, Takuwa Y, et al. Involvement of sphingosine-1-phosphate and S1P₁ in angiogenesis: analyses using a new S1P₁ antagonist of non-sphingosine-1-phosphate analog. *Biochem Pharmacol* 2009; 77: 1011–20.
- 101 Goetzl EJ, Dembrow D, Van Brocklyn JR, Graler M, Huang MC. An IgM-kappa rat monoclonal antibody specific for the type 1 sphingosine 1-phosphate G protein-coupled receptor with antagonist and agonist activities. *Immunol Lett* 2004; 93: 63–9.
- 102 Koide Y, Hasegawa T, Takahashi A, Endo A, Mochizuki N, Nakagawa M, et al. Development of novel EDG3 antagonists using a 3D database search and their structure-activity relationships. *J Med Chem* 2002; 45: 4629–38.
- 103 Jongsma M, Hendriks-Balk MC, Michel MC, Peters SL, Alewijnse AE. BML-241 fails to display selective antagonism at the sphingosine-1-phosphate receptor, S1P₃. *Br J Pharmacol* 2006; 149: 277–82.
- 104 Koide Y, Uemoto K, Hasegawa T, Sada T, Murakami A, Takasugi H, et al. Pharmacophore-based design of sphingosine 1-phosphate-3 receptor antagonists that include a 3,4-dialkoxybenzophenone scaffold. *J Med Chem* 2007; 50: 442–54.
- 105 Zhang H, Desai NN, Olivera A, Seki T, Brooker G, Spiegel S. Sphingosine-1-phosphate, a novel lipid, involved in cellular proliferation. *J Cell Biol* 1991; 114: 155–67.
- 106 Rosen H, Gonzalez-Cabrera PJ, Sanna MG, Brown S. Sphingosine 1-phosphate receptor signaling. *Annu Rev Biochem* 2009; 78: 743–68.
- 107 Rivera R, Chun J. Biological effects of lysophospholipids. *Rev Physiol Biochem Pharmacol* 2008; 160: 25–46.
- 108 Skoura A, Hla T. Lysophospholipid receptors in vertebrate development, physiology, and pathology. *J Lipid Res* 2009; 50: S293–8.
- 109 Ishii I, Fukushima N, Ye X, Chun J. Lysophospholipid receptors: signaling and biology. *Annu Rev Biochem* 2004; 73: 321–54.
- 110 Brinkmann V, Lynch KR. FTY720: targeting G-protein-coupled receptors for sphingosine 1-phosphate in transplantation and autoimmunity. *Curr Opin Immunol* 2002; 14: 569–75.
- 111 Brown BA, Kantesaria PP, McDevitt LM. Fingolimod: a novel immunosuppressant for multiple sclerosis. *Ann Pharmacother* 2007; 41: 1660–8.
- 112 Swaney JS, Chapman C, Correa LD, Bain G, Bunday RA, Prodanovich PP, et al. A novel, small molecule, LPA₁ receptor-selective antagonist, inhibits bleomycin-induced inflammation and fibrosis in the mouse. *FASEB Summer Research Conference*. 2009.
- 113 Nagiec M, Funckes-Shippy C, Hiesch R, Radi Z, Walker J, Gierse J, et al. Targeting S1P signaling for inflammation. *FASEB Summer Research Conference*. 2009.
- 114 Sabbadini R, Cavalli A, Wocjiak J, Visentin B, Matteo R, Campbell M, et al. Antibodies against bioactive lipids. *FASEB Summer Research Conference*. 2009.
- 115 Caballero S, Swaney J, Moreno K, Afzal A, Kielczewski J, Stoller G, et al. Anti-sphingosine-1-phosphate monoclonal antibodies inhibit angiogenesis and sub-retinal fibrosis in a murine model of laser-induced choroidal neovascularization. *Exp Eye Res* 2009; 88: 367–77.
- 116 Prestwich GD. Dual activity LPA receptor pan-antagonist/autotaxin inhibitors as anti-cancer agents: evaluation *in vivo* using engineered human tumors. *FASEB Summer Research Conference*. 2009.
- 117 Zhang H, Xu X, Gajewiak J, Tsukahara R, Fujiwara Y, Liu J, et al. Dual activity lysophosphatidic acid receptor pan-antagonist/autotaxin inhibitor reduces breast cancer cell migration *in vitro* and causes tumor regression *in vivo*. *Cancer Res* 2009; 69: 5441–9.
- 118 Goetzl EJ, Lynch KR. Preface: the omnific lysophospholipid growth factors. *Ann N Y Acad Sci* 2000; 905: xi–xiv.
- 119 Chun J, Goetzl EJ, Hla T, Igarashi Y, Lynch KR, Moolenaar W, et al. International Union of Pharmacology. XXXIV. Lysophospholipid Receptor Nomenclature. *Pharmacol Rev* 2002; 54: 265–9.
- 120 Fells JI, Tsukahara R, Liu J, Tigyi G, Parrill AL. Structure-based drug design identifies novel LPA₃ antagonists. *Bioorg Med Chem* 2009; 17: 7457–64.
- 121 Bolli MH, Abele S, Binkert C, Bravo R, Buchmann S, Bur D, et al. 2-imino-thiazolidin-4-one derivatives as potent, orally active S1P₁ receptor agonists. *J Med Chem* 2010; 53: 4198–211.
- 122 An S, Bleu T, Zheng Y, Goetzl EJ. Recombinant human G protein-coupled lysophosphatidic acid receptors mediate intracellular calcium mobilization. *Mol Pharmacol* 1998; 54: 881–8.
- 123 Murakami A, Takasugi H, Ohnuma S, Koide Y, Sakurai A, Takeda S, et al. Sphingosine 1-phosphate (S1P) regulates vascular contraction via S1P₃ receptor: investigation based on a new S1P₃ receptor antagonist. *Mol Pharmacol* 2010; 77: 704–13.
- 124 Kim K, Kim YL, Sackett SJ, Kim HL, Han M, Park DS, et al. Sphingosine 1-phosphate (S1P) induces shape change in rat C6 glioma cells through the S1P₂ receptor: development of an agonist for S1P receptors. *J Pharm Pharmacol* 2007; 59: 1035–41.