

Original Article

# Novel $\kappa$ -opioid receptor agonist MB-1C-OH produces potent analgesia with less depression and sedation

Le-sha ZHANG<sup>1</sup>, Jun WANG<sup>1,2</sup>, Jian-chun CHEN<sup>3</sup>, Yi-min TAO<sup>1</sup>, Yu-hua WANG<sup>2,4</sup>, Xue-jun XU<sup>1</sup>, Jie CHEN<sup>1</sup>, Yun-gen XU<sup>2</sup>, Tao XI<sup>2</sup>, Xiao-wu HU<sup>3,\*</sup>, Yu-jun WANG<sup>1,\*</sup>, Jing-gen LIU<sup>1</sup>

<sup>1</sup>State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica and Collaborative Innovation Center for Brain Science, Chinese Academy of Sciences, Shanghai 201203, China; <sup>2</sup>Biotechnology Center, School of Life Science and Technology, China Pharmaceutical University, Nanjing 210009, China; <sup>3</sup>Department of Neurosurgery, Changhai Hospital, Second Military Medical University, Shanghai 200433, China; <sup>4</sup>School of Pharmacy, Nanjing University of Chinese Medicine, Nanjing 210046, China

**Aim:** To characterize the pharmacological profiles of a novel  $\kappa$ -opioid receptor agonist MB-1C-OH.

**Methods:** [<sup>3</sup>H]diprenorphine binding and [<sup>35</sup>S]GTP $\gamma$ S binding assays were performed to determine the agonistic properties of MB-1C-OH. Hot plate, tail flick, acetic acid-induced writhing, and formalin tests were conducted in mice to evaluate the antinociceptive actions. Forced swimming and rotarod tests of mice were used to assess the sedation and depression actions.

**Results:** In [<sup>3</sup>H]diprenorphine binding assay, MB-1C-OH did not bind to  $\mu$ - and  $\delta$ -opioid receptors at the concentration of 100  $\mu$ mol/L, but showed a high affinity for  $\kappa$ -opioid receptor ( $K_i=35$  nmol/L). In [<sup>35</sup>S]GTP $\gamma$ S binding assay, the compound had an  $E_{max}$  of 98% and an  $EC_{50}$  of 16.7 nmol/L for  $\kappa$ -opioid receptor. Subcutaneous injection of MB-1C-OH had no effects in both hot plate and tail flick tests, but produced potent antinociception in the acetic acid-induced writhing test ( $ED_{50}=0.39$  mg/kg), which was antagonized by pretreatment with a selective  $\kappa$ -opioid receptor antagonist Nor-BNI. In the formalin test, subcutaneous injection of MB-1C-OH did not affect the flinching behavior in the first phase, but significantly inhibited that in the second phase ( $ED_{50}=0.87$  mg/kg). In addition, the sedation or depression actions of MB-1C-OH were about 3-fold weaker than those of the classical  $\kappa$  agonist (-)U50,488H.

**Conclusion:** MB-1C-OH is a novel  $\kappa$ -opioid receptor agonist that produces potent antinociception causing less sedation and depression.

**Keywords:**  $\kappa$ -opioid receptor; MB-1C-OH; (-)U50,488H; Nor-BNI; morphine; pain; acetic acid-induced writhing; formalin test; depression; sedation

Acta Pharmacologica Sinica (2015) 36: 565–571; doi: 10.1038/aps.2014.145; published online 30 Mar 2015

## Introduction

Opioid receptors have been classified into  $\mu$ -,  $\kappa$ -, and  $\delta$ -opioid receptors based on their different pharmacological profiles<sup>[1]</sup>. Currently, most opioid analgesics, such as morphine, mainly induce analgesia via binding to the  $\mu$ -opioid receptor, but they are also associated with a spectrum of undesirable side effects, such as physical dependence, respiratory depression, and constipation<sup>[2]</sup>. Numerous studies have demonstrated that drugs targeting  $\kappa$ -opioid receptor produced potent antinociception with less adverse and fewer addictive side effects than  $\mu$ -opioid receptor agonists<sup>[3–5]</sup>. Endoh *et al* reported that TRK-820, a  $\kappa$ -opioid receptor agonist, was 68- to 328-fold more potent than (-)U50,488H, and 41- to 349-fold more potent than


morphine in producing antinociception<sup>[6]</sup>. We previously found that LPK-26 produced potent antinociceptive effects without inducing physical dependence<sup>[7]</sup>. However, selective  $\kappa$ -opioid agonists frequently exhibit severe central nervous system side effects such as sedation and depression, which limit their clinical use<sup>[8]</sup>.

Increasing evidence from a wide variety of visceral pain models, inflammatory pain models and thermal hyperalgesia induced by capsaicin show that  $\kappa$ -opioid agonists could produce their analgesic effects via peripheral targeting<sup>[9–13]</sup>. Thus, peripherally acting  $\kappa$ -opioid agonists may be more promising candidate pharmacotherapies for pain relief because they do not cause central nervous system side effects.

In this study, a novel synthetic compound, MB-1C-OH (empirical formula: C<sub>24</sub>H<sub>25</sub>ClN<sub>2</sub>O<sub>3</sub>, molecular weight: 424), in which a hydroxyl group was introduced in the parent framework of MB-1C, was examined (Figure 1). MB-1C-OH was characterized using *in vitro* and *in vivo* experiments. Specifi-

\* To whom correspondence should be addressed.  
E-mail yjwang@mail.simm.ac.cn (Yu-jun WANG);  
huxiaowu25@sina.com (Xiao-wu HU)  
Received 2014-08-09 Accepted 2014-11-10

cally, MB-1C-OH's binding affinity for opioid receptors ( $\mu$ ,  $\kappa$ , and  $\delta$ ) and ability to stimulate guanosine5'-O-(3-[ $^{35}$ S]thio) triphosphate ([ $^{35}$ S]GTP $\gamma$ S) binding to G-proteins were determined. The antinociceptive effects of MB-1C-OH were evaluated in the hot plate, tail flick, acetic acid-induced writhing and formalin tests.



**Figure 1.** Chemical structures of (-)U50,488H and MB-1C-OH.

## Materials and methods

### Cell culture

Chinese hamster ovary (CHO) cells were transfected with human  $\kappa$ -, rat  $\mu$ -, or rat  $\delta$ -opioid receptors using Lipofectamine (Invitrogen) as per the manufacturer's protocol. CHO cells stably expressing human  $\kappa$ -, rat  $\mu$ -, or rat  $\delta$ -opioid receptors were maintained in F12 medium (Gibco) with 10% fetal calf serum and 0.25 mg/mL G418 (Roche). Cells were incubated in a humidified atmosphere consisting of 5% CO<sub>2</sub>, 95% air at 37°C. For the receptor binding assay and the [ $^{35}$ S]GTP $\gamma$ S binding assay, cells were seeded into 175-cm<sup>3</sup> flasks. When cell growth reached 70% confluence, cells were washed with phosphate-buffered saline and the membrane was prepared as follows.

### Cell membrane preparation

CHO cells were detached by incubation with phosphate-buffered saline containing 1 mmol/L EDTA and centrifuged at 1000×g for 10 min. The cell pellet was suspended in ice-cold homogenization buffer composed of 50 mmol/L HEPES, pH 7.4; 1 mmol/L MgCl<sub>2</sub>; and 1 mmol/L EGTA. Cells were homogenized with 10 strokes using a glass Dounce homogenizer. After centrifugation at 40000×g for 10 min (4°C), pellets were resuspended in homogenization buffer, homogenized, and centrifuged again as described. This procedure was repeated twice more. The final pellets were resuspended in 50 mmol/L Tris-HCl buffer, pH 7.4. The protein concentration was determined, and aliquots were stored at -80°C.

### Receptor binding assay

Ligand binding experiments were performed with [ $^3$ H]diprenorphine for opioid receptors. Competition inhibition of [ $^3$ H]diprenorphine binding to opioid receptors by MB-1C-OH or (-)U50,488H was performed in the absence or presence at various concentration of each drug. The receptor binding assay was carried out in triplicate using 50 mmol/L Tris-

HCl buffer (pH 7.4) at 37°C for 30 min in a final volume of 0.5 mL with 30  $\mu$ g of membrane protein. Naloxone (10  $\mu$ mol/L, Sigma) was used to define nonspecific binding. Bound and free [ $^3$ H]diprenorphine were separated by filtration under reduced pressure with GF/B filters (Whatman). The amount of radioactivity on the filters was determined using a liquid scintillation counter (Beckman LS6500).

### GTP $\gamma$ S binding assay

[ $^{35}$ S]GTP $\gamma$ S binding was performed as described previously<sup>[14,15]</sup>. Briefly, membranes (15  $\mu$ g/sample) were incubated with 0.1 nmol/L [ $^{35}$ S]GTP $\gamma$ S in a binding buffer composed of 50 mmol/L Tris-HCl, pH 7.5; 1 mmol/L EDTA; 5 mmol/L MgCl<sub>2</sub>; 100 mmol/L NaCl; and 40  $\mu$ mol/L GDP at 30°C for 1 h in the presence of increasing concentrations of MB-1C-OH. Nonspecific binding was determined in the presence of non-radioactive GTP $\gamma$ S (10  $\mu$ mol/L). Reactions were terminated by rapid filtration, and the amount of bound radioactivity was determined by liquid scintillation counting as described above. The percentage of stimulated [ $^{35}$ S]GTP $\gamma$ S binding was calculated as  $100 \times (\text{cpm}_{\text{sample}} - \text{cpm}_{\text{nonspecific}}) / (\text{cpm}_{\text{basal}} - \text{cpm}_{\text{nonspecific}})$ .

### Animals

Male Kunming strain mice (approximately 20 g) were obtained from the Laboratory Animal Center, Chinese Academy of Sciences (Shanghai, China). Mice were housed in groups and maintained in a 12/12 h light/dark cycle in a temperature controlled environment with free access to food and water. Ten to fifteen mice were used per treatment group. All animal treatments were strictly in accordance with the institutional guidelines of Animal Care and Use Committee, Shanghai Institute of Materia Medica, Chinese Academy of Sciences.

### Hot plate test

The hot plate test was performed according to the method we described previously<sup>[4,16]</sup>. Briefly, the hot plate temperature was set at 55°C. Mice were placed on the heated smooth surface. The amount of time that elapsed before the mice showed the first signs of discomfort (hind-paw lifting, licking or shaking, and jumping) was recorded. Prior to drug administration, the nociceptive response for each mouse was measured two times. The mean of these two times was used as the basal response time for each mouse. A cut-off time of 60 s was used in the test to avoid tissue damage. Each test was undertaken 15 min after the drug was administered. The Maximum Possible Effect (% MPE) was calculated as:  $100 \times (\text{response time after drug treatment} - \text{basal response time before drug treatment}) / (\text{cut-off time} - \text{basal latency before response time without drug treatment})$ .

### Acetic acid-induced writhing test

The acetic acid-induced writhing test was performed in mice according to our previously reported procedure<sup>[4,16]</sup>. Abdominal constriction was induced by the injection of 0.6% acetic acid (10 mL/kg, ip). An abdominal constriction was defined as a wave of contraction of the abdominal musculature fol-

lowed by extension of the hind limbs. Acetic acid solution was injected ip 15 min after the administration of the drug, and the number of abdominal constrictions was counted for 15 min after acetic acid administration. The Maximum Possible Effect (% MPE) was calculated as:  $100 \times (\text{mean abdominal constriction in vehicle treated group} - \text{constriction times in drug-treated animal}) / \text{mean abdominal constriction in vehicle treated group}$ . In all experiments with mice, drugs were given by a subcutaneous (sc) route prior to acetic acid administration. To determine the *in vivo* agonist profile of MB-1C-OH, mice were pretreated with a  $\kappa$ -(or-binaltorphimine (Nor-BNI), 10 mg/kg, sc, -15 min)<sup>[7,16,17]</sup>,  $\mu$ - ( $\beta$ -funaltrexamine ( $\beta$ -FNA), 10 mg/kg, sc, -24 h)<sup>[18,19]</sup> or  $\delta$ -(naltrindole, 3.0 mg/kg, sc, -30 min)<sup>[20]</sup> opioid receptor antagonist before MB-1C-OH administration.

#### Tail flick test

The tail-flick assay was performed according to our previously reported procedure<sup>[4,16]</sup>. Briefly, a beam of light was focused on the dorsal surface of the tail approximately 1–2 cm from the tip, and the amount of time that it took the mouse to withdraw its tail from the noxious stimulus was recorded. Before drug administration, the nociceptive response of each mouse was measured twice with a between trial interval of 5 min, and the mean of the two trials was recorded as the basal response time without drug treatment. Mice that did not have a basal response time between 2 and 7 s were excluded from further studies. A cut-off time of 14 s was employed as the maximum possible duration in the test after drug treatment in order to prevent tissue damage. The test was conducted 15 min after the administration of the drug. The Maximum Possible Effect (% MPE) was calculated as follows:  $100 \times (\text{response time after drug treatment} - \text{basal response time before drug treatment}) / (\text{cut-off time} - \text{basal latency before drug treatment})$ .

#### Formalin test

The formalin test was performed according to the method described previously<sup>[21]</sup>. Briefly, 20  $\mu$ L of 1.0% formalin was injected to the plantar surface of the right hind paw. The mice were observed for 60 min after the injection of formalin, and the amount of time that the mice spent licking or flinching the injected hind paw was recorded. The first 10 min after an injection of formalin was recorded as the early phase, and the period between 10 min and 60 min was recorded as the late phase. The drugs were administered 15 min before the injection of formalin. The total time that the mice spent licking or flinching the injured paw (pain behavior) was measured with a stopwatch.

#### Forced swimming test (FST)

The forced swimming test was performed according to the method described previously<sup>[22]</sup>. Mice were put singly into transparent Plexiglas cylinders (height: 24 cm, diameter: 13 cm) containing water at a depth of 10 cm, maintained at 22°C for 6 min. The amount of time that the mice remained immobile was recorded in seconds during the final 4 min of the test. The percentage of depression was calculated

as:  $100 \times (\text{immobility time}) / 240$ , and then, the ED<sub>50</sub> value was calculated using software which was based on the Litchfield and Wilcoxon method<sup>[23]</sup>.

#### Rotarod test

The rotarod test was conducted using a procedure described previously<sup>[16,24]</sup>. The maximum time tested was prolonged from 60 s to 300 s. In brief, the machine was set at a fixed rotational rate of 8 rounds per minute. The animals were trained twice a day for three consecutive days to maintain their position for 300 s without falling off the rotarod. The formal test began 15 min after the administration of the drug. The duration time for each mouse to remain on the Rotarod was recorded in seconds. The percentage of sedation was calculated as  $100 \times (300 - \text{duration time after test drug treatment}) / 300$  and was used to determine the ED<sub>50</sub> values of the test drugs.

#### Drugs

[<sup>3</sup>H]diprenorphine (1.85 TBq/mmol) and guanosine 5-O-(3-[<sup>35</sup>S]thio)triphosphate ([<sup>35</sup>S]GTP $\gamma$ S) (38.11 TBq/mmol) were purchased from Amersham Biosciences (Piscataway, NJ, USA). (-)U50,488H; naloxone; naltrindole; nor-binaltorphimine;  $\beta$ -funaltrexamine; GTP $\gamma$ S and GDP were obtained from Sigma-Aldrich (St Louis, MO, USA). Morphine hydrochloride was purchased from Qinghai Pharmaceutical General Factory (Xining, China).

#### Statistical analysis

Curve-fitting analyses were performed using GraphPad Prism 4.02 software. Results are expressed as the mean  $\pm$  SEM of at least three separate experiments. Statistical significance was determined using ANOVA (one- or two-way) followed by *post hoc* comparison using Dunnett's tests. When only two groups were compared, statistical significance was determined using unpaired Student's *t*-tests.

## Results

#### Affinity, selectivity, and efficacy of MB-1C-OH

Competitive inhibition of [<sup>3</sup>H]diprenorphine binding to opioid receptors by MB-1C-OH was performed to examine the binding affinities of MB-1C-OH to the opioid receptors ( $\mu$ ,  $\kappa$ , and  $\delta$ ). The K<sub>i</sub> of MB-1C-OH was 35 nmol/L to inhibit the binding of [<sup>3</sup>H]diprenorphine to the  $\kappa$ -opioid receptor, whereas MB-1C-OH at concentrations as high as 100  $\mu$ mol/L showed no affinity for the  $\mu$ - or  $\delta$ -opioid receptors (Table 1). In the [<sup>35</sup>S]GTP $\gamma$ S binding assay, MB-1C-OH had an E<sub>max</sub> (percentage of maximal stimulation) of 98.19% for the  $\kappa$ -opioid receptor. The EC<sub>50</sub> of MB-1C-OH to stimulate [<sup>35</sup>S]GTP $\gamma$ S binding to the  $\kappa$ -opioid receptor was 16.7 nmol/L (Table 1).

#### Antinociceptive effects of MB-1C-OH were mediated by the $\kappa$ -opioid receptor

As shown in Table 2, in the hot plate and tail flick tests, MB-1C-OH had no significant antinociceptive effects. MB-1C-OH (10 mg/kg, sc) did not produce significant analgesic effects in

both mouse hot plate and tail flick tests, which was similar to the data that we got from dose of 6 mg/kg (data not shown). Morphine and (-)U50,488H produced antinociception in the hot plate test with ED<sub>50</sub> values of 6.95 and 4.42 mg/kg, respectively, and in the tail flick test with ED<sub>50</sub> values of 3.46 and 5.02 mg/kg, respectively.

In the mouse acetic acid-induced writhing test, MB-1C-OH produced significant antinociception with an ED<sub>50</sub> value of 0.39 mg/kg, which was approximately 2.3-fold smaller than the ED<sub>50</sub> for (-)U50,488H (0.89 mg/kg), and 2.1-fold smaller than the ED<sub>50</sub> for morphine (0.84 mg/kg) (Table 2). The antinociceptive effect of MB-1C-OH in the acetic acid-induced writhing test was also completely antagonized by the  $\kappa$ -opioid receptor antagonist nor-BNI but not by the  $\mu$ - or  $\delta$ -opioid receptor antagonists (Figure 2). One-way ANOVA tests ( $F_{3,36}=22.34$ ,  $P<0.01$ ) and the following *post hoc* comparison using Dunnett's tests ( $P<0.01$ ) revealed a significant effect of the ability of nor-BNI to inhibit the antinociceptive effect of MB-1C-OH. These results suggested that the  $\kappa$ -opioid receptor is involved in MB-1C-OH-mediated antinociceptive effects.

In the formalin test, MB-1C-OH dose-dependently inhibited formalin-induced pain only in the second phase with an ED<sub>50</sub> of 0.87 mg/kg (Figure 3, Table 3), which was similar to the pattern of antinociception produced by (-)U50,488H but different from the antinociception produced by morphine. In Figure 3, one-way ANOVA tests ( $F_{4,45}=22.53$ ,  $P<0.01$ ) and the following *post hoc* comparison using Dunnett's tests revealed that doses of 1.25 mg/kg ( $P<0.01$ ), 2.5 mg/kg ( $P<0.01$ ) and

6.25 mg/kg ( $P<0.01$ ) of MB-1C-OH could significantly inhibit formalin-induced pain in second phase. As shown in Table 3, (-)U50,488H produced antinociception on the second phase with an ED<sub>50</sub> of 0.41 mg/kg. Morphine acted on both phases with ED<sub>50</sub> values of 5.05 mg/kg in the first phase and 3.06 mg/kg in the second phase.

#### Depressive side effect of MB-1C-OH in the forced swimming test

MB-1C-OH and (-)U50,488H dose-dependently increased the immobility time during the forced swimming test (Figure 4A). As shown in Table 4, the depressive ED<sub>50</sub> of MB-1C-OH in the FST was 9.49 mg/kg, which was 4 times higher than that of (-)U50,488H (2.35 mg/kg). The ratio of the depressive ED<sub>50</sub> to the antinociceptive ED<sub>50</sub> of MB-1C-OH was 24.33, which was 9.2 times higher than that of (-)U50,488H (2.64), suggesting that MB-1C-OH has a lower potential to cause depression than the classical  $\kappa$ -opioid agonist (-)U50,488H.

#### Sedative side effect of MB-1C-OH in the rotarod test

$\kappa$ -Opioid agonists often produce the undesirable side effect of sedation in therapeutic use. MB-1C-OH also caused a dose-dependent inhibition in rotarod performance (Figure 4B). As shown in Table 4, the sedative ED<sub>50</sub> values of MB-1C-OH and (-)U50,488H were 9.29 mg/kg and 3.32 mg/kg, respectively. The ratio of the sedative ED<sub>50</sub> to the antinociceptive ED<sub>50</sub> of MB-1C-OH was 23.82, which was 6.4 times higher than that of (-)U50,488H (3.73), suggesting that MB-1C-OH has a lower risk for causing sedation than (-)U50,488H.

**Table 1.** Affinity values ( $K_i$ ) for the binding to opioid receptors and EC<sub>50</sub> values as well as maximal effects in stimulating [<sup>35</sup>S]GTP $\gamma$ S binding to membranes of MB-1C-OH and (-)U50,488H in CHO cells stably expressing opioid receptors.

Compounds	Receptor binding $K_i$ (nmol/L)			[ <sup>35</sup> S]GTP $\gamma$ S binding assay	
	[ <sup>3</sup> H]diprenorphine ( $\mu$ )	[ <sup>3</sup> H]diprenorphine ( $\kappa$ )	[ <sup>3</sup> H]diprenorphine ( $\delta$ )	$E_{max}$	EC <sub>50</sub> (nmol/L)
MB-1C-OH	N.B	35.13±1.74	N.B	98.19%	16.70±0.32
(-)U50,488H <sup>a</sup>	1397.00±488.00	6.08±0.10	N.B	100%	3.75±0.31

<sup>a</sup> The data of (-)U50,488H were cited from our previous data<sup>[7]</sup>.


Membranes stably expressing opioid receptors in Chinese hamster ovary cells were incubated with varying concentrations of MB-1C-OH and (-)U50,488H in the presence of [<sup>3</sup>H]diprenorphine (1 nmol/L), or [<sup>35</sup>S]GTP $\gamma$ S (0.1 nmol/L) as described in Materials and methods. Data were derived from the curves in Figure 2 and 3. Data were presented as the mean±SEM for at least three independent experiments performed in triplicate. N.B did not bind at 100  $\mu$ mol/L.

**Table 2.** ED<sub>50</sub> values of the maximal possible effect (% MPE) produced by MB-1C-OH, morphine, and (-)U50,488H evaluated with hot plate, tail flick and acetic acid-induced writhing tests.


Compounds	Hot plate test (mg/kg, sc)	Tail flick test (mg/kg, sc)	Acetic acid-induced writhing test (mg/kg, sc)
MB-1C-OH	-	-	0.39 (0.24–0.66)
Morphine	6.95 (5.68–8.50) <sup>a</sup>	3.46 (3.21–3.72)	0.84 (0.56–1.27) <sup>a</sup>
(-)U50,488H	4.42 (2.51–7.75) <sup>a</sup>	5.02 (4.58–5.51)	0.89 (0.54–1.44) <sup>a</sup>

<sup>a</sup> The data of (-)U50,488H and morphine associated with hot plate and acetic acid-induced writhing tests were cited from our previous data<sup>[7]</sup>.

The data was obtained at 15 min after drug administration (in parentheses are 95% confidence limits).



**Figure 2.** Effects of pretreatment with Nor-BNI,  $\beta$ -FNA, and naltrindole on MB-1C-OH-induced antinociception in the acetic acid-induced writhing assay. Mice were pretreated with the  $\mu$ -opioid receptor antagonist,  $\beta$ -FNA (10 mg/kg, sc) 24 h before MB-1C-OH administration; the  $\delta$ -opioid receptor antagonist, naltrindole (3.0 mg/kg, sc) 30 min before MB-1C-OH administration; or the selective  $\kappa$ -opioid receptor antagonist, Nor-BNI (10 mg/kg, sc) 15 min before MB-1C-OH administration and then injected with MB-1C-OH (1.88 mg/kg, sc). Acetic acid solution was intraperitoneally injected 15 min after drug administration. The maximum possible effect (% MPE) was calculated as described in the Materials and methods section, and the number of times the mouse writhed is also shown. Data are presented as the mean $\pm$ SEM from 10 animals.  $^{\circ}P < 0.01$  compared with the control group.




**Figure 3.** MB-1C-OH-induced antinociceptive effects in the formalin test. The animals were pretreated with MB-1C-OH into subcutaneous. After 15 min, the animals were injected with formalin (20  $\mu$ L/paw). The amount of time the mice spent licking or flinching during first phase and second phase was recorded. All data are expressed as the mean $\pm$ SEM (at least five animals in each group).  $^{\circ}P < 0.01$  compared with the vehicle group (one-way ANOVA with Dunnett's test).

**Table 3.** ED<sub>50</sub> of each drug maximal possible effect (% MPE) measured by the formalin test in mice.

Compounds	Antinociceptive ED <sub>50</sub> (mg/kg, sc)	
	Phase I	Phase II
MB-1C-OH	-	0.87 (0.69–1.10)
Morphine	5.05 (4.33–5.90)	3.06 (2.86–3.28)
(-)-U50,488H	-	0.41 (0.32–0.61)

The data was calculated obtained at 15 min after drug administration (in parentheses are 95% confidence limits).



**Figure 4.** The depressive effect of MB-1C-OH in the forced swimming test and the sedative effect of MB-1C-OH in the rotarod test. (A) Mice were administered (sc) saline, (-)U50,488H or MB-1C-OH. After 15 min, they were put singly into transparent Plexiglas cylinders containing water at a depth of 10 cm for 6 min. The immobility time was recorded during the final 4 min. Data for each group are presented as the mean $\pm$ SEM from 5–12 animals. (B) Mice were subcutaneously administered various doses of MB-1C-OH. Then, 15 min later, mice were put singly on a rotarod for 300 s. The duration time before each mouse fell off the rotarod was recorded. Data for each group are presented as the mean $\pm$ SEM from 10 animals.

## Discussion

The present study demonstrated that MB-1C-OH had high affinity for the  $\kappa$ -opioid receptor with a  $K_i$  value of 35 nmol/L in the receptor binding assay. With the [<sup>35</sup>S]GTP $\gamma$ S binding assay, a classical functional measurement for receptor activation that can be used to determine the potencies and efficacies of opioid ligands at opioid receptors<sup>[25]</sup>, we found that MB-

**Table 4.** Depression effects in mouse forced swimming test and sedation effects in mouse rotarod test.

Compounds	Depression ED <sub>50</sub> (mg/kg, sc)	Depression ED <sub>50</sub> / antinociceptive ED <sub>50</sub>	Sedation ED <sub>50</sub> (mg/kg, sc)	Sedation ED <sub>50</sub> / antinociceptive ED <sub>50</sub>
MB-1C-OH	9.49 (8.84–10.18)	24.33	9.29 (8.75–9.87)	23.82
(-)-U50,488H	2.35 (2.21–2.50)	2.64	3.32 (2.65–3.94) <sup>a</sup>	3.73

<sup>a</sup> The sedation ED<sub>50</sub> of (-)-U50,488H were cited from our previous data<sup>[47]</sup>.

ED<sub>50</sub> values of maximal possible effect (% MPE) obtained from mice acetic acid writhing test (in parentheses are 95% confidence limits).

1C-OH stimulated [<sup>35</sup>S]GTPγS binding to membrane receptors with an  $E_{max}$  of 98% and an EC<sub>50</sub> value of 16.7 nmol/L. These results indicate that MB-1C-OH is a highly selective, potent κ-opioid receptor agonist.

It has been well demonstrated that κ-agonists produce potent antinociception in different animal pain models<sup>[9, 11–13]</sup>. Our findings were similar to these studies in that we found that MB-1C-OH produced a significant antinociceptive effect in the acetic acid-induced writhing test via κ- but not μ- or δ-opioid receptors. Furthermore, compared with morphine and (-)-U50,488H, MB-1C-OH displayed more potent antinociception in the acetic acid-induced writhing test. In the formalin test, MB-1C-OH also significantly inhibited formalin-induced flinching behavior in the second phase. These results suggested that MB-1C-OH, a full κ-receptor agonist, produced potent antinociceptive effects and were similar to our previous findings<sup>[4, 7, 16]</sup>.

More importantly, we found that MB-1C-OH had no effects on the mouse hot plate and tail flick tests. Because the hot plate and tail flick tests both measure responses to thermal pain and often reflect central drug actions mediated by supraspinal and spinal mechanisms<sup>[26, 27]</sup>, our results suggested that the antinociceptive effects of MB-1C-OH may occur via a peripheral- rather than a central-acting mechanism. We further evaluated the antinociception produced by MB-1C-OH using the acetic acid-induced writhing and formalin tests. These tests are both widely used to measure analgesic activity<sup>[28–30]</sup>. The formalin test occurs in two phases. The first phase is characterized by neurogenic pain caused by the direct chemical stimulation of nociceptors. The second phase is characterized by inflammatory pain triggered by a combination of stimuli<sup>[31, 32]</sup>. Drugs that act primarily on the central nervous system inhibit both phases equally, whereas peripherally acting drugs only inhibit the late phase<sup>[33]</sup>. MB-1C-OH produced significant inhibition of flinching behavior in the second phase but not in the first phase, suggesting that its antinociceptive effect may be related to peripheral action on inflammatory pain. The precise mechanisms of MB-1C-OH-induced antinociception are of great interest and need to be elucidated in future studies. Many recent studies have shown that κ-opioid agonists produced analgesic effect via peripheral κ-opioid receptor<sup>[9–13]</sup> without producing undesirable side effects.

Activation of the κ-receptor produces severe, undesirable central nervous system side effects, which limit the clinical utility of κ-receptor agonists for pain relief<sup>[34, 35]</sup>. In the present

study, we found that the depression and sedation ED<sub>50</sub> values of MB-1C-OH are much higher than the ED<sub>50</sub> value for antinociception. In contrast, the doses of (-)-U50,488H that produce depression and sedation were similar to those that produce antinociception. These results demonstrated that MB-1C-OH produced less risk for sedation and depression than the classical κ-agonist (-)-U50,488H.

In summary, MB-1C-OH, a novel selective κ-opioid receptor agonist, produced potent antinociceptive effects and little sedation and depression. Taken together, MB-1C-OH may be a promising analgesic for pain relief.

#### Acknowledgements

This research was supported by grants from the National Basic Research Program: a grant from the Ministry of Science and Technology of China 2009CB522005 (Jing-gen LIU) and grants from National Natural Science Foundation of China 81130087 (Jing-gen LIU) and 81171296 (Xiao-wu HU).

#### Author contribution

Jing-gen LIU, Yu-jun WANG, and Tao XI designed and supervised the research project; Yun-gen XU provided the compound; Le-sha ZHANG, Jun WANG, Yi-min TAO, Yu-hua WANG, Xue-jun XU, Jian-chun CHEN, Xiao-wu HU and Jie CHEN performed the experiments; Le-sha ZHANG, Jun WANG, and Yu-jun WANG analyzed the data and wrote the manuscript; Yu-jun WANG revised the manuscript.

#### References

- Trescot AM, Datta S, Lee M, Hansen H. Opioid pharmacology. *Pain Physician* 2008; 11: S133–53.
- Ballantyne JC. Opioid analgesia: perspectives on right use and utility. *Pain Physician* 2007; 10: 479–91.
- Clark JA, Pasternak GW. U50,488: a kappa-selective agent with poor affinity for mu1 opiate binding sites. *Neuropharmacology* 1988; 27: 331–2.
- Sun JF, Wang YH, Li FY, Lu G, Tao YM, Cheng Y, et al. Effects of ATPM-ET, a novel κ agonist with partial μ activity, on physical dependence and behavior sensitization in mice. *Acta Pharmacol Sin* 2010; 31: 1547–52.
- Wang YH, Sun JF, Tao YM, Chi ZQ, Liu JG. The role of kappa-opioid receptor activation in mediating antinociception and addiction. *Acta Pharmacol Sin* 2010; 31: 1065–70.
- Endoh T, Matsuura H, Tajima A, Izumimoto N, Tajima C, Suzuki T, et al. Potent antinociceptive effects of TRK-820, a novel kappa-opioid receptor agonist. *Life Sci* 1999; 65: 1685–94.

- 7 Tao YM, Li QL, Zhang CF, Xu XJ, Chen J, Ju YW, et al. LPK-26, a novel kappa-opioid receptor agonist with potent antinociceptive effects and low dependence potential. *Eur J Pharmacol* 2008; 584: 306–11.
- 8 Negus SS, Mello NK, Portoghese PS, Lin CE. Effects of kappa opioids on cocaine self-administration by rhesus monkeys. *J Pharmacol Exp Ther* 1997; 282: 44–55.
- 9 Riviere PJ. Peripheral kappa-opioid agonists for visceral pain. *Br J Pharmacol* 2004; 141: 1331–4.
- 10 Stein C, Lang LJ. Peripheral mechanisms of opioid analgesia. *Curr Opin Pharmacol* 2009; 9: 3–8.
- 11 Binder W, Machelska H, Mousa S, Schmitt T, Riviere PJ, Junien JL, et al. Analgesic and antiinflammatory effects of two novel kappa-opioid peptides. *Anesthesiology* 2001; 94: 1034–44.
- 12 Jolivald CG, Jiang Y, Freshwater JD, Bartoszyk GD, Calcutt NA. Dynorphin A. kappa opioid receptors and the antinociceptive efficacy of asimadoline in streptozotocin-induced diabetic rats. *Diabetologia* 2006; 49: 2775–85.
- 13 Ko MC, Tuchman JE, Johnson MD, Wiesenauer K, Woods JH. Local administration of mu or kappa opioid agonists attenuates capsaicin-induced thermal hyperalgesia via peripheral opioid receptors in rats. *Psychopharmacology (Berl)* 2000; 148: 180–5.
- 14 Liu JG, Prather PL. Chronic exposure to mu-opioid agonists produces constitutive activation of mu-opioid receptors in direct proportion to the efficacy of the agonist used for pretreatment. *Mol Pharmacol* 2001; 60: 53–62.
- 15 Wen Q, Yu G, Li YL, Yan LD, Gong ZH. Pharmacological mechanisms underlying the antinociceptive and tolerance effects of the 6,14-bridged oripavine compound O30418. *Acta Pharmacol Sin* 2011; 32: 1215–24.
- 16 Portoghese AS, Lipkowski AW, Takemori AE. Bimorphinans as highly selective, potent kappa opioid receptor antagonists. *J Med Chem* 1987; 30: 238–9.
- 17 Wang YJ, Tao YM, Li FY, Wang YH, Xu XJ, Chen J, et al. Pharmacological characterization of ATPM [(–)-3-aminothiazolo[5,4-b]-N-cyclopropylmethylmorphinan hydrochloride], a novel mixed  $\kappa$ -agonist and  $\mu$ -agonist/-antagonist that attenuates morphine antinociceptive tolerance and heroin self-administration behavior. *J Pharmacol Exp Ther* 2009; 329: 306–13.
- 18 Jiang Q, Heyman JS, Porreca F. Mu antagonist and kappa agonist properties of b-funaltrexamine (b-FNA): long lasting spinal antinociception. *NIDA Res Monogr* 1989; 95: 199–205.
- 19 Vanderah TW, Schteingart CD, Trojnar J, Junien JL, Lai J, Riviere PJ. FE200041 (D-Phe-D-Phe-D-Nle-D-Arg-NH<sub>2</sub>): A peripheral efficacious kappa opioid agonist with unprecedented selectivity. *J Pharmacol Exp Ther* 2004; 310: 326–33.
- 20 Saitoh A, Yoshikawa Y, Onodera K, Kamei J. Role of delta-opioid receptor subtypes in anxiety-related behaviors in the elevated plus-maze in rats. *Psychopharmacology (Berl)* 2005; 182: 327–34.
- 21 Hunskaar S, Hole K. The formalin test in mice: dissociation between inflammatory and non-inflammatory pain. *Pain* 1987; 30: 103–14.
- 22 David DJ, Renard CE, Jolliet P, Hascoet M, Bourin M. Antidepressant-like effects in various mice strains in the forced swimming test. *Psychopharmacology (Berl)* 2003; 166: 373–82.
- 23 Litchfield JT Jr, Wilcoxon F. A simplified method of evaluating dose-effect experiments. *J Pharmacol Exp Ther* 1949; 96: 99–113.
- 24 Shiotsuki H, Yoshimi K, Shimo Y, Funayama M, Takamatsu Y, Ikeda K, et al. A rotarod test for evaluation of motor skill learning. *J Neurosci Methods* 2010; 189: 180–5.
- 25 Zhu J, Luo LY, Li JG, Chen C, Liu-Chen LY. Activation of the cloned human kappa opioid receptor by agonists enhances [<sup>35</sup>S]GTPgammaS binding to membranes: determination of potencies and efficacies of ligands. *J Pharmacol Exp Ther* 1997; 282: 676–84.
- 26 Nemirovsky A, Chen L, Zelman V, Jurna I. The antinociceptive effect of the combination of spinal morphine with systemic morphine or buprenorphine. *Anesth Analg* 2001; 93: 197–203.
- 27 Rani S, Gupta MC. Evaluation and comparison of antinociceptive activity of aspartame with sucrose. *Pharmacol Rep* 2012; 64: 293–8.
- 28 Matsumoto A, Anan H, Maeda K. An immunohistochemical study of the behavior of cells expressing interleukin-1 alpha and interleukin-1 beta within experimentally induced periapical lesions in rats. *J Endod* 1998; 24: 811–6.
- 29 Vanderlei ES, Patoilo KK, Lima NA, Lima AP, Rodrigues JA, Silva LM, et al. Antinociceptive and anti-inflammatory activities of lectin from the marine green alga *Caulerpa cupressoides*. *Int Immunopharmacol* 2010; 10: 1113–8.
- 30 Shields SD, Cavanaugh DJ, Lee H, Anderson DJ, Basbaum AI. Pain behavior in the formalin test persists after ablation of the great majority of C-fiber nociceptors. *Pain* 2010; 151: 422–9.
- 31 Tjølsen A, Berge OG, Hunskaar S, Rosland JH, Hole K. The formalin test: an evaluation of the method. *Pain* 1992; 51: 5–17.
- 32 Puig S, Sorkin LS. Formalin-evoked activity in identified primary afferent fibers: systemic lidocaine suppresses phase-2 activity. *Pain* 1996; 64: 345–55.
- 33 Bitencourt Fda S, Figueiredo JG, Mota MR, Bezerra CC, Silvestre PP, Vale MR, Nascimento KS, et al. Antinociceptive and anti-inflammatory effects of a mucin-binding agglutinin isolated from the red marine alga *Hypnea cervicornis*. *Naunyn Schmiedeberg Arch Pharmacol* 2008; 377: 139–48.
- 34 Wadenberg ML. A review of the properties of spiradolone: a potent and selective kappa-opioid receptor agonist. *CNS Drug Rev* 2003; 9: 187–98.
- 35 Dortch-Carnes J, Potter DE. Bremazocine: a kappa-opioid agonist with potent analgesic and other pharmacologic properties. *CNS Drug Rev* 2005; 11: 195–212.