

## LETTER TO THE EDITOR

A novel fusion of *SQSTM1* and *FGFR1* in a patient with acute myelomonocytic leukemia with t(5;8)(q35;p11) translocation

*Blood Cancer Journal* (2014) 4, e265; doi:10.1038/bcj.2014.86; published online 12 December 2014

Hematological malignancies with *FGFR1* abnormality (8p11 myeloproliferative syndrome; EMS) are rare atypical stem cell disorders characterized by eosinophilia, T-cell proliferation and progression to acute myeloid leukemia.<sup>1,2</sup> In EMS, fibroblast growth factor receptor 1 (*FGFR1*) gene at 8p11 is disrupted by chromosomal translocation, resulting in the formation of chimeric products with various partner genes. To date, 13 *FGFR1* partners have been identified: *ZMYM2* (*ZNF198*) at 13q12; *CNTR* at 9q33; *FGFR1OP* at 6q27; *BCR* at 22q11; *HERVK* at 19p13.3; *FGFR1OP2* at 12p11; *TRIM24* (*TIF1*) at 7q34; *CPSF6* at 12q15; *MYO18A* at 17q23; *LRRFIP1* at 2q37; *CUX1* at 7q34; *TPR* at 1q25; and *RANBP2/NUP358* at 2q12.<sup>2–5</sup> In all reported cases in which chimeric transcripts were cloned, N-terminal portion of the predicted fusion protein is composed of a partner gene with a dimerization domain, which may induce the constitutive activation of *FGFR1* tyrosine kinase in the C-terminal portion, leading to the cellular transformation.<sup>2</sup>

Here, we report the identification of sequestosome 1 (*SQSTM1*) gene at 5q35 as a novel fusion partner of the *FGFR1* in a patient with acute myelomonocytic leukemia presenting 8p11 chromosomal abnormality.

A 76-year-old man was referred to our hospital for anemia with hemoglobin level of 5.0 g/dl. The white blood cell count was  $4.1 \times 10^9$  cells/l with 11% blasts, 1% eosinophils, 29% neutrophils and 30% monocytes. Bone marrow (BM) was hypercellular with 43% blasts and 17% monocytes. Dysplastic changes were observed in peripheral neutrophils and BM megakaryocytes. Immunophenotyping revealed that leukemic cells were positive for CD13, CD33, CD34 and HLA-DR and negative for other myeloid and lymphoid markers. He was diagnosed as acute myelomonocytic leukemia and received blood transfusion therapy for anemia, but chemotherapy was not done because of poor performance status. He died of progressive disease 15 months after the diagnosis.

Chromosome analysis of the BM cells showed the karyotype as 46,XY,t(5;8)(q35;p11) (20/20 cells) (Figure 1a). To characterize the 8p11 translocation, Southern blot analysis of DNA from the patient's peripheral leukemic cells was performed using a genomic probe containing *FGFR1* exon 9 and 10, as chromosomal breaks occur within intron 8 in most cases with EMS. Abnormal bands were observed in sample with *Bam*HI or *Hind*III digestion (Figure 1b), confirming that *FGFR1* was rearranged by the translocation.

To determine the *FGFR1* partner gene, we searched for a *FGFR1* chimeric transcript by 5'-rapid amplification of cDNA end (RACE) method using 5'-Full RACE core set (Takara Bio, Otsu, Japan). Total RNA was extracted from the leukemic cells and the first cDNA strand was synthesized with phosphorylated primer in *FGFR1* exon 12 (5'-GCCAGATACTCCATGCCTCG-3') by reverse transcription. After ligation, cDNA was amplified by PCR with *FGFR1*-10R1 (5'-ACACCACCTGCCAAAGCAGC-3') and *FGFR1*-10F1 (5'-AGGCTATCGGGCTGGACAAGG-3') primers in exon 10. The PCR product was further amplified with *FGFR1*-10R2 (5'-CAGGGGTTGCCTAAGACCAG-3') and *FGFR1*-10F2 (5'-CGTGTGACCAAGTGCTGTG-3')

primers in exon 10. An abnormal band was amplified and sequencing of the PCR product revealed that *FGFR1* was fused to a 5' foreign sequence that was identified as exon 9 of *SQSTM1* at chromosome 5q35, showing that *SQSTM1* was juxtaposed to *FGFR1* as the result of the chromosomal translocation.

To ascertain the formation of chimeric transcript, we performed reverse transcription PCR (RT-PCR) analysis, using *SQSTM1*-8F1 (5'-CCTGAAGAACGTTGGGAGAG-3') for exon 8 and *FGFR1*-10R1 primers. An amplified product was obtained from the patient's leukemic cells (Figure 1c). Sequencing analysis of the PCR product revealed that the *SQSTM1* exon 9 was fused to *FGFR1* exon 9 (Figures 1c and d).


A reciprocal *FGFR1*-*SQSTM1* fusion transcript was also detected in using forward *FGFR1*-7F1 (5'-GTAACCTATCGGACTCTCCC-3') for exon 7 and reverse *SQSTM1*-11R1 (5'-CGGGGGATGCTTTGAATAC TGG-3') for exon 11 primers (Figures 1c and d).

To identify the chromosomal breakpoints, genomic DNA was amplified by the long-and-accurate PCR method using LA Taq polymerase (Takara Bio). The following primers were used: *FGFR1*-7F1 and *FGFR1*-7F2 (5'-TCTGCATGTTGACCGTTCTG-3') for exon 7 outer and inner; *FGFR1*-10R1 and *FGFR1*-10R2 for exon 10 outer and inner; *SQSTM1*-9F1 (5'-GATATCGATGTGGAGCACGGAGGG-3'), *SQSTM1*-9F2 (5'-CTCCAGAGAGTCCAGCACAGAGG-3') for exon 9 outer and inner; and *SQSTM1*-10R1 (5'-TGTGGGTACAAGGCAGCTCC-3'), *SQSTM1*-10R2 (5'-CTTGGCCCTTCGGATTCTGGC-3') for exon 10 outer and inner, respectively. Genomic fragments corresponding to the der(5) and der(8) chromosomes were amplified by primers around the predicted breakpoints on both chromosomes (Figures 1e and f). The sequence analysis of the PCR products identified the chromosome 5 and 8 breakpoints within *SQSTM1* intron 9 and *FGFR1* intron 9, respectively (data not shown).


As sequence databases contain alternative transcripts for *SQSTM1* coding N-terminal truncated protein (NCBI Reference Sequence: NM\_001142298.1 and NM\_00114299.1), we performed RT-PCR using primers located in *SQSTM1* alternative exons 1, 2 or 4 and *FGFR1* exon 9 and detected only a chimeric transcript from *SQSTM1* exon 4 coding the full N-terminal end of *SQSTM1* (data not shown).

*SQSTM1* at chromosome 5q35 encodes a multifunctional protein that binds ubiquitin and regulates activation of the NF- $\kappa$ B signaling pathway, associating with oxidative stress and autophagy.<sup>6–9</sup> Mutations in this gene were reported to cause Paget disease of bone.<sup>10</sup> The *SQSTM1*-*FGFR1* transcript encodes a predicted chimeric protein of 718 amino acids, containing the N-terminal protein-protein interaction domain, PB1, of *SQSTM1* (ref. 11) and the tyrosine kinase domain of *FGFR1* (Figure 2). *SQSTM1*-*FGFR1* may induce the constitutive activation of tyrosine kinase by PB1-mediated dimerization, thus leading to the cellular transformation.

Recently, the generation of *SQSTM1*-*NUP214* and *SQSTM1*-*ALK* fusion genes were reported in one case with adult T-cell acute lymphoblastic leukemia presenting t(5;9)(q35;q34) translocation and two cases with anaplastic lymphoma kinase (ALK)-positive large B-cell lymphoma, respectively.<sup>12–14</sup> *SQSTM1*-*ALK* was shown to possess transforming activity in 3T3 fibroblasts, probably by the constitutive activation of ALK tyrosine kinase by PB1-mediated



**Figure 1.** Cytogenetic and molecular analysis. **(a)** Partial karyotyping of patient's bone marrow cells. The derivative chromosome 5 and 8 are shown by arrows. **(b)** Southern blot analysis, presenting rearrangement within *FGFR1* gene. Arrows indicate the abnormal bands. **(c)** RT-PCR detection of the fusion transcripts in the patient. **(d)** Partial sequence of *SQSTM1-FGFR1* and reciprocal *FGFR1-SQSTM1* fusion cDNA. The amino-acid translations spanning the fusion are shown under the sequence. **(e)** Detection of der(5) and der(8) chromosomes in the patient's leukemic cells. Amplified genomic DNA fragments from the patient's sample are indicated by arrows. **(f)** Genomic organization of *FGFR1* and *SQSTM1*, encompassing the breakpoints. Horizontal arrows indicate primers used in PCR. Lanes C and P represent normal control and the patient's sample, respectively.



**Figure 2.** Schematic representation of *FGFR1*, *SQSTM1* and the predicted *SQSTM1-FGFR1* fusion protein. Relevant protein domains are shown. The breakpoints within proteins are indicated by arrows.

dimerization.<sup>13</sup> To our knowledge, our case is the third instance of hematologic malignancy, in which *SQSTM1* was involved in chromosomal translocation. *SQSTM1* may be a recurrent target of chromosomal aberration.

Our patient, diagnosed as acute myelomonocytic leukemia, lacked typical EMS features, such as myeloproliferation,

eosinophilia or lymphadenopathy involved by T-cell proliferation. Multilineage dysplasia indicated the leukemic transformation at early myeloid precursor level, but involvement of the T-cell lineage was not determined. *SQSTM1-FGFR1* and/or reciprocal *FGFR1-SQSTM1* fusion gene may exert to promote proliferation of immature myelomonocytic cells preferentially. Further investigation is needed to clarify this point.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

Y Nakamura, Y Ito, N Wakimoto, E Kakegawa, Y Uchida and M Bessho  
 Department of Hematology, Saitama Medical University, Saitama, Japan  
 E-mail: ynakam@saitama-med.ac.jp

#### REFERENCES

- Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H *et al.* *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. IARC: Lyon, 2008.
- Jackson CC, Medeiros LJ, Miranda RN. 8p11 myeloproliferative syndrome: a review. *Hum Pathol* 2010; **41**: 461–476.
- Wasag B, Lierman E, Meeus P, Cools J, Vandenberghe P. The kinase inhibitor TKI258 is active against the novel CUX-FGFR1 fusion detected in a patient with

- T-lymphoblastic leukemia/lymphoma and t(7;8)(q22;p11). *Haematologica* 2011; **96**: 922–926.
- 4 Li F, Zhai Y-P, Tang Y-M, Wang L-P, Wan P-J. Identification of a novel partner gene, *TPR*, fused to *FGFR1* in 8p11 myeloproliferative syndrome. *Gene Chromosome Cancer* 2012; **51**: 890–897.
- 5 Gervais C, Dano L, Perrusson N, Helias C, Jeandidier E, Galois A-C *et al.* A translocation t(2;8)(q12;p11) fuses *FGFR1* to a novel partner gene, *RANBP2/NUP358*, in a myeloproliferative/myelodysplastic neoplasm. *Leukemia* 2013; **27**: 1186–1188.
- 6 Geetha T, Wooten MW. Structure and functional properties of the ubiquitin binding protein p62. *FEBS Lett* 2002; **512**: 19–24.
- 7 Seibenhener ML, Geetha T, Wooten MW. Sequestosome 1/p62—more than a scaffold. *FEBS Lett* 2007; **581**: 175–179.
- 8 Kirkin V, McEwan DG, Novak I, Dikic I. A role for ubiquitin in selective autophagy. *Mol Cell* 2009; **34**: 259–269.
- 9 Komatsu M, Kurokawa H, Waguri S, Taguchi K, Kobayashi A, Ichimura Y *et al.* The selective autophagy substrate p62 activates the stress responsive transcription factor Nrf2 through inactivation of Keap1. *Nat Cell Biol* 2010; **12**: 213–223.
- 10 Laurin N, Brown JP, Morissette J, Raymond V. Recurrent mutation of the gene encoding sequestosome 1 (SQSTM1/p62) in Paget disease of bone. *Am J Hum Genet* 2002; **70**: 1582–1588.
- 11 Wilson MI, Gill DJ, Perisic O, Quinn MT, Williams RL. PB1 domain-mediated heterodimerization in NADPH oxidase and signaling complexes of atypical protein kinase C with Par6 and p62. *Mol Cell* 2003; **12**: 39–50.
- 12 Gorello P, Starza RL, Giacomo DD, Messina M, Puzzolo MC, Crescenzi B *et al.* SQSTM1-NUP214: a new gene fusion in adult T-cell acute lymphoblastic leukemia. *Haematologica* 2010; **95**: 2161–2163.
- 13 Takeuchi K, Soda M, Togashi Y, Ota Y, Sekiguchi Y, Hatano S *et al.* Identification of a novel fusion, SQSTM1-ALK, in ALK-positive large B-cell lymphoma. *Haematologica* 2011; **96**: 464–467.
- 14 d'Amore ES, Visco C, Menin A, Famengo B, Bonvini P, Lazzari E. STAT3 pathway is activated in ALK-positive large B-cell lymphoma carrying SQSTM1-ALK rearrangement and provides a possible therapeutic target. *Am J Surg Pathol* 2013; **37**: 780–786.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>