LETTER TO THE EDITOR

Amyloid cast tubulopathy: a unique form of immunoglobulin-induced renal disease

Blood Cancer Journal (2016) 6, e474; doi:10.1038/bcj.2016.74; published online 23 September 2016

Excess production of monoclonal immunoglobulins (Ig) occurs in ~1% of all individuals beyond the age of 50.¹ The overall prevalence of kidney disease in this condition, that is, of Ig-induced kidney disease (Ig-KD) is over 20%, and its presence is associated with a wide variety of nephropathological lesions (Table 1) as well as underlying pathogenic mechanisms. The most classic Ig-KD is cast nephropathy, also called myeloma kidney. It corresponds to an Ig-induced tubular disease (Ig-TD) that generally occurs in the setting of multiple myeloma. The other types of lesions, however, are associated with a wider variety

of plasma cell dyscrasias including monoclonal gammopathy of undetermined significance. On very rare occasions, Ig-TD is limited to the proximal nephron. Under such circumstances, the lesion usually corresponds to cytosolic κ 1-restricted crystalline deposits or to λ -restricted lysosomal abnormalities.^{1,2} On the basis of a study by Larsen *et al.*,² the latter entity could be underdiagnosed as it corresponds to a more subtle renal injury in which lysosomes are typically increased in number and mottled in texture. Cytosolic fibrillary deposits in the absence of crystal formation have also been identified in a few cases.^{3–6} In the current report, we describe an even rarer, if not unique, form of Ig-TD. This entity was identified in a 54-year-old diabetic man admitted for renal failure (creatinine 5.49 mg/dl) and found to have an IgG λ gammopathy due to multiple myeloma in the absence of Fanconi syndrome. As seen in Figure 1, the lesion consisted of numerous spherical deposits that were confined to the cytoplasm of the proximal nephron (upward arrows) and to the lumen of many nephron segments where they formed aggregates (downward arrows). The deposits also shared a unique combination of features in that they were pale by hematoxylin and eosin, and periodic acid-Schiff (PAS) (a and b), positive by trichrome and Congo red (c and d), fibrillary (e and f) and λ -restricted (not shown). From these findings, we concluded that the pathological picture observed belonged to the highly uncommon category of isolated fibrillary Ig-TD of which a subset is termed amyloid tubulopathy (AT) in the presence of Congo red-positive deposits. Such an entity has only been reported twice thus far.^{5,6} In the current case, surprisingly, it was also accompanied by amyloid casts (AC), a type of lesion that has already been reported^{6–10} but that was described previously as spiculated structures within the periphery of atypical casts—instead of intraluminal nodular structures—and that was generally identified in the absence of AT. The other subset of isolated fibrillary Ig-TD is Congo red-negative, is also quite uncommon^{3,4} and could correspond to a variant of fibrillary or immunotactoid glomerulopathy. At times, plasma cell dyscrasias can induce two types of renal lesions in the same individual. In our patient, AC and AT could have still resulted from a single process in which Ig β -fibrillary structures formed in the urinary space were endocytosed by proximal tubular cells. This hypothesis would be consistent with the absence of amyloid deposition in other renal structures and the limited ability of β -fibrils to permeate the glomerular filtration barrier. It would also suggest that AT, AC and AC-associated AT all correspond to distinct entities. On the basis of our observations, it appears that isolated Ig-TD could now come into as much as nine different entities, expanding further the list of lesions that can develop in the face of excess monoclonal Ig production. In Table 1, we have regrouped these different entities based on the type of Ig deposits formed, that is, fibrillary, crystalloid or plain Ig deposits, and are using the term amyloid cast tubulopathy (ACT) to designate the new form of Ig-TD identified through this case.

Table 1. Types of Ig-KD

<i>Ig-GD (with or without tubulointerstitial deposition)</i>	
Fibrillary Ig	
Amyloidotic (L, HC, LCHC)	
Predominantly vascular	
Predominantly glomerular	
Non-Amyloidotic	
Immunotactoid (LCHC)	
With glomerulonephritis (LCHC)	
With cryoglobulinemia type I (LCHC)	
Fibrillary <i>per se</i> (LCHC)	
Crystalloid Ig	
With cryoglobulinemia type I	
Crystal-storing histiocytosis	
Plain (Ig or C)	
With glomerulosclerosis (LC, HC, LCHC)	
With glomerulonephritis	
Ig glomerulonephritis (LCHC)	
C3 glomerulonephritis (C3)	
C4 glomerulonephritis (C4)	
<i>Isolated Ig-TD</i>	
Fibrillary Ig	
Amyloidotic	
Amyloid tubulopathy	
Amyloid cast	
Amyloid cast tubulopathy	
Interstitial	
Non-Amyloidotic: immunotactoid or fibrillary?	
Crystalloid Ig	
Plain Ig	
Cast nephropathy	
Lysosomal	
Interstitial fibrosis with or without interstitial nephritis	

Abbreviations: HC, heavy chain; LC, light chain. Deposits can also be found in the tubulointerstitial compartment. In crystalline Ig-TD, deposits occur in the lumen, cytosol and lysosomes of proximal tubular cells, are negative by PAS and birefringent. In non-amyloidotic Ig-TD, deposits occur in the cytosol, are spherical, contain tightly joined fibrils of ~8 nm diameter and lead to cellular vacuolization. Crystalline and lysosomal Ig-TD are often associated with Fanconi syndrome.

Figure 1. Histological and ultrastructural characteristics of renal lesions in the current case. (a) Hematoxylin and eosin. (b) Periodic acid-Schiff. (c) Trichrome. (d) Congo red. Isolated cytosolic deposits are shown by arrows pointing downward and through insets in which they are magnified further. Clumped intraluminal deposits are shown by arrows pointing upward. A dense plasma cell infiltrate is also seen in a. (e and f) Electron microscopy. Micrographs were taken at $\times 5000$ - and $\times 40\,000$ -magnification, respectively). The box in e represents the field that was magnified in f. Amyloid fibrils measured ~ 8 nm in diameter.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

The patient has given consent for the case to be published.

AUTHOR CONTRIBUTIONS

IAI wrote the first draft and reviewed the patient's chart. APG wrote an advanced draft and reviewed the literature. EL interpreted the renal biopsy and took the micrographs. PI wrote the final draft, reviewed the literature and assembled Figure 1.

I-A Iliuta¹, AP Garneau¹, E Latulippe² and P Isenring¹
¹Department of Nephrology, Québec, QC, Canada and
²Department of Medicine, Pathology Group, L'Hôtel-Dieu
 du CHU de Québec, Laval University, Québec,
 QC, Canada
 E-mail: paul.isenring@crhdq.ulaval.ca

REFERENCES

- Sanders PW. Mechanisms of light chain injury along the tubular nephron. *J Am Soc Nephrol* 2012; **23**: 1777–1781.
- Larsen CP, Bell JM, Harris AA, Messias NC, Wang YH, Walker PD. The morphologic spectrum and clinical significance of light chain proximal tubulopathy with and without crystal formation. *Mod Pathol* 2011; **24**: 1462–1469.
- Corbett RW, Cook HT, Duncan N, Moss J. Fibrillary inclusions in light chain proximal tubulopathy associated with myeloma. *Clin Kidney J* 2012; **5**: 75–76.
- Yao Y, Wang SX, Zhang YK, Wang Y, Liu L, Liu G. Acquired Fanconi syndrome with proximal tubular cytoplasmic fibrillary inclusions of λ light chain restriction. *Intern Med* 2014; **53**: 121–124.
- Larsen CP, Borrelli SB, Walker PD. Amyloid proximal tubulopathy: a novel form of light chain proximal tubulopathy. *Clin Kidney J* 2012; **5**: 130–132.
- El-Zoghby Z, Lager D, Gregoire J, Lewin M, Sethi S. Intratubular amyloidosis. *Kidney Int* 2007; **72**: 1282–1288.
- Nasr SH, Alobeid BB, Otrakji JA, Markowitz GS. Myeloma cast nephropathy, direct renal infiltration by myeloma, and renal extramedullary hematopoiesis. *Kidney Int* 2008; **73**: 517–518.
- Melato M, Falconieri G, Pascali E, Pezzoli A. Amyloid casts within renal tubules: a singular finding in AC myelomatosis. *Virchows Arch A Pathol Anat Histol* 1980; **387**: 133–145.
- Friman C, Tornroth T, Wegelius O. IgD myeloma associated with multiple extra-medullary amyloid-containing tumours and amyloid casts in the renal tubules. *Ann Clin Res* 1970; **2**: 161–166.
- Sethi S, Hanna MH, Fervenza FC. Unusual casts in a case of multiple myeloma. *Am J of Kidney Dis* 2009; **54**: 970–974.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016