LETTER TO THE EDITOR

A phase 1/2 study of chemosensitization with plerixafor plus G-CSF in relapsed or refractory acute myeloid leukemia

Blood Cancer Journal (2017) 7, e542; doi:10.1038/bcj.2017.21; published online 10 March 2017

The interaction of acute myeloid leukemia (AML) blasts with the bone marrow (BM) microenvironment provides potent protection against both spontaneous apoptosis and chemotherapy.¹ Similar to normal hematopoietic stem cells (HSCs), AML blasts express many of the same adhesion molecules such as CXCR4, VLA-4, VLA-5 and CD44, which allow them to interact with the marrow microenvironment. We and others have demonstrated in murine models that CXCR4 inhibitors can mobilize AML cells from the BM into the peripheral blood and enhance the anti-leukemic effects of chemotherapy.^{2,3} We have also demonstrated in a previous phase 1/2 study that plerixafor, a small molecule inhibitor of CXCR4, can be safely combined with chemotherapy in patients with relapsed or refractory AML with encouraging response rates.⁴

Mobilization of HSCs by granulocyte-colony stimulating factor (G-CSF) occurs through downregulation of mRNA and protein levels of CXCL12, the ligand for CXCR4.⁵ For mobilization of autologous HSCs, G-CSF acts synergistically when combined with plerixafor and mobilizes higher numbers of CD34⁺ cells compared with either agent alone.^{6,7} In AML, 'priming' with G-CSF concurrent with chemotherapy may result in superior outcomes for patients receiving induction therapy for AML.⁸ We hypothesized that disruption of the interaction between leukemic blasts with the marrow microenvironment using G-CSF in combination with plerixafor would effectively mobilize and sensitize AML blasts to chemotherapy.

In this phase 1/2 study (ClinicalTrials.gov NCT00906945), we evaluated the combination of G-CSF and plerixafor in conjunction with mitoxantrone, etoposide and cytarabine (MEC). Eligible participants were adults, age 18–70 years old, with relapsed or refractory AML. Subjects with a peripheral blood blast count of $\geq 20 \times 10^3/\text{mm}^3$, acute promyelocytic leukemia, active central nervous system leukemia or who had been previously treated with MEC chemotherapy were excluded from the study. The primary endpoint in phase I was to determine the maximum tolerated dose and in phase II was to determine the complete response rate (CR+CRi) of plerixafor plus G-CSF in combination with MEC in patients with relapsed or refractory AML. The phase I was performed using a standard 3+3 design escalating to a maximum plerixafor dose of 0.75 mg/kg/day. For the phase II, a bivariate design was used in two stages to allow an interim analysis of toxicity and response rates.⁹ Treatment consisted of G-CSF 10 mcg/kg by subcutaneous injection daily on days 1–8. Plerixafor was administered intravenously (IV) on days 3–8. Chemotherapy consisting of mitoxantrone 8 mg/m²/day IV, etoposide 100 mg/m²/day IV and cytarabine 1000 mg/m²/day (MEC) was administered on days 4–8, ~4 h after administration of plerixafor.

Thirty-five patients with a median age of 56 years (range 29–70) were enrolled and treated on this study. The majority received treatment for first relapse ($n=21$, 60%) with 10 patients (29%) having had a prior allogeneic hematopoietic cell transplantation (Table 1). In the phase I, plerixafor was successfully escalated from 0.24 to 0.75 mg/kg/day in five dose cohorts. The plerixafor

0.75 mg/kg/day dose was brought forward in the phase II expansion with a total of 20 patients treated in the first stage of the phase II. After an interim analysis, we observed that 6 out of 20 patients treated at the phase II dose achieved a CR/CRi (30%), which is less than the 7 out of 20 responses (35%) indicated in the study design for proceeding with the second stage of the phase II. As a result, the study was terminated for futility.

With a median follow-up of 34.6 months, the median overall survival for all subjects was 7.6 months with a 1-year overall survival of 37% (95% confidence interval: 21.2–53). The median time to neutrophil recovery ($\text{ANC} \geq 1000/\text{mm}^3$) was 40 days (range 23–62) from the start of treatment (36 days from MEC). The median time to platelet recovery ($\text{platelets} \geq 100 \text{ K}/\text{mm}^3$) was 32 days (range 30–62). Adverse events were typical of those observed for patients with relapsed or refractory AML with the most common non-hematologic adverse events of nausea (69%), vomiting (37%), febrile neutropenia (57%), headache (40%), fatigue (34%), fever (29%) and electrolyte abnormalities (hypocalcemia 37%, hypokalemia 31%) (Supplementary Table S1).

Table 1. Baseline patient characteristics ($n=35$)

<i>Patient characteristic</i>	
Age, median (range)	56 (29–70)
<i>Gender</i>	
Male (%)	18 (51)
Female (%)	17 (49)
<i>ECOG performance status</i>	
0 (%)	15 (43)
1 (%)	15 (43)
2 (%)	4 (11)
Missing (%)	1 (3)
<i>Onset of AML</i>	
De novo (%)	23 (66)
Therapy-related (%)	12 (34)
Prior MDS/MPN (%)	4 (11)
<i>Indication for treatment</i>	
Primary refractory (%)	10 (29)
First relapse (%)	21 (60)
Second relapse (%)	4 (11)
Prior unsuccessful salvage chemotherapy (%)	7 (20)
Prior allogeneic HCT (%)	10 (29)
<i>Cytogenetics</i>	
Favorable (%)	3 (9)
Intermediate (%)	19 (54)
Poor (%)	13 (37)
WBC k/cumm (range)	2.5 (0.2–14.7)
BM blast % (range)	33 (0–96)

Abbreviations: AML, acute myeloid leukemia; BM, bone marrow; ECOG, Eastern Cooperative Oncology Group; HCT, hematopoietic cell transplantation; MDS, myelodysplastic syndrome; MPN, myeloproliferative neoplasm; WBC, white blood cells.

To determine the effect of G-CSF and plerixafor on leukemic cell mobilization, peripheral blood samples were collected at baseline, after G-CSF only on day 3 (pre-P), and at 2, 4, 6 and 24 h after IV plerixafor administration (Figure 1a). We were unable to detect a dose–response relationship between plerixafor and leukemic mobilization because of substantial interpatient variability in mobilization and the limited numbers of patients analyzed (Figures 1b and c). Peak mobilization of both total leukocytes and AML blasts occurred ~4–6 h after the administration of plerixafor on day 3 of G-CSF treatment and cell counts remained elevated at 24 h after administration (Figures 1b and c). In 31 evaluable patients, total leukocytes and blast cell counts increased a median of 2.4- and 3.5-fold, respectively, after 2 days of G-CSF (Figure 1d). Circulating levels of both leukocytes and blasts were further increased an additional 1.8- or 2.8-fold, respectively, at 6 h after plerixafor administration on day 3 (Figure 1d). The magnitude of AML blast cell mobilization was significantly greater than the total leukocyte mobilization at all treatment time points. Therefore similar to normal stem cell mobilization, plerixafor augments the mobilization of AML blasts by G-CSF.

We measured the expression of CXCR4 on AML blasts in response to administration of G-CSF and plerixafor using two different mAb clones. Plerixafor inhibits the binding of clone 12G5 to CXCR4. In contrast, the 1D9 mAb binds to the N-terminus of CXCR4 and is not affected by plerixafor. Similar to our previous trial,⁴ we observed a decrease in 12G5 binding on AML blasts from pretreatment to 6 h and an increase from 6 to 24 h toward baseline, indicating transient CXCR4 blockade by plerixafor *in vivo* (Figures 1e and f). In contrast, when CXCR4 was measured using 1D9, we found that there was an almost immediate (within 2 h) and dramatic increase of surface CXCR4 expression after plerixafor administration. Others and we have previously demonstrated that this upregulation of surface CXCR4 is primarily due to the plerixafor-mediated inhibition of CXCR4 internalization by its ligand CXCL12.^{4,10}

In addition to showing CXCR4 blockade by plerixafor *in vivo*, we also observed downregulation of the G-CSF receptor (CD114) and integrin alpha 6 (CD49f) on AML blasts by G-CSF (Figure 1f). In contrast, we observed no significant cell surface modulation of CD49d, CD62L, CD117 or CD135 on AML blasts following treatment with G-CSF and plerixafor (data not shown). However, we observed a slight but highly variable increase in the expression of the cell proliferation marker Ki67 in AML blasts following 2 days of G-CSF treatment (Figure 1f).

In this study, we sought to maximize blockage of the CXCL12/CXCR4 axis through (i) addition of G-CSF, which downregulates CXCL12 expression and acts synergistically with plerixafor in normal stem cell mobilization; (ii) intravenous instead of subcutaneous dosing of plerixafor to improve the kinetics of administration; and (iii) dose escalation of plerixafor to maximize CXCR4 blockade. Although we demonstrated the safety and feasibility of combining G-CSF and plerixafor with chemotherapy, combination therapy did not improve remission rates compared with historical controls.

We believe that primarily two factors contributed to the failure of G-CSF to enhance the efficacy of plerixafor in combination with MEC salvage chemotherapy in our clinical trial. First, a high proportion of patients with unfavorable prognosis were recruited to the study in comparison with our previous study, which was conducted at a single institution (Supplementary Table S2). Second, stimulation of cells with G-CSF has been shown to activate multiple signal transduction pathways that regulate the proliferation, differentiation and survival of myeloid cells. We believe that prosurvival signals mediated by G-CSF counteract both (i) the 'priming' proliferative effect of G-CSF that increases the susceptibility of leukemic cells to cell cycle-specific chemotherapeutic agents and (ii) the pro-apoptotic effects of CXCR4 inhibition by plerixafor.

In our previous study of plerixafor with MEC, blast cell mobilization was not associated with remission.⁴ We hypothesize that inhibition of CXCR4-mediated prosurvival signaling is more important than the physical detachment and mobilization of AML blasts in enhancing the efficacy of MEC chemotherapy.

Plerixafor has a relatively short half-life of 4–5 h and is known to be a weak partial agonist of CXCR4.¹¹ As high CXCR4 expression is a marker of poor prognosis in AML, plerixafor-mediated upregulation of CXCR4 may undermine the intended anti-apoptotic effect of CXCR4 blockade by also enhancing the re-homing of AML blasts to the BM, thus reducing the efficacy of CXCR4 inhibitor-based chemosensitization. A number of other CXCR4 inhibitors have been developed for clinical use including other small molecule inhibitors, antibodies and peptidomimetics. Compared with plerixafor, many of these agents provide more potent and sustained inhibition of CXCR4.^{12–14} These newer inhibitors also can directly induce apoptosis of AML and other tumor cells line *in vitro* which supports the assertion that antileukemic activity of CXCR4 inhibitors may be in part independent of the mobilization.¹⁵ We are currently exploring the use of these alternative CXCR4 inhibitors, as well as inhibitors of other pathways, which mediate tumor–stromal interactions in both AML and other hematologic malignancies.

CONFLICT OF INTEREST

MPR, KM, WE, WS, RV, KSG, AFC, CNA, PW and JFD have declared that no conflict of interest exists. GLU, MPR and RMS have received honoraria and MK has received research funding from Sanofi/Genzyme.

ACKNOWLEDGEMENTS

We thank the Alvin J Siteman Cancer Center at Washington University School of Medicine and Barnes-Jewish Hospital in St Louis, Missouri, for the use of their core facilities. The Siteman Cancer Center is supported in part by Cancer Center Support Grant P30 CA91842 from the National Cancer Institute at the National Institutes of Health. Research reported in this publication was supported by from research funding Genzyme/Sanofi, National Institutes of Health grants K23 CA140707 (GLU), R01 CA152329 (JFD), R21 CA141523 (JFD) and R21 CA132269 (JFD). The content is solely the responsibility of the authors and does not necessarily represent the official view of the NIH.

AUTHOR CONTRIBUTIONS

GLU, MPR and JFD designed and performed the research, analyzed and interpreted data, and drafted the manuscript. WS performed statistical analyses. RMS, MYK, WE, RV, KSG, AFC, CNA and PW contributed patients to this study. TRF, TR and KM collected and analyzed data.

GL Uy^{1,5}, MP Rettig^{1,5}, RM Stone², MY Konopleva³, M Andreeff³, K McFarland¹, W Shannon⁴, TR Fletcher¹, T Reineck¹, W Eades¹, K Stockerl-Goldstein¹, CN Abboud¹, MA Jacoby¹, P Westervelt¹ and JF DiPersio¹

¹Division of Oncology, Department of Medicine, Washington University School of Medicine, St Louis, MO, USA;

²Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA;

³Department of Leukemia, The University of Texas MD Anderson Cancer Center, Houston, TX, USA and

⁴Division of General Medical Sciences, Department of Medicine, Washington University School of Medicine, St Louis, MO, USA

E-mail: jdpersi@wustl.edu

⁵These authors contributed equally to this work.

REFERENCES

1. Tabe Y, Konopleva M. Role of microenvironment in resistance to therapy in AML. *Curr Hematol Malig Rep* 2015; **10**: 96–103.

- 2 Nervi B, Ramirez P, Rettig MP, Uy GL, Holt MS, Ritchey JK *et al.* Chemosensitization of acute myeloid leukemia (AML) following mobilization by the CXCR4 antagonist AMD3100. *Blood* 2009; **113**: 6206–6214.
- 3 Zeng Z, Shi YX, Samudio IJ, Wang RY, Ling X, Frolova O *et al.* Targeting the leukemia microenvironment by CXCR4 inhibition overcomes resistance to kinase inhibitors and chemotherapy in AML. *Blood* 2009; **113**: 6215–6224.
- 4 Uy GL, Rettig MP, Motabi IH, McFarland K, Trinkaus KM, Hladnik LM *et al.* A phase 1/2 study of chemosensitization with the CXCR4 antagonist plerixafor in relapsed or refractory acute myeloid leukemia. *Blood* 2012; **119**: 3917–3924.
- 5 Greenbaum AM, Link DC. Mechanisms of G-CSF-mediated hematopoietic stem and progenitor mobilization. *Leukemia* 2011; **25**: 211–217.
- 6 DiPersio JF, Micallef IN, Stiff PJ, Bolwell BJ, Maziarz RT, Jacobsen E *et al.* Phase III prospective randomized double-blind placebo-controlled trial of plerixafor plus granulocyte colony-stimulating factor compared with placebo plus granulocyte colony-stimulating factor for autologous stem-cell mobilization and transplantation for patients with non-Hodgkin's lymphoma. *J Clin Oncol* 2009; **27**: 4767–4773.
- 7 DiPersio JF, Stadtmauer EA, Nademanee A, Micallef IN, Stiff PJ, Kaufman JL *et al.* Plerixafor and G-CSF versus placebo and G-CSF to mobilize hematopoietic stem cells for autologous stem cell transplantation in patients with multiple myeloma. *Blood* 2009; **113**: 5720–5726.
- 8 Lowenberg B, van Putten W, Theobald M, Gmur J, Verdonck L, Sonneveld P *et al.* Effect of priming with granulocyte colony-stimulating factor on the outcome of chemotherapy for acute myeloid leukemia. *N Engl J Med* 2003; **349**: 743–752.
- 9 Bryant J, Day R. Incorporating toxicity considerations into the design of two-stage phase II clinical trials. *Biometrics* 1995; **51**: 1372–1383.
- 10 Kawaguchi A, Orba Y, Kimura T, Iha H, Ogata M, Tsuji T *et al.* Inhibition of the SDF-1 α -CXCR4 axis by the CXCR4 antagonist AMD3100 suppresses the migration of cultured cells from ATL patients and murine lymphoblastoid cells from HTLV-I Tax transgenic mice. *Blood* 2009; **114**: 2961–2968.
- 11 Zhang WB, Navenot JM, Haribabu B, Tamamura H, Hiramatsu K, Omagari A *et al.* A point mutation that confers constitutive activity to CXCR4 reveals that T140 is an inverse agonist and that AMD3100 and ALX40-4C are weak partial agonists. *J Biol Chem* 2002; **277**: 24515–24521.
- 12 Borthakur G, Ofran Y, Nagler A, Rowe JM, Foran JM, Uy GL *et al.* The peptidic CXCR4 antagonist, BL-8040, significantly reduces bone marrow immature leukemia progenitors by inducing differentiation, apoptosis and mobilization: results of the dose escalation clinical trial in acute myeloid leukemia. *Blood* 2015; **126**: 2546–2546.
- 13 Peng SB, Zhang X, Paul D, Kays LM, Ye M, Vaillancourt P *et al.* Inhibition of CXCR4 by LY2624587, a fully humanized anti-CXCR4 antibody induces apoptosis of hematologic malignancies. *PLoS One* 2016; **11**: e0150585.
- 14 Becker PS, Foran JM, Altman JK, Yacoub A, Castro JE, Sabbatini P *et al.* Targeting the CXCR4 pathway: safety, tolerability and clinical activity of ulocuplumab (BMS-936564), an anti-CXCR4 antibody, in relapsed/refractory acute myeloid leukemia. *Blood* 2014; **124**: 386–386.
- 15 Kashyap MK, Kumar D, Jones H, Amaya-Chanaga CI, Choi MY, Melo-Cardenas J *et al.* Ulocuplumab (BMS-936564 / MDX1338): a fully human anti-CXCR4 antibody induces cell death in chronic lymphocytic leukemia mediated through a reactive oxygen species-dependent pathway. *Oncotarget* 2016; **7**: 2809–2822.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017

Supplementary Information accompanies this paper on Blood Cancer Journal website (<http://www.nature.com/bcj>)