

Potential tumorigenic programs associated with *TP53* mutation status reveal role of VEGF pathway

H Joshi^{*1}, G Bhanot^{2,3}, A-L Børresen-Dale^{4,5} and V Kristensen^{1,4,5}

¹Medical Division (EpiGen), Akershus University Hospital and University of Oslo, Lorenskog 1478, Norway; ²Department of Molecular Biology and Biochemistry, Rutgers University, Piscataway, NJ 08854, USA; ³Department of Physics and BioMaPS Institute, Rutgers University, Piscataway, NJ 08854, USA; ⁴Institute of Clinical Medicine, University of Oslo, Oslo 0450, Norway; ⁵Department of Genetics, Institute for Cancer Research, Oslo University Hospital Radiumhospitalet, Oslo 0310, Norway

BACKGROUND: Targeting differentially activated or perturbed tumour pathways is the key idea in individualised cancer therapy, which is emerging as an important option in treating cancers with poor prognostic profiles. *TP53* mutation status is known as a core determinant of survival in breast cancer. The pathways disrupted in association with *TP53* mutation status in tumours are not well characterised. **METHOD:** In this study, we stratify breast cancers based on their *TP53* mutation status and identify the set of dysregulated tumorigenic pathways and corresponding candidate driver genes using breast cancer gene expression profiles. Expressions of these genes were evaluated for their effect on patient survival first in univariate models, followed by multivariate models with *TP53* status as a covariate. **RESULTS:** The most strongly differentially enriched pathways between breast cancers stratified by *TP53* mutation status include in addition to *TP53* signalling, several known cancer pathways involved in renal, prostate, pancreatic, colorectal, lung and other cancers, and signalling pathways such as calcium signalling, MAPK, ERBB and vascular endothelial growth factor (VEGF) signalling pathways. We found that mutant *TP53* in conjunction with active estrogen receptor (ER) signalling significantly influence survival. We also found that upregulation of *VEGFA* mRNA levels in association with active ER signalling is a significant marker for poor survival, even in the presence of wild-type *TP53*.

CONCLUSION: Mutation status of *TP53* in breast cancer involves wide ranging derangement of several pathways. Among the candidate genes of the significantly deranged pathways, we identified *VEGFA* expression as an important marker of survival even when controlled by *TP53* mutation status. Interestingly, independent of the *TP53* mutation status, the survival effect of *VEGFA* was found significant in patients with active ER signalling (ER/PgR +), but not in those with ER/PgR – status. Therefore, we propose more studies to focus on the role of complex interplay between *TP53*, ER and VEGF signalling from therapeutic and prognostic context in breast cancer.

British Journal of Cancer (2012) 107, 1722–1728. doi:10.1038/bjc.2012.461 www.bjancer.com

Published online 18 October 2012

© 2012 Cancer Research UK

Keywords: breast cancer; *TP53* mutation status; estrogen receptor signalling; vascular endothelial growth factor signalling; dysregulated pathways; survival

The fact that nearly 30% of early-diagnosed breast cancer cases might eventually develop recurrent or metastatic disease (O'Shaughnessy, 2005) – underscores the priority to explore the mechanisms of advanced disease. The *TP53* protein is an important clinical biomarker of breast cancer because of its association with tumour progression (Norberg *et al*, 2001), metastatic potential (D'Assoro *et al*, 2010), early relapse (Aas *et al*, 1996), response to chemotherapy (Aas *et al*, 1996; Kandioler-Eckersberger *et al*, 2000; Bertheau *et al*, 2007), and ultimately, to prognosis and survival (Børresen *et al*, 1995; Berns *et al*, 2000; Olivier *et al*, 2006). It is also of relevance to molecular subtypes of breast cancer (Miller *et al*, 2005; Langerød *et al*, 2007). Whereas ~70% of breast cancers with wild-type *TP53* are mostly of the Luminal A subtype, mutant *TP53* is common in the remaining 30%, which have a poorer prognosis and are classified as triple negative or luminal B. The focus of this work is to identify

diagnostic, prognostic and therapeutic biomarkers associated with pathways perturbed by *TP53* mutations and understand their relationship to patient survival in breast cancer, under current therapeutic protocols.

TP53 is a key regulator of programmed cell death, cell cycle, DNA repair and genomic stability. In response to stimulus-specific post-transcriptional modification, *TP53* regulates genes, which activate specific cellular programs. The *TP53* protein has three major functional domains: a transactivation domain at its N-terminal, a central DNA-binding domain (which includes mutation hotspots) and tetramerization and regulatory domains at the C-terminal. The location and type of *TP53* mutation affect the ability of *TP53* to regulate its target genes, leading to aberrant functions (Blandino *et al*, 1999) with clinical implications (Kim and Deppert, 2006). Characterisation of the differential activation of key pathways and candidate genes according to the *TP53* mutation status may therefore identify mechanisms correlated with *TP53* mutation status in breast cancer.

In this study, we stratify breast cancers based on their *TP53* mutation status and identify the set of dysregulated tumorigenic pathways and their candidate driver genes by using gene

*Correspondence: Dr H Joshi; E-mail: Dr.Himanshu.Joshi@gmail.com

Received 29 May 2012; revised 21 August 2012; accepted 18 September 2012; published online 18 October 2012

expression data sets obtained from tumours. The goal is to infer the class-specific candidate gene signature by identifying weak to moderate, but coherent gene expressions that significantly influence tumorigenic pathways and survival.

RESULTS

We first categorised breast cancer samples by their corresponding TP53 mutation status, as described in Supplementary Table 1 and performed analysis as shown in the flow-chart (Supplementary Figure 1).

Candidate driver pathways differentially perturbed by TP53 mutations

Enrichment analysis of pathways between mutation status classes was performed using globaltest (Goeman *et al*, 2011) and SAM-GS


(Dinu *et al*, 2007) on the primary and combined validation data set. Globaltest, although being sensitive to genes with smaller regression coefficients, its results might be influenced by the standardisation and normalisation procedures. SAM-GS on the other hand is shown to have relatively higher power in the lower alpha-level region, thus can better focus on pathways of greatest interest (Liu *et al*, 2007). Therefore, we use a combination of the two approaches here. The list of differentially enriched KEGG (Kanehisa and Goto, 2000) pathways identified by each of the methods on each of the data set is shown together in Supplementary Table 2. A set of 40 pathways inferred as commonly significant by both the methods in both data sets (Table 1) – are graphically presented as an enrichment map color-coded according to globaltest FDR corrected *P*-values (Supplementary Figure 2). The most dysregulated pathways included a group of key signalling pathways – such as p53 signalling, calcium signalling, MAPK, ErbB, vascular endothelial growth factor (VEGF) signalling and various cancer pathways.

Table 1 Consensus list of differentially enriched pathways between two TP53 mutation status classes (wild-type TP53 profiles compared with the mutant TP53 profiles), based on pathway analysis performed by using two approaches – globaltest and SAM-GS on primary (*n* = 111 samples) and validation data sets (a combined cross-platform data set with *n* = 327)

KEGGID	KEGG pathway name	Primary data set		Validation data set	
		Asymptotic global test BH corrected <i>P</i> -value	SAM-GS FDR adj <i>P</i> -value	Asymptotic global test BH corrected <i>P</i> -value	SAM-GS FDR adj <i>P</i> -value
hsa:00230	Purine metabolism	1.8E – 09	< 10e – 6	2.37E – 36	< 10e – 6
hsa:04115	p53-signalling pathway	1.8E – 09	< 10e – 6	2.43E – 34	< 10e – 6
hsa:05211	Renal cell carcinoma	3.72E – 09	< 10e – 6	1.32E – 17	< 10e – 6
hsa:05200	Pathways in cancer	1.1E – 08	< 10e – 6	6.86E – 29	< 10e – 6
hsa:05215	Prostate cancer	1.1E – 08	< 10e – 6	1.32E – 29	< 10e – 6
hsa:04020	Calcium-signalling pathway	4.12E – 08	< 10e – 6	4.81E – 27	< 10e – 6
hsa:00260	Glycine, serine and threonine metabolism	4.73E – 08	< 10e – 6	1.44E – 25	< 10e – 6
hsa:05212	Pancreatic cancer	5.65E – 08	< 10e – 6	1.15E – 39	< 10e – 6
hsa:04340	Hedgehog-signalling pathway	6.02E – 08	< 10e – 6	5.76E – 21	< 10e – 6
hsa:05222	Small-cell lung cancer	7.93E – 08	< 10e – 6	6.75E – 40	< 10e – 6
hsa:04120	Ubiquitin-mediated proteolysis	0.00000012	< 10e – 6	3.26E – 40	< 10e – 6
hsa:04910	Insulin signalling pathway	0.00000012	< 10e – 6	5.83E – 27	< 10e – 6
hsa:00051	Fructose and mannose metabolism	1.28E – 07	< 10e – 6	2.51E – 30	< 10e – 6
hsa:05218	Melanoma	0.00000014	< 10e – 6	1.7E – 17	< 10e – 6
hsa:04150	mTOR-signalling pathway	1.68E – 07	< 10e – 6	9.66E – 26	< 10e – 6
hsa:00380	Tryptophan metabolism	1.96E – 07	< 10e – 6	1.48E – 08	< 10e – 6
hsa:04144	Endocytosis	2.39E – 07	< 10e – 6	4.96E – 24	< 10e – 6
hsa:00330	Arginine and proline metabolism	0.00000025	< 10e – 6	1.29E – 18	< 10e – 6
hsa:05214	Glioma	0.00000025	< 10e – 6	1.47E – 14	< 10e – 6
hsa:04010	MAPK-signalling pathway	0.00000031	< 10e – 6	2.44E – 34	< 10e – 6
hsa:04012	ErbB-signalling pathway	3.65E – 07	< 10e – 6	2.68E – 17	< 10e – 6
hsa:04520	Adherens junction	4.03E – 07	< 10e – 6	9.78E – 13	< 10e – 6
hsa:05217	Basal cell carcinoma	0.00000048	< 10e – 6	6.47E – 11	< 10e – 6
hsa:00600	Sphingolipid metabolism	4.94E – 07	< 10e – 6	4.67E – 14	< 10e – 6
hsa:05120	Epithelial cell signalling in <i>Helicobacter pylori</i> infection	5.79E – 07	< 10e – 6	1.45E – 11	< 10e – 6
hsa:04722	Neurotrophin-signalling pathway	6.72E – 07	< 10e – 6	1.09E – 21	< 10e – 6
hsa:04912	GnRH-signalling pathway	8.22E – 07	< 10e – 6	6E – 18	< 10e – 6
hsa:05219	Bladder cancer	8.23E – 07	< 10e – 6	1.61E – 17	< 10e – 6
hsa:05210	Colorectal cancer	0.00000116	< 10e – 6	3.9E – 11	< 10e – 6
hsa:04070	Phosphatidylinositol-signalling system	0.00000117	< 10e – 6	2.16E – 12	< 10e – 6
hsa:04110	Cell cycle	0.00000125	< 10e – 6	3.7E – 27	< 10e – 6
hsa:04370	VEGF-signalling pathway	0.00000153	< 10e – 6	1.01E – 07	< 10e – 6
hsa:05221	Acute myeloid leukaemia	0.00000205	< 10e – 6	6.36E – 12	< 10e – 6
hsa:00270	Cysteine and methionine metabolism	0.0000036	< 10e – 6	1.24E – 25	< 10e – 6
hsa:04530	Tight junction	0.00000531	< 10e – 6	6.85E – 18	< 10e – 6
hsa:04350	TGF-β-signalling pathway	0.00000725	< 10e – 6	5.93E – 14	< 10e – 6
hsa:04310	Wnt-signalling pathway	0.0000103	< 10e – 6	8.24E – 19	< 10e – 6
hsa:00590	Arachidonic acid metabolism	0.0000146	< 10e – 6	1.16E – 11	< 10e – 6
hsa:05213	Endometrial cancer	0.000018	< 10e – 6	0.00000131	< 10e – 6
hsa:04142	Lysosome	0.0000489	< 10e – 6	6.09E – 18	< 10e – 6

Abbreviations: BH, Benjamini-Hochberg; FDR, false discovery rate; MAPK, mitogen-activated protein kinase; mTOR, mammalian target of rapamycin; SAM-GS, significance analysis of microarrays for genesets; TGF, tumour growth factor; VEGF, vascular endothelial growth factor. The full pathway lists that show significance of differential enrichment in each individual data set are shown with their respective *P*-values of significance in Supplementary Table 2.

A



B




Figure 1 TP53 mutation status-specific network of potential candidate driver genes shown based on their known and predicted functional interactions. (A) Network for wild-type TP53 breast cancer profiles. (B) Network for mutant TP53 breast cancer profiles. Significant association of gene means significant non-zero regression coefficient of a gene in a significantly differentially enriched KEGG pathway. Gene upregulation means its class-specific upward biased expression pattern, inferred by the rank-sum statistic of the modified Kolmogorov–Smirnov test. Relevant biological processes represented by these genes are also highlighted in background.

Candidate genes deregulated according to the TP53 mutation status

Candidate genes were identified by applying a combination of two mutually complementary approaches: pathway-based gene-search that infers class-specific association (globaltest) and pathway-independent search that identifies individual genes with class-specific upregulation (modified Kolmogorov–Smirnov approach) on both primary and validation data sets (Supplementary Tables 3 and 4). Combining genesets inferred by these two approaches would help to account for the genes with smaller as well as larger effects on the overall biological condition. A consensus geneset (Supplementary Table 5) of 112 genes consists of genes inferred as significant at least by either of the two statistical tests (but not necessarily by the same test) in both primary and validation data sets, as shown in the Venn diagram (Supplementary Figure 3). Class-specific predicted functional networks based on these genesets are plotted in Figures 1A and B for wild-type and mutant TP53 samples, respectively. These networks reflect the key genes and corresponding processes that have potential functional implication in association with the one of the TP53 mutation status class. Wild-type TP53 samples showed significance of genes involved in estrogen receptor (ER) signalling, whereas mutant TP53 samples in proliferative processes. Besides, GO terms—response to insulin stimulus and mammary gland development in wild-type and protein kinase activity, mitotic cell cycle, microtubule cytoskeleton in mutant TP53 class were over-represented (Supplementary Figure 4).

Association of EMT and stemness to TP53 mutation status

Aberrant TP53 function is shown to induce epithelial-mesenchymal transition (EMT) and thereby confers stemness properties to the cancer cells (Dhar *et al*, 2008). Therefore, we compared our inferred TP53 status-specific candidate genesets with the published EMT and stemness marker sets. We found that mutant TP53-marker geneset was significantly associated with embryonic stem cell (ESC) and its TP53 targets (p53ESC) genesets (P -value < 0.05). Whereas wild-type TP53 signature was found significantly associated with PRC2 targets (P -value: 0.003) (Table 2). Top 1000 upregulated genes (according the signal-to-noise ratio) in mutant TP53 class were significantly associated with EMT, ESC and induced pluripotent stem cell marker genesets. Moreover, KEGG pathways involved in stemness and EMT properties such as TGF β , wnt signalling were found differentially enriched (Supplementary Table 6b).

Vascular endothelial growth factor A upregulation with wild-type TP53 associates with activation of pro-angiogenic and pro-metastatic biological processes

Among the inferred candidate genes that were found upregulated and/or significantly associated to one of the TP53 mutation status class, 47 genes showed univariate significance to overall patient survival. Vascular endothelial growth factor A (VEGFA)

Table 2 Association between the inferred TP53 mutation status-specific signatures with previously reported EMT and stemness markers. Statistical significance of differential expressed geneset overlapping the stemness and epithelial-mesenchymal transition (EMT) marker genes^a. Statistical significance was computed by applying hypergeometric test^b

EMT and stemness geneset and its transcript size	wtTP53 signature		Mutant TP53 signature		Top 1000 genes ranked acc to absolute SNR (wt vs mtTP53 BC)		Top 1000 mtTP53-upregulated genes ranked acc to SNR	
	Number of overlapping genes	P-value	Number of overlapping genes	P-value	Number of overlapping genes	P-value	Number of overlapping genes	P-value
EMT (n = 497)	0	NS	1	NS	11	NS	15	0.031
ESC (n = 553)	0	NS	14	2.65E - 13	22	2.60E - 04	35	4.34E - 11
PRC2 (n = 1016)	7	3.25E - 03	0	NS	25	NS	19	NS
iPSC (n = 597)	1	NS	3	NS	17	4.50E - 02	22	1.50E - 03
p53esc (n = 912)	2	NS	5	2.66E - 02	12	NS	15	NS

Abbreviations: BC, breast cancer; ESC, embryonic stem cell; iPSC, induced pluripotent stem cell; NS, not significant; p53esc, p53 targets identified in murine embryonic stem cells; PRC2, polycomb repressive complex 2; SNR, signal-to-noise ratio. ^aSources of the genesets are described in the Supplementary Table 6A. ^bStatistical significance was evaluated by Fisher's exact test, in instances where number of overlapping genes ≤ 5 .

maintained significance in multivariate model (Supplementary Table 7), even after adjusting for TP53 mutation status. VEGFA might be induced by estrogen receptor in breast cancer cells (Buteau-Lozano *et al*, 2002; Applanat *et al*, 2008). Besides, wild-type TP53 could block VEGFA function induced by active estrogen receptor signalling (Liang *et al*, 2005). However, implications of VEGFA in wild-type TP53/ER+ patients are less understood. We therefore analysed this subgroup separately by using the globaltest and moderated *t*-test (Smyth, 2004).

Using moderated *t*-test of differential expression on a cross-platform compiled data set, we found 516 gene features (Supplementary Table 8a) differentially expressed between VEGFA upregulation (VEGFA+) vs VEGFA normal/- samples (VEGFA-/N). IGF1 and PPARG were found differentially down-regulated in samples with VEGFA upregulation. A GO analysis identified pathways associated with blood vessel morphogenesis, cell migration and regulation of VEGF signalling pathway. The complete list of over-represented GO terms and predicted functional interactions are shown in Supplementary Table 9 and Supplementary Figure 5, respectively. Notably, VEGFA+ vs VEGFA-/N comparison for mutant TP53 subgroup does not show any remarkable difference apart from differential expression of VEGFA itself and pH regulator CA9 (Supplementary Table 8b).

Tumours overexpressing VEGFA (both ER+ wild-type TP53 and mutant TP53 irrespective of ER status) show a differential enrichment of the mTOR-signalling pathway compared with normal/downregulated VEGFA samples. VEGFA+/ER+ wild-type TP53 samples showed significant association of EIF4EBP1, MAPK1 (*P*-value < 0.05) and weak association of MTOR, ULK3 and RPTOR. Conversely, PIK3CA and IGF1 were significantly associated with VEGFA N/- tumours (Supplementary Figure 5 and Supplementary Table 10a). Interestingly, different sets of genes, although involved in the same pathways were found associated with VEGFA status in the mutant TP53 subgroup (Supplementary Table 10b).

TP53 mutation, ER status and VEGFA upregulation influence survival

Samples were stratified according to the ER status in each TP53 mutation class. While comparing the ER+/mutant TP53 to the ER+/wild-type TP53 samples, we noted a death hazard ratio (HR) of 2.15 (95% CI: 1.25-3.70) and likelihood *P*-value < 0.01. On the other hand, ER- samples showed weaker significance (*P* = 0.2; HR: 2.6; 95% CI: 1.14-5.91). As progesterone receptor (PgR) positivity is a better marker of active ER signalling (Bardou *et al*, 2003), we also used PgR status as an indicator of active ER signalling. PgR+ samples showed a

significant survival difference between mutant and wild-type tumours (*P* = 1.53e - 05, HR: 7.2, 95% CI: 3.03-17.1). However, PgR-tumours do not show significant survival differences (*P* > 0.1) (Figure 2A and B). On the basis of these findings, we propose that active ER signalling can influence the effect of mutant TP53 on survival.

As VEGFA expression is observed here as a significant influencer on survival even after controlling for TP53 status, we reanalyzed the above effects by adding VEGFA expression status as a covariate. Among ER+ group, the overall patient survival was significantly influenced by TP53 mutation status and VEGFA (model significance = 0.0005) with their corresponding HR = 2.02 and 2.08, compared with baseline risk for wild-type TP53 and VEGFA normal/downregulation. Even stronger effect was observed after excluding samples with non-missense mutant TP53 (*P*-value = 0.0001, HR = 2.38 and 2.11, respectively). Survival effect of TP53 mutation status and VEGFA was stronger in PgR+ cases (HR = 2.35, 95% CI: 1.17-4.74 for VEGFA upregulation, HR = 5.2, 95% CI: 2.43-11.1 for mutant TP53 status, and overall likelihood ratio test *P* = 2.76e - 6), but non-significant effect in PgR-cases (Figure 2C and D). Although active ER signalling in general is known to predict better prognosis, these findings show that irrespective of the TP53 mutation status, ER+ cases with high mRNA levels of VEGFA indicates poor prognosis. Interestingly, despite of the lowest occurrence of cases with upregulated VEGFA in ER+/wtTP53 subgroup (Supplementary Figure 6), its prognostic significance underscores further exploration.

DISCUSSION

Our findings show predominance of ER signalling in breast cancers with wild-type TP53, marked by the upregulation of ESR1, GATA-binding protein 3, retinoic acid receptor alpha (RAR α) and CA12. Estrogen receptor α , a direct transcriptional activator of RAR α (Han *et al*, 1997), mediates anti-proliferative response by vitamin A metabolite (all-trans-retinoic acid) in breast cancer cells (Dawson *et al*, 1995). Retinoic acid receptor α is a rate-limiting factor for ER transcriptional activity (Ross-Innes *et al*, 2010). Co-expression of BCL2, ERBB4, IGF1R, IRS1 was also found in this group. Our observation of consistent upregulation of CA12, AGR3, IL6ST and STC2 genes is in agreement with their previously reported association with ER+ breast cancers. Our findings also showed upregulation of SIRT3, a mitochondrial p53 activity regulator, necessary for averting TP53-mediated growth arrest (Li *et al*, 2010). Predicted functional network (Figure 1A) provides a hint that genes involved in ER signalling form a core group of




Figure 2 Overall patient survival differs significantly according to the *TP53* mutation status and *VEGFA* expression status in PgR + and PgR - subgroups of patients. Survival differences between wild-type *TP53* and mutant *TP53* in each of the subgroups are shown in Kaplan-Meier plots shown in **A** and **B**. Survival differences of four classes: (1) wild-type *TP53* and *VEGFA* normal/downregulation (wt*TP53* *VEGFA* N/-); (2) wild-type *TP53* and *VEGFA* upregulation (wt*TP53* *VEGFA* +); (3) mutant *TP53* and *VEGFA* normal/downregulation (mt*TP53* *VEGFA* N/-); and (4) mutant *TP53* and *VEGFA* upregulation (mt*TP53* *VEGFA* +) - in PgR + and PgR - subgroups are shown in **C** and **D**. Significance of overall model is based on the likelihood ratio test *P*-value.

interactions in *TP53* wild-type tumours. A strong relationship between ER signalling and *TP53* can be observed in our results. This relationship also has got implications on proliferation and treatment responsiveness. The presence of wild-type *TP53* improves sensitivity to Tamoxifen (Berns *et al*, 2000) and inhibits ER cross-talk with the EGFR/HER2 pathways (Fernandez-Cuesta *et al*, 2010). Experimental observations have provided evidence about potential direct ER-*TP53* interactions (Liu *et al*, 2006). However, these complex interactions and their effects on transactivation activity of *TP53* and ER α in ER+ breast cancer remains to be understood. Given that *TP53* status is an important predictor of response in patients receiving therapy targeting the ER pathway (SERM), we expect that *TP53* retains a subset of functions necessary for the response to such therapy.

Genes in pathways related to cell cycle, angiogenesis, chromosomal instability and metastasis were significantly affected in mutant *TP53* tumours. We found the gene *BUB1* and spindle-checkpoint associated kinases were significantly associated with *TP53* mutant tumours. In the presence of dysfunctional *TP53*, their aberrant expression can cause genomic instability, leading to aneuploidy and malignant transformation (Gjoerup *et al*, 2007). Other genes associated with mutant *TP53* included ones involved in proliferation, angiogenesis and metastasis-*VEGFA*, *HIF1 α* , *E2F1*, *CDK6* and *EGFR*.

VEGFA upregulation is an important indicator of pro-angiogenic and pro-metastatic activity. Dysregulation of *TP53*-*VEGF* signalling may potentially be a key event in breast cancers with

mutant *TP53*. Mutant *TP53* may facilitate this tumorigenic programme by: passing the direct survival advantage to malignant cells, by facilitating the VEGF-mediated enhanced cell migration, angiogenesis and metastasis or by overcoming the regulation by *ETS1* (Dittmer, 2003). Active ER signalling and mutant *TP53* are also reported to activate VEGF and mark poor prognosis (Berns *et al*, 2003). In our data, we see that mutant *TP53* and VEGF upregulation significantly affects patient survival in ER+/PgR+ samples, but not in ER-/PgR- samples. Activation of *VEGFA* may also be attributed to the expression of EGFR (Maity *et al*, 2000) or *CDK6*, which can correlate with the expression of mutant *TP53* (Wyllie *et al*, 2003) and potentially delay cell senescence. Thus, besides the direct effects of lost *TP53* function, other related opportunistic mechanisms, such as dysregulated proliferative effects of *VEGFA* may contribute the overall manifestation.

ER+/wild-type *TP53* samples showed relatively low occurrence of *VEGFA* upregulation but poor survival profile. ER-mediated induction of VEGF (Berns *et al*, 2003; Applanat *et al*, 2008) and VEGF regulation by *TP53* (Liang *et al*, 2005) suggests a complex interplay between these three signalling mechanisms. This group also showed the differential enrichment of mTOR signalling. Co-activation of VEGF and mTOR pathway components has been previously reported (Trinh *et al*, 2009). Thus, *VEGFA* may represent a biomarker of interest to identify the target subset of ER+ breast cancer patients who might benefit from early administration of *VEGFA* or mTOR-targeted therapy.

MATERIALS AND METHODS

Agilent chip based gene expression data for a subset of 111 breast cancer cases from (Enerly *et al*, 2011) GEO (accession number GSE19783) was used as the primary data set. *TP53* mutations for the primary data in coding regions of exons 2–11 and clinical data were obtained from (Naume *et al*, 2007). Expression data used for validation was obtained from GEO (accession number GSE3494) and from Stanford Microarray Database. Clinical and *TP53* data for these data sets were obtained from (Miller *et al*, 2005; Langerød *et al*, 2007).

Methods used to merge data sets to form a validation data set

Two expression data sets (Miller *et al*, 2005; Langerød *et al*, 2007) from independent studies and different technology platform were preprocessed, quantile normalised and combined based on UniGene identifiers. Batch effects were corrected by applying parametric empirical Bayes method (Johnson *et al*, 2007).

Differential enrichment of pathways and candidate genes

The globaltest (Goeman *et al*, 2011) uses a regression model where genes are covariates and sample classes are response variables. Significant association of gene means significant non-zero regression coefficient of a gene in a geneset (here a particular KEGG pathway). SAM-GS is another geneset enrichment analysis method based on the *t*-like statistic for assessing the permutation-based significance of association between an individual pathway and a phenotype of interest. KEGG pathways inferred as significant by globaltest at FDR corrected *P*-value of $10e-5$ and validated by SAM-GS (Dinu *et al*, 2007) at FDR corrected *P*-value cutoff = $10e-6$ on both primary and validation data sets were analysed by *post-hoc* covariate test to identify significant genes. Gene upregulation means its class-specific upward biased expression pattern, inferred by the rank-sum statistic of the modified Kolmogorov-Smirnov test (Yang *et al*, 2010).

Class-specific predicted functional interactions between genes in the genesets were obtained from STRING database (Jensen *et al*, 2009).

Pathways enrichment and GO analysis

Gene Ontology (GO) analysis was performed for each *TP53* mutation status-specific genesets using DAVID (Huang *et al*, 2009) by Fisher's exact test with human whole genome as a background. Differentially enriched pathways and GO terms were graphically

REFERENCES

- Aas T, Børresen AL, Geisler S, Smith-Sørensen B, Johnsen H, Varhaug JE, Akslen LA, Lønning PE (1996) Specific P53 mutations are associated with de novo resistance to doxorubicin in breast cancer patients. *Nat Med* 2: 811–814
- Applanat MP, Buteau-Lozano H, Herve MA, Corpet A (2008) Vascular endothelial growth factor is a target gene for estrogen receptor and contributes to breast cancer progression. *Adv Exp Med Biol* 617: 437–444
- Bardou V-J, Arpino G, Elledge RM, Osborne CK, Clark GM (2003) Progesterone receptor status significantly improves outcome prediction over estrogen receptor status alone for adjuvant endocrine therapy in two large breast cancer databases. *J Clin Oncol* 21: 1973–1979
- Berns EM, Foekens JA, Vossen R, Look MP, Devilee P, Henzen-Logmans SC, Van Staveren IL, Van Putten WL, Inganas M, Meijer-Van Gelder ME, Cornelisse C, Claassen CJ, Portengen H, Bakker B, Klijn JG (2000) Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer. *Cancer Res* 60: 2155–2162
- Berns EMJJ, Klijn JGM, Look MP, Grebenchtchikov N, Vossen R, Peters H, Geurts-Moespot A, Portengen H, Van Staveren IL, Meijer-Van Gelder ME, Bakker B, Sweep FCGJ, Foekens JA (2003) Combined vascular endothelial growth factor and TP53 status predicts poor response to

presented as Enrichment map (Merico, 2009), with nodes color-coded by FDR-adjusted *P*-value of significance and node-size proportionate to number of genes in the pathway. Fraction of overlapping genes between any two pathways is represented by the edge thickness, with cutoff overlap coefficient of 0.1.

Association of TP53 biology with EMT and stemness marker signatures

Inferred class-specific genesets were tested by hypergeometric test for their association to the published EMT and stemness marker genesets shown in Supplementary Table 6a. A larger genelist inferred by using signal-to-noise ratio between *TP53* mutation status classes was also tested for its association to these published genesets.

Survival analysis

A combined cohort of 438 cases obtained by merging clinical data from three individual clinical data sets (Supplementary Table 1) was used. Kaplan–Meier estimation of survival and computation of Cox proportional hazards frailty model for the death event was performed by using R package *survival* (Therneau and Lumley, 2009). Inferred candidate genes were assessed for their uni-/multivariate effect on survival. The effect of *TP53* mutation status together with genes that maintain significance in a multivariate model (*VEGFA* expression status) and predicted subtype (Parker *et al*, 2009) was computed with and without stratification by ER/PgR status.

Discretisation of gene expression

The mRNA expression levels of candidate genes were discretised into two levels using mean (μ) + 0.5*standard deviation (s.d.) as a cutoff in each data set.

Analysis was performed by using R (R Development Core Team, 2011).

ACKNOWLEDGEMENTS

This paper is a part of the doctoral thesis work of HJ. His work was supported by grant number 2789119 from Helse Sør-Øst and internal grants from Akershus University Hospital (number: 2679030 and 2699015 to VNK).

Supplementary Information accompanies the paper on British Journal of Cancer website (<http://www.nature.com/bjc>)

- tamoxifen therapy in estrogen receptor-positive advanced breast cancer. *Clin Cancer Res* 9: 1253–1258
- Bertheau P, Turpin E, Rickman DS, Espié M, De Reyniès A, Feugeas J-P, Plassa L-F, Soliman H, Varna M, De Roquancourt A, Lehmann-Che J, Beuzard Y, Marty M, Misset J-L, Janin A, De Thé H (2007) Exquisite sensitivity of TP53 mutant and basal breast cancers to a dose-dense epirubicin – cyclophosphamide regimen. *PLoS Med* 4: 10
- Blandino G, Levine AJ, Oren M (1999) Mutant p53 gain of function: differential effects of different p53 mutants on resistance of cultured cells to chemotherapy. *Oncogene* 18(2): 477–485
- Børresen AL, Andersen TI, Eyfjörd JE, Cornelis RS, Thorlacius S, Borg A, Johansson U, Theillet C, Scherneck S, Hartman S (1995) TP53 mutations and breast cancer prognosis: particularly poor survival rates for cases with mutations in the zinc-binding domains. *Genes Chromosomes Cancer* 14: 71–75
- Buteau-Lozano H, Ancelin M, Lardeux B, Milanini J, Perrot-Applanat M (2002) Transcriptional regulation of vascular endothelial growth factor by estradiol and tamoxifen in breast cancer cells: a complex interplay between estrogen receptors alpha and beta. *Cancer Res* 62(17): 4977–4984
- D'Assoro AB, Leontovich A, Amato A, Ayers-Ringler JR, Quatraro C, Hafner K, Jenkins RB, Libra M, Ingle J, Stivala F, Galanis E, Salisbury JL

- (2010) Abrogation of p53 function leads to metastatic transcriptome networks that typify tumor progression in human breast cancer xenografts. *Int J Oncol* 37: 1167–1176
- Dawson MI, Chao WR, Pine P, Jong L, Hobbs PD, Rudd CK, Quick TC, Niles RM, Zhang XK, Lombardo A (1995) Correlation of retinoid binding affinity to retinoic acid receptor alpha with retinoid inhibition of growth of estrogen receptor-positive MCF-7 mammary carcinoma cells. *Cancer Res* 55(19): 4446–4451
- Dhar G, Banerjee S, Dhar K, Tawfik O, Mayo MS, Vanveldhuizen PJ, Banerjee SK (2008) Gain of oncogenic function of p53 mutants induces invasive phenotypes in human breast cancer cells by silencing CCN5/WISP-2. *Cancer Res* 68(12): 4580–4587
- Dinu I, Potter JD, Mueller T, Liu Q, Adewale AJ, Jhangri GS, Einecke G, Famulski KS, Halloran P, Yasui Y (2007) Improving gene set analysis of microarray data by SAM-GS. *BMC Bioinformatics* 8: 242
- Dittmer J (2003) The Biology of the Ets1 Proto-Oncogene. *Mol Cancer* 2: 29
- Enerly E, Steinfeld I, Kleivi K, Leivonen S-K, Aure MR, Russnes HG, Rønneberg JA, Johnsen H, Navon R, Røddland E, Mäkelä R, Naume B, Perälä M, Kallioniemi O, Kristensen VN, Yakhini Z, Børresen-Dale A-L (2011) miRNA-mRNA Integrated Analysis Reveals Roles for miRNAs in Primary Breast Tumors. *PLoS One* 6: 13
- Fernandez-Cuesta L, Anaganti S, Hainaut P, Olivier M (2010) p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines. *Int J Cancer* 128(8): 1813–1821
- Gjoerup OV, Wu J, Chandler-Militello D, Williams GL, Zhao J, Schaffhausen B, Jat PS, Roberts TM (2007) Surveillance mechanism linking Bub1 loss to the p53 pathway. *Proc Natl Acad Sci USA* 104: 8334–8339
- Goeman JJ, Van Houwelingen HC, Finos L (2011) Testing against a high-dimensional alternative in the generalized linear model: asymptotic type I error control. *Biometrika* 98: 381–390
- Han QX, Allegretto EA, Shao ZM, Kute TE, Ordonez J, Aisner SC, Rishi AK, Fontana JA (1997) Elevated expression of retinoic acid receptor-alpha (RAR alpha) in estrogen-receptor-positive breast carcinomas as detected by immunohistochemistry. *Am J Surg Pathol Part B* 6(1): 42–48
- Huang DW, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4(1): 44–57
- Jensen LJ, Kuhn M, Stark M, Chaffron S, Creevey C, Muller J, Doerks T, Julien P, Roth A, Simonovic M, Bork P, Von Mering C (2009) STRING 8 – a global view on proteins and their functional interactions in 630 organisms. *Nucl Acids Res* 37: D412–D416
- Johnson WE, Li C, Rabinovic A (2007) Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics* 8: 118–127
- Kandioler-Eckersberger D, Ludwig C, Rudas M, Kappel S, Janschek E, Wenzel C, Schlagbauer-Wadl H, Mittlböck M, Gnant M, Steger G, Jakesz R (2000) TP53 mutation and p53 overexpression for prediction of response to neoadjuvant treatment in breast cancer patients. *Clin Cancer Res* 6: 50–56
- Kanehisa M, Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 28(1): 27–30
- Kim E, Deppert W (2006) The versatile interactions of p53 with DNA: when flexibility serves specificity. *Cell Death Differ* 13(6): 885–889
- Langerød A, Zhao H, Borgan Ø, Nesland JM, Bukholm IR, Ikdahl T, Kåresen R, Børresen-Dale A-L, Jeffrey SS (2007) TP53 mutation status and gene expression profiles are powerful prognostic markers of breast cancer. *Breast Cancer Res* 9(3): R30
- Li S, Banck M, Mujtaba S, Zhou M-M, Sugrue MM, Walsh MJ (2010) p53-Induced Growth Arrest Is Regulated by the Mitochondrial SirT3 Deacetylase. *PLoS One* 5: 12
- Liang Y, Wu J, Stancel GM, Hyder SM (2005) p53-dependent inhibition of progesterin-induced VEGF expression in human breast cancer cells. *J Steroid Biochem Mol Biol* 93(2-5): 173–182
- Liu Q, Dinu I, Adewale AJ, Potter JD, Yasui Y (2007) Comparative evaluation of gene-set analysis methods. *BMC Bioinformatics* 8: 431
- Liu W, Konduri SD, Bansal S, Nayak BK, Rajasekaran SA, Karuppaiy SM, Rajasekaran AK, Das GM (2006) Estrogen receptor-alpha binds p53 tumor suppressor protein directly and represses its function. *J Biol Chem* 281(15): 9837–9840
- Maity A, Pore N, Lee J, Solomon D, O'Rourke DM (2000) Epidermal growth factor receptor transcriptionally up-regulates vascular endothelial growth factor expression in human glioblastoma cells via a pathway involving phosphatidylinositol 3'-kinase and distinct from that induced by hypoxia. *Cancer Res* 60(20): 5879–5886
- Merico D (2009) Enrichment map: a network-based method for gene-set enrichment visualization and interpretation. *PLoS One* 5: e13984
- Miller LD, Smeds J, George J, Vega VB, Vergara L, Ploner A, Pawitan Y, Hall P, Klaar S, Liu ET, Bergh J (2005) An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival. *Proc Natl Acad Sci USA* 102(38): 13550–13555
- Naume B, Zhao X, Synnestevedt M, Borgen E, Russnes HG, Lingjaerde OC, Strømberg M, Wiedswang G, Kvalheim G, Kåresen R, Nesland JM, Børresen-Dale A-L, Sørli T (2007) Presence of bone marrow micro-metastasis is associated with different recurrence risk within molecular subtypes of breast cancer. *Mol Oncol* 1(2): 160–171
- Norberg T, Klaar S, Kärf G, Nordgren H, Holmberg L, Bergh J (2001) Increased p53 mutation frequency during tumor progression – results from a breast cancer cohort. *Cancer Res* 61: 8317–8321
- O'Shaughnessy J (2005) Extending survival with chemotherapy in metastatic breast cancer. *Oncologist* 10(Suppl 3): 20–29
- Olivier M, Langerød A, Carrieri P, Bergh J, Klaar S, Eyfjord J, Theillet C, Rodriguez C, Lidereau R, Bièche I, Varley J, Bignon Y, Uhrhammer N, Winqvist R, Jukkola-Vuorinen A, Niederacher D, Kato S, Ishioka C, Hainaut P, Børresen-Dale A-L (2006) The clinical value of somatic TP53 gene mutations in 1,794 patients with breast cancer. *Clin Cancer Res* 12: 1157–1167
- Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T, Davies S, Fauron C, He X, Hu Z, Quackenbush JF, Stijleman IJ, Palazzo J, Marron JS, Nobel AB, Mardis E, Nielsen TO, Ellis MJ, Perou CM, Bernard PS (2009) Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol* 27(8): 1160–1167
- R Development Core Team (2011) R: A Language and Environment for Statistical Computing. Vienna Austria R Foundation for Statistical Computing 1: ISBN 3-900051-07-0
- Ross-Innes CS, Stark R, Holmes KA, Schmidt D, Spyrou C, Russell R, Massie CE, Vowler SL, Eldridge M, Carroll JS (2010) Cooperative interaction between retinoic acid receptor-alpha and estrogen receptor in breast cancer. *Genes Dev* 24: 171–182
- Smyth GK (2004) Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 3: Article 3
- Therneau T, Lumley T (2009) Survival: Survival Analysis Including Penalised Likelihood. R package Version 2.35-8. R package version 2.35-8. <http://CRAN.R-project.org/package=survival>
- Trinh XB, Tjalma WA, Vermeulen PB, Van den Eynden G, Van der Auwera I, Van Laere SJ, Helleman J, Berns EM, Dirix LY, van Dam PA (2009) The VEGF pathway and the AKT/mTOR/p70S6K1 signalling pathway in human epithelial ovarian cancer. *Br J Cancer* 100(6): 971–978
- Wyllie F, Houghton M, Bartek J, Rowson J, Wynford-Thomas D (2003) Mutant p53 can delay growth arrest and loss of CDK2 activity in senescing human fibroblasts without reducing p21(WAF1) expression. *Exp Cell Res* 285(2): 236–242
- Yang Y, Kort EJ, Ebrahimi N, Zhang Z, Teh BT (2010) Dual KS: defining gene sets with tissue set enrichment analysis. *Cancer Inform* 9: 1–9



This work is licensed under the Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>