Keywords: pancreatic cancer; pancreatic intraepithelial neoplasia; non-small-cell lung cancer; acinar to ductal metaplasia; oncogenic Kras; Kras-driven cancer; oncogenic Kras signalling; metabolism; glycolysis; glutaminolysis; PI3K; Pdk1; Akt; MAPK

Oncogenic KRAS signalling in pancreatic cancer

S Eser^{1,2}, A Schnieke³, G Schneider¹ and D Saur^{*,1,2}

¹Department of Internal Medicine II, Klinikum rechts der Isar, Technische Universität München, Ismaningerstr. 22, 81675 München, Germany; ²German Cancer Consortium (DKTK) and German Cancer Research Center (DKFZ), Im Neuenheimer Feld 280, 69120 Heidelberg, Germany and ³Livestock Biotechnology, Technische Universität München, Liesel-Beckmann Str. 1., 85354 Freising, Germany

Pancreatic ductal adenocarcinoma (PDAC) is almost universally fatal. The annual number of deaths equals the number of newly diagnosed cases, despite maximal treatment. The overall 5-year survival rate of <5% has remained stubbornly unchanged over the last 30 years, despite tremendous efforts in preclinical and clinical science. There is unquestionably an urgent need to further improve our understanding of pancreatic cancer biology, treatment response and relapse, and to identify novel therapeutic targets. Rigorous research in the field has uncovered genetic aberrations that occur during PDAC development and progression. In most cases, PDAC is initiated by oncogenic mutant KRAS, which has been shown to drive pancreatic neoplasia. However, all attempts to target KRAS directly have failed in the clinic and KRAS is widely assumed to be undruggable. This has led to intense efforts to identify druggable critical downstream targets and nodes orchestrated by mutationally activated KRAS. This includes context-specific KRAS effector pathways, synthetic lethal interaction partners and KRAS-driven metabolic changes. Here, we review recent advances in oncogenic KRAS signalling and discuss how these might benefit PDAC treatment in the future.

PANCREATIC DUCTAL ADENOCARCINOMA IS DRIVEN BY ONCOGENIC KRAS

Pancreatic ductal adenocarcinoma (PDAC), the predominant form of pancreatic cancer, develops via acinar-ductal metaplasia (ADM) and neoplastic precursor lesions, such as pancreatic intraepithelial neoplasia (PanIN), intraductal papillary mucinous neoplasia (IPMN), mucinous cystic neoplasia and atypical flat lesions (AFLs; Morris *et al*, 2010; Aichler *et al*, 2012). Mutationally activated KRAS is present in >90% of PDAC and represents the most frequent (>90%) and the earliest genetic alteration, being found in low-grade PanIN 1A lesions (Morris *et al*, 2010; Kanda *et al*, 2012).

The KRAS proto-oncogene encodes an ~21 kDa small GTPase, which cycles between GTP-bound active and GDP-bound inactive states. The switch to the active state is promoted by guanine nucleotide exchange factors (GEFs), which aid exchange of GDP for GTP. KRAS inactivation is mediated by GTPase-activating proteins (GAPs), which induce hydrolysis of GTP. Activating mutations of KRAS found in human PDAC (point mutations at codon G12 (98% of all KRAS mutations in PDAC), G13 and Q61)

impair intrinsic GTPase activity of the KRAS protein and can block the interaction between KRAS and GAPs. This leads to constitutive activation of KRAS and persistent stimulation of downstream signalling pathways that drive many of the hallmarks of cancer, sustained proliferation, metabolic reprogramming, anti-apoptosis, remodelling of the tumour microenvironment, evasion of the immune response, cell migration and metastasis (Pylayeva-Gupta *et al*, 2011).

Targeting of mutant Kras^{G12D} or Kras^{G12V} specifically to the murine pancreas is sufficient to initiate development of ADM, PanINs, IPMNs and AFLs, which progress with long latency to invasive metastatic PDAC, thus recapitulating the human disease (Hingorani *et al*, 2003; Guerra *et al*, 2007; Seidler *et al*, 2008; Morris *et al*, 2010; Pylayeva-Gupta *et al*, 2011). The low frequency of spontaneous progression of precursor lesions to invasive PDAC suggests that additional genetic aberrations are needed for disease progression (Morris *et al*, 2010). Pancreatic intraepithelial neoplasia and PDAC development can indeed be accelerated in Kras-driven mouse models by introducing inactivating mutations in tumour suppressor genes *Cdkn2a*, *Trp53*, or *Dpc4/Smad4*,

*Correspondence: Professor D Saur; E-mail: Dieter.Saur@lrz.tum.de

Received 28 November 2013; revised 19 March 2014; accepted 26 March 2014; published online 22 April 2014

© 2014 Cancer Research UK. All rights reserved 0007–0920/14

all of which occur frequently in human lesions as they progress to invasive PDAC (Jones *et al*, 2008; Morris *et al*, 2010; Biankin *et al*, 2012).

These mouse studies showed that oncogenic Kras is capable of initiating PDAC, but could not investigate whether continuous Kras activity is required for maintenance of PanINs and PDAC. Recently developed mouse models in which the Kras oncogene can be switched on and off have impressively demonstrated that continuous oncogenic Kras signalling is essential for both progression and maintenance of PDAC (Ying *et al*, 2012; Collins *et al*, 2012a). In addition, it became evident that sustained oncogenic Kras signalling is also necessary for the growth and maintenance of metastatic lesions (Collins *et al*, 2012b). However, individual tumour cells can remain dormant over a long time period after Kras inactivation. Accordingly, reactivation of oncogenic Kras in these dormant cancer cells leads to rapid disease progression (Collins *et al*, 2012b). This accords with studies of human pancreatic cancer cell lines that are addicted to oncogenic KRAS for sustained proliferation and survival (Zimmermann *et al*, 2013).

Together, these findings place oncogenic KRAS squarely at the top of the list of therapeutic targets. However, a subset of human PDAC cells is known to resist KRAS inactivation, a phenotype associated with an epithelial to mesenchymal transition (Singh *et al*, 2009). This highlights the importance of the genetic heterogeneity of human PDAC. Whole-exome and genome sequencing has revealed that human PDAC is an extremely heterogeneous disease with diverse molecular subtypes (Jones *et al*, 2008; Biankin *et al*, 2012; Cowley *et al*, 2013). Such *in vitro* findings therefore suggest that only a subset of pancreatic cancer patients will benefit from KRAS inhibition. This view is supported by an outstanding gene expression profiling study, which revealed three distinct subtypes of pancreatic cancer. One, termed the 'classical subtype', represents 41.2% of the analysed pancreatic cancer cases, has high expression of epithelial genes, and was found to be strongly dependent on constitutive KRAS signalling (Collisson *et al*, 2011). The 'quasi-mesenchymal' and the 'exocrine-like' subtypes were found in 36.5% and 22.3%, respectively (Collisson *et al*, 2011). These distinctions clearly have a broad impact on clinical practice, and it is vital that the molecular determinants be defined.

Because oncogenic KRAS drives PDAC maintenance, at least in a large subset of PDAC patients, there have been considerable efforts to develop direct inhibitors. However, all clinical attempts to directly interfere with KRAS oncoprotein activity have failed, and KRAS is still widely considered undruggable (Berndt *et al*, 2011). A new approach is to identify and characterise druggable crucial downstream effectors of oncogenic KRAS. We now focus on effector pathways of oncogenic KRAS that contribute to PDAC initiation, progression and maintenance and how these pathways can be addressed therapeutically.

EFFECTOR PATHWAYS OF ONCOGENIC KRAS

KRAS signalling is highly complex and dynamic, engaging various downstream effectors, such as canonical Raf/Mek/Erk, phosphatidylinositol 3-kinase (PI3K)/3-phosphoinositide-dependent protein kinase-1 (Pdk1)/Akt, RalGDS/p38MAPK, Rac and Rho, Rassf1, NF1, p120GAP and PLC- ϵ (Castellano and Downward, 2011; Pylayeva-Gupta *et al*, 2011). It is believed that oncogenic KRAS signalling in PDAC passes through three major pathways: Raf/Mek/Erk, PI3K/Pdk1/Akt and the Ral guanine nucleotide exchange factor pathway (Lim *et al*, 2005; Feldmann *et al*, 2010; Collisson *et al*, 2012; Eser *et al*, 2013).

Recent findings in mice showed that pancreatic cancer initiation, progression and maintenance depend on tumour cell autonomous

Kras-PI3K-Pdk1 signalling. Pancreas-specific expression of PIK3CA^{H1047R} (p110 α ^{H1047R}), a constitutively active oncogenic class IA phosphatidylinositol 3-kinase, in Ptf1a-positive cells enabled selective activation of the PI3K/Pdk1/Akt pathway and phenocopied Kras^{G12D}-driven pancreatic carcinogenesis with striking similarity. In this model, oncogenic signalling flowed through Pdk1 and Akt without cross-activation of Kras, showing that the PI3K-Pdk1 axis represents a pathway capable of inducing cell plasticity, ADM, PanIN and PDAC formation (Eser *et al*, 2013). It is important to mention that contrasting findings have been reported using another murine model. Expression of p110 α ^{H1047R} in Pdx1-positive cells using a tamoxifen activatable *Pdx1-CreER* mouse line failed to induce PanIN and PDAC formation (Collisson *et al*, 2012). Different target cells of distinct Cre-driver lines, differences in recombination efficacy, or different expression and signalling levels of p110 α might explain these opposing results.

Importantly, genetic proof of the importance of PI3K-Pdk1 signalling was shown in the classical Kras^{G12D}-driven PDAC model. Genetic inactivation of *Pdk1* (*PDPK1*) completely blocked the development of ADM, PanIN and PDAC (Eser *et al*, 2013). This was evident in several mouse models using different Cre-driver lines, clearly demonstrating that the PI3K/Pdk1 pathway is essentially engaged by oncogenic Kras and needed for PDAC formation.

Collisson and colleagues showed that selective activation of the Raf-Mek-Erk pathway by expression of a conditional mutant oncogenic *Braf*^{V600E} allele in murine pancreas induces PanIN and PDAC development. In this model, activation of the oncogene resulted in a more aggressive phenotype with more PanINs compared with the classical Kras^{G12D} model (Collisson *et al*, 2012). Thus, activation of the canonical MAPK pathway at the level of Raf is sufficient to drive neoplastic changes in pancreas. However, the contribution of Braf to Kras-driven pancreatic carcinogenesis remains unclear. Interestingly, *Craf* has no role in PDAC development in the *Ptf1a*^{Cre/+}; *LSL-Kras*^{G12D/+} PDAC model, although it is known to be important for Kras^{G12D}-driven non-small-cell lung carcinogenesis (Blasco *et al*, 2011; Karreth *et al*, 2011; Eser *et al*, 2013).

The Ral guanine nucleotide exchange factor (RalGEFs) pathway has also been implicated in pancreatic carcinogenesis and progression. RalGEFs, which load GTP to small GTPases of the RAS superfamily, are necessary for RAS-induced transformation of several human cell types (Lim *et al*, 2005). They activate the RAS-like small GTPases RAL-A and RAL-B, which function as mediators of tumour growth and metastasis in human PDAC cell lines, respectively (Lim *et al*, 2006). High levels of active GTP-bound RAL-A and RAL-B have been found in human PDAC (Lim *et al*, 2006). Inhibition of cyclin-dependent kinase 5 (CDK5) had a marked anti-tumourigenic effect on KRAS mutant PDAC cell lines *in vitro* and *in vivo*, which was attributed to reduced RAS-RAL signalling (Feldmann *et al*, 2010). CDK5 inhibition also reduced levels of RHO-GTP and RAC-GTP in these cell lines (Feldmann *et al*, 2010). These proteins belong to the RAS superfamily of small GTPases and are deregulated in PDAC. Interestingly, Rac1 is dispensable for pancreas development, but has an important role in regulating the actin cytoskeleton during metaplastic transdifferentiation in the early stages of pancreatic carcinogenesis (Heid *et al*, 2011). However, expression of a constitutively active form of Rac1 without concomitant expression of oncogenic Kras does not induce pancreatic carcinogenesis (Eser *et al*, 2013).

CONTEXT SPECIFICITY OF ONCOGENIC KRAS SIGNALLING

The therapeutic efficacy of inhibitors that target distinct effector pathways of oncogenic KRAS is likely to vary significantly between

KRAS-driven tumour types. Recent data show that different downstream effectors are engaged by the Kras oncoprotein in PDAC and non-small-cell lung cancer (NSCLC) (Eser *et al*, 2013). The concept of tissue- and context-specific oncogenic signalling is illustrated by the need for Craf in KRAS^{G12D}-driven NSCLC (Blasco *et al*, 2011; Karreth *et al*, 2011), but not KRAS^{G12D}-driven pancreatic carcinogenesis (Eser *et al*, 2013). Furthermore, KRAS^{G12D}-driven PDAC depends completely on signalling via the PI3K effector Pdk1, whereas KRAS^{G12D}-driven NSCLC is unaffected by loss of *Pdk1* (Eser *et al*, 2013). This has significant implications for clinical therapy. Indeed, previous pharmacological studies in PDAC and NSCLC have suggested tissue-specific differences in KRAS signalling. Engelman *et al* (2008) found no substantial response of Kras^{G12D}-driven NSCLC to PI3K-mTOR inhibition by NPV-Bez235 *in vivo*. Although genetic ablation of the RAS/PIK3CA interaction induced regression of Kras-driven NSCLC *in vivo*, treatment with either a class IA PI3K inhibitor or a p110 α isoform-specific inhibitor alone showed modest anti-tumour effects (Castellano *et al*, 2013). However, when either one of these agents was combined with a Mek1/2 inhibitor, striking tumour shrinkage was observed (Engelman *et al*, 2008; Castellano *et al*, 2013). In pancreatic cancer, a clinically available inhibitor of class IA PI3Ks efficiently blocked tumour progression in Kras^{G12D}-driven PDAC *in vivo* (Eser *et al*, 2013). Input from receptor tyrosine kinases (RTKs) in Kras mutant tumours also seems to be tissue dependent, because elimination of the epidermal growth factor receptor (EGFR) in a mouse model of Kras^{G12D}-driven NSCLC failed to recapitulate the inhibitory effect seen on Kras^{G12D}-driven PDAC initiation (Navas *et al*, 2012). Kras signalling is tissue specific, with the important implication that efficacy of a treatment cannot be extrapolated from one Kras-driven tumour type to another. Indeed, it is well known that BRAF-driven melanoma and colon cancer differ markedly in their response to targeted therapies (Chapman *et al*, 2011; Prahallad *et al*, 2012). In line with this view, Kras-driven NSCLC depends on coordinated input from the *Ras* oncogene as well as the insulin-like growth factor 1 receptor (IGF1R), but not EGFR (Molina-Arcas *et al*, 2013). Accordingly, EGFR was predominantly activated in Kras wild-type NSCLC (Molina-Arcas *et al*, 2013). These data underscore the need to define tissue- and context-specific molecular hubs and vulnerabilities to develop effective treatment strategies.

CROSS-SIGNALLING AND SIGNALLING LOOPS

It has recently been recognised that autocrine and paracrine signalling loops are important amplifiers of oncogenic Kras signalling in several tumour types (Ardito *et al*, 2012; Navas *et al*, 2012; Molina-Arcas *et al*, 2013).

As described above, different RTKs are engaged in Kras-initiated carcinogenesis in different tissue types, which might explain tissue specificity and the differential response to targeted therapies. In mouse models of PDAC, activation of EGFR by a Kras-induced autocrine-positive feedback loop is essential for Kras-induced ADM and initiation of PanIN lesions (Ardito *et al*, 2012; Navas *et al*, 2012). Although not understood in detail, this positive feedback loop is believed to intensify the level of Kras signalling up to a threshold necessary for transformation of pancreatic acinar cells (Ardito *et al*, 2012; Navas *et al*, 2012). However, additional genetic aberrations, such as functional loss of p53, frequently observed in human PDAC, lead to the development of PDAC even without EGFR expression in this model (Navas *et al*, 2012). How loss of p53 function uncouples Kras^{G12D} from EGFR input in the pancreas is currently unknown. It is also unclear whether EGFR signalling is important for PanIN progression and

PDAC maintenance. Treatment with the EGFR inhibitor Erlotinib has shown benefit in a subpopulation (<10%) of PDAC patients (Moore *et al*, 2007). This raises the question whether a subset of human PDAC remains addicted to KRAS signalling amplification via EGFR. However, the response to Erlotinib could also be due to inhibition of 'off-target' kinases with a higher affinity for the drug compared with EGFR (Conradt *et al*, 2011).

Negative feedback loops and inhibitory cross-signalling between different KRAS downstream pathways have a role in health and disease. Under physiological conditions, these inhibitory circuits fine-tune the level of signalling in response to growth factors, providing an appropriate response to external stimuli. Interestingly, persistence of negative feedback is found in tumour cells and seems to represent a major selection pressure for mutations in modulators of these feedback programmes (Chandarlapaty, 2012). Furthermore, persistent feedback inhibition provides an explanation of oncogene addiction as hyperdependency on the oncoprotein to sustain a certain strength of pathway output that acts to counter intrinsic inhibition (Pratils *et al*, 2009; Chandarlapaty, 2012). This has important implications for the design of targeted therapeutic strategies. Inhibition of oncoproteins or certain effector pathways might reduce negative feedback and actually increase signal output, as shown recently in BRAF mutant colon cancer (Prahallad *et al*, 2012). Selective inhibition of the BRAF oncoprotein reduced a negative feedback loop blocking EGFR. EGFR signalling was consequently activated allowing continued proliferation of colon cancer cells via EGFR-mediated PI3K/AKT pathway activation (Prahallad *et al*, 2012). This contrasts with BRAF mutated melanoma, where BRAF inhibition efficiently blocks MAPK activation without affecting EGFR activation (Chapman *et al*, 2011; Prahallad *et al*, 2012). The mechanisms leading to tissue-specific EGFR expression and activation have yet to be uncovered.

It has long been known that there is inhibitory crosstalk between the PI3K/AKT and MAPK pathways at the level of AKT and RAF that modulates proliferation in human cancer cells (Zimmermann and Moelling, 1999). This interaction was found to be highly complex depending on several parameters such as cell type, RTK input and time course of RTK activation (Moelling *et al*, 2002). A novel RTK-independent crosstalk between these two pathways was found in an intriguing study by Zmajkovicova *et al* (2013). They showed that Mek1, phosphorylated by Erk at T292, is essential for the activity of a MAGI1/Mek1/PTEN complex that negatively regulates PI3K signalling (Zmajkovicova *et al*, 2013). Therefore, inhibition of the MAPK pathway is likely to interfere with PTEN tumour suppressor function and might lead to PI3K pathway activation. Loss of PTEN function and subsequent activation of the PI3K pathway has been found in PDAC and shown to accelerate tumour formation in mouse models of pancreatic cancer by amplification of PI3K/Akt pathway activation (Ying *et al*, 2011). This leads to increased signalling of the NF κ B survival pathway and pro-tumourigenic changes in the tumour microenvironment (Ying *et al*, 2011). These examples of cross-signalling between KRAS downstream pathways in cancer might provide the basis for understanding primary therapeutic resistance and tissue-specific signalling requirements in different tumour entities driven by identical oncogenes.

KRAS SIGNALLING, INFLAMMATION AND THE TUMOUR MICROENVIRONMENT

The tumour microenvironment has been implicated as a major player in PDAC and could conceivably determine tissue-specific signalling pathways by paracrine activation of specific cytokine receptors or RTKs. A prominent desmoplastic reaction is found in PDAC that distinguishes it from other KRAS-driven tumour

Chronic inflammation caused by repeated and/or sustained pancreatic injury through environmental or genetic factors is known to increase pancreatic cancer risk substantially (Yadav and Lowenfels, 2013). Although incompletely understood, the sustained inflammatory micromilieu contributes to a compromised anti-tumour immune response through the infiltration of immunosuppressive regulatory T cells and myeloid-derived suppressor cells (Pylayeva-Gupta *et al*, 2011; Steele *et al*, 2013). In addition, these inflammatory stimuli activate stellate cells and fibroblasts, causing fibrotic remodelling of pancreatic tissue, which in turn enhances oncogenic Kras signalling (Pylayeva-Gupta *et al*, 2011). Indeed, expression of oncogenic Kras in the adult murine pancreas has been shown to cause neoplastic changes only in the context of pancreatic inflammation (Guerra *et al*, 2007). Thus, oncogenic Kras signalling is enhanced by inflammatory stimuli and also itself drives inflammation and desmoplasia in pancreatic neoplasia.

Nearly 100 years ago, Otto Warburg recognised that altered metabolism is a hallmark of cancer. Tumour cells metabolise about 10 times more glucose than lactate than do normal cells, a phenomenon now known as aerobic glycolysis or the Warburg effect (Warburg, 1956). Oncogenic KRAS drives metabolic reprogramming in tumour cells by increasing aerobic glycolysis (e.g., by increasing expression of glycolytic enzymes, such as Hk1, Hk2, Glut1, Pfkf and Ldha), an effect that might be exploited therapeutically (Ying *et al*, 2012). Pancreatic cancer cells also depend on a particular type of glutamine metabolism that differs substantially from normal cells, again offering a promising therapeutic target (Son *et al*, 2013). The identification of tumour-cell-unique and potentially targetable metabolic pathways is a new and exciting field. How soon this can translate into clinically applicable therapeutic strategies is, however, uncertain. Factors such as context specificity and PDAC heterogeneity will certainly be important.

Direct inhibition of the KRAS oncoprotein in PDAC is another hopeful strategy. So far, all attempts to develop inhibitors of KRAS post-translational modification, such as farnesyl- and geranyl-transferase inhibitors that interfere with membrane association and subcellular localisation, have been unsuccessful in the clinic (Berndt *et al*, 2011). Yet, there are some promising new methods, including small molecule inhibitors that block SOS-mediated nucleotide exchange and thus KRAS activation (Maurer *et al*, 2012; Sun *et al*, 2012), and KRAS^{G12C} inhibitors that allosterically shift the affinity of KRAS to favour GDP over GTP (Ostrem *et al*, 2013). Inhibition of the interaction between KRAS and the prenyl-binding protein PDE δ to suppress oncogenic RAS signalling by altering its localisation to endomembranes has also shown interesting results in human PDAC cells *in vitro* and *in vivo* (Zimmermann *et al*, 2013).

www.bjcancer.com | DOI:10.1038/bjc.2014.215

The identification of synthetic lethal interactions of oncogenic KRAS provides another means of targeting mutationally activated KRAS signalling. Defining such interactions depends on comprehensive screening efforts, as recently shown for the synthetic lethal interaction of BCL-XL with MEK inhibition in KRAS-driven cancers (Corcoran *et al.*, 2013). However, concerns about the robustness of such screens require that the targets identified are validated independently.

CONCLUDING REMARKS

Oncogenic KRAS signalling is the main driving force behind PDAC. The signalling networks engaged by oncogenic KRAS are highly complex and characterised by the activation of several effector pathways. These are interconnected at various levels by cross-signalling and feedback loops (Figure 1). KRAS-driven signalling networks differ between tumour entities, such as PDAC, NSCLC and colon cancer, and most likely between subtypes of each entity. In different contexts KRAS signalling involves input from different upstream signals and engagement of different downstream effector pathways. Dissection and thorough understanding of these diverse signalling requirements is essential for the development of effective sub-entity-specific targeted strategies. These are urgently needed to improve the poor prognosis for patients suffering from KRAS-driven cancer.

ACKNOWLEDGEMENTS

We apologise to all colleagues whose work was not included in this review owing to space constraints. We thank Alexander Kind for his suggestions during the preparation of the manuscript. The authors are supported by Deutsche Forschungsgemeinschaft (DFG SA1374/4-1 to DS; SFB824, TP C9 to GS and DS; DFG SCHN971/3-1 to AS), German Cancer Consortium (DKTK) to DS, Deutsche Krebshilfe (#108985 to DS and #110230 to AS), the Helmholtz Alliance Preclinical Comprehensive Cancer Center (PCCC) to DS and Wilhelm-Sander Stiftung (Antrag Nr. 2012.84.1 to GS).

REFERENCES

- Aichler M, Seiler C, Tost M, Sivek J, Mazur PK, Da Silva-Buttkus P, Bartsch DK, Langer P, Chiblak S, Durr A, Hofler H, Kloppel G, Muller-Decker K, Brielmeier M, Esposito I (2012) Origin of pancreatic ductal adenocarcinoma from atypical flat lesions: a comparative study in transgenic mice and human tissues. *J Pathol* **226**: 723–734.
- Ardito CM, Gruner BM, Takeuchi KK, Lubeseder-Martellato C, Teichmann N, Mazur PK, Delgiorio KE, Carpenter ES, Halbrook CJ, Hall JC, Pal D, Briel T, Herner A, Trajkovic-Arsic M, Sipos B, Liou GY, Storz P, Murray NR, Threadgill DW, Sibilio M, Washington MK, Wilson CL, Schmid RM, Raines EW, Crawford HC, Sivek JT (2012) EGF receptor is required for KRAS-induced pancreatic tumorigenesis. *Cancer Cell* **22**: 304–317.
- Berndt N, Hamilton AD, Sefti SM (2011) Targeting protein prenylation for cancer therapy. *Nat Rev Cancer* **11**: 775–791.
- Biankin AV, Waddell N, Kassahn KS, Gingras MC, Muthuswamy LB, Johns AL, Miller DK, Wilson PJ, Patch AM, Wu J, Chang DK, Cowley MJ, Gardiner BB, Song S, Harliwong I, Idrisoglu S, Nourse C, Nourbakhsh E, Manning S, Wani S, Gongora M, Pajic M, Scarlett CJ, Gill AJ, Pinho AV, Rooman I, Anderson M, Holmes O, Leonard C, Taylor D, Wood S, Xu Q, Nones K, Fink JL, Christ A, Bruxner T, Cloonan N, Kolle G, Newell F, Pinese M, Mead RS, Humphris JL, Kaplan W, Jones MD, Colvin EK, Nagrial AM, Humphrey ES, Chou A, Chin VT, Chantrill LA, Mawson A, Samra JS, Kench JG, Lovell JA, Daly RJ, Merrett ND, Toon C, Epari K, Nguyen NQ, Barbour A, Zeps N, Kakkar N, Zhao F, Wu YQ, Wang M, Muzny DM, Fisher WE, Brunicardi FC, Hodges SE, Reid JG, Drummond J, Chang K, Han Y, Lewis LR, Dinh H, Buhay CJ, Beck T, Timms L, Sam M, Begley K, Brown A, Pai D, Panchal A, Buchner N, De Borja R, Denroche RE, Yung CK, Serra S, Onetto N, Mukhopadhyay D, Tsao MS, Shaw PA, Petersen GM, Gallinger S, Hruban RH, Maitra A, Iacobuzio-Donahue CA, Schulick RD, Wolfgang CL, Morgan RA, Lawlor RT, Capelli P, Corbo V, Scardoni M, Tortora G, Tempero MA, Mann KM, Jenkins NA, Perez-Mancera PA, Adams DJ, Largaespa DA, Wessels LF, Rust AG, Stein LD, Tuveson DA, Copeland NG, Musgrave EA, Scarpa A, Eshleman JR, Hudson TJ, Sutherland RL, Wheeler DA, Pearson JV, McPherson JD, Gibbs RA, Grimmond SM (2012) Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. *Nature* **491**: 399–405.
- Blasco RB, Francoz S, Santamaria D, Canamero M, Dubus P, Charron J, Baccarini M, Barbacid M (2011) c-Raf, but not B-Raf, is essential for development of K-Ras oncogene-driven non-small cell lung carcinoma. *Cancer Cell* **19**: 652–663.
- Castellano E, Downward J (2011) RAS Interaction with PI3K: more than just another effector pathway. *Genes Cancer* **2**: 261–274.
- Castellano E, Sheridan C, Thin MZ, Nye E, Spencer-Dene B, Diefenbacher ME, Moore C, Kumar MS, Murillo MM, Gronroos E, Lassailly F, Stamp G, Downward J (2013) Requirement for interaction of PI3-kinase p110alpha with RAS in lung tumor maintenance. *Cancer Cell* **24**: 617–630.
- Chandrapatya S (2012) Negative feedback and adaptive resistance to the targeted therapy of cancer. *Cancer Discov* **2**: 311–319.
- Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, Dummer R, Garbe C, Testori A, Maio M, Hogg D, Lorigan P, Lebbe C, Jouary T, Schadendorf D, Ribas A, O'Day SJ, Sosman JA, Kirkwood JM, Eggermont AM, Dreno B, Nolop K, Li J, Nelson B, Hou J, Lee RJ, Flaherty KT, McArthur GA (2011) Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* **364**: 2507–2516.
- Collins MA, Bednar F, Zhang Y, Brisset JC, Galban S, Galban CJ, Rakshit S, Flannagan KS, Adsay NV, Pasca di Magliano M (2012a) Oncogenic Kras is required for both the initiation and maintenance of pancreatic cancer in mice. *J Clin Invest* **122**: 639–653.
- Collins MA, Brisset JC, Zhang Y, Bednar F, Pierre J, Heist KA, Galban CJ, Galban S, di Magliano MP (2012b) Metastatic pancreatic cancer is dependent on oncogenic Kras in mice. *PLoS One* **7**: e49707.
- Collisson EA, Sadanandam A, Olson P, Gibb WJ, Truitt M, Gu S, Cooc J, Weinkle J, Kim GE, Jakkula L, Feiler HS, Ko AH, Olshen AB, Danenberg KL, Tempero MA, Spellman PT, Hanahan D, Gray JW (2011) Subtypes of pancreatic ductal adenocarcinoma and their differing responses to therapy. *Nat Med* **17**: 500–503.
- Collisson EA, Trejo CL, Silva JM, Gu S, Korkola JE, Heiser LM, Charles RP, Rabinovich BA, Hann B, Dankort D, Spellman PT, Phillips WA, Gray JW, McMahon M (2012) A central role for RAF->MEK->ERK signaling in the genesis of pancreatic ductal adenocarcinoma. *Cancer Discov* **2**: 685–693.
- Conradt L, Godt K, Schaab C, Tebbe A, Eser S, Diersch S, Michalski CW, Kleeff J, Schnieke A, Schmid RM, Saur D, Schneider G (2011) Disclosure of erlotinib as a multikinase inhibitor in pancreatic ductal adenocarcinoma. *Neoplasia* **13**: 1026–1034.
- Conroy T, Desseigne F, Ychou M, Bouche O, Guimbaud R, Becouarn Y, Adenis A, Raoul JL, Gourgou-Bourgade S, de la Fouchardiere C, Bennouna J, Bachet JB, Khemissa-Akouf F, Pere-Verge D, Delbaldo C, Assenat E, Chauffert B, Michel P, Montoto-Grillot C, Ducreux M (2011) FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N Engl J Med* **364**: 1817–1825.
- Corcoran RB, Cheng KA, Hata AN, Faber AC, Ebi H, Coffee EM, Greninger P, Brown RD, Godfrey JT, Cohoon TJ, Song Y, Lifshits E, Hung KE, Shioda T, Dias-Santagata D, Singh A, Settleman J, Benes CH, Mino-Kenudson M, Wong KK, Engelman JA (2013) Synthetic lethal interaction of combined BCL-XL and MEK inhibition promotes tumor regressions in KRAS mutant cancer models. *Cancer Cell* **23**: 121–128.
- Cowley MJ, Chang DK, Pajic M, Johns AL, Waddell N, Grimmond SM, Biankin AV (2013) Understanding pancreatic cancer genomes. *J Hepatobiliary Pancreat Sci* **20**(6): 549–556.
- Engelman JA, Chen L, Tan X, Crosby K, Guimaraes AR, Upadhyay R, Maira M, McNamara K, Perera SA, Song Y, Chirieac LR, Kaur R, Lightbown A, Simendinger J, Li T, Padera RF, Garcia-Echeverria C, Weissleder R, Mahmood U, Cantley LC, Wong KK (2008) Effective use of PI3K and MEK inhibitors to treat mutant Kras G12D and PIK3CA H1047R murine lung cancers. *Nat Med* **14**: 1351–1356.
- Erkan M, Adler G, Apte MV, Bachem MG, Buchholz M, Detlefsen S, Esposito I, Friess H, Gress TM, Habisch HJ, Hwang RF, Jaster R, Kleeff J, Kloppel G, Kordes C, Logsdon CD, Masamune A, Michalski CW, Oh J,

- Phillips PA, Pinzani M, Reiser-Erkan C, Tsukamoto H, Wilson J (2012) StellaTUM: current consensus and discussion on pancreatic stellate cell research. *Gut* **61**: 172–178.
- Eser S, Reiff N, Messer M, Seidler B, Gottschalk K, Dobler M, Hieber M, Arbeiter A, Klein S, Kong B, Michalski CW, Schlitter AM, Esposito I, Kind AJ, Rad L, Schnieke AE, Baccarini M, Alessi DR, Rad R, Schmid RM, Schneider G, Saur D (2013) Selective requirement of PI3K/PDK1 signaling for Kras oncogene-driven pancreatic cell plasticity and cancer. *Cancer Cell* **23**: 406–420.
- Feldmann G, Mishra A, Hong SM, Bisht S, Strock CJ, Ball DW, Goggins M, Maitra A, Nelkin BD (2010) Inhibiting the cyclin-dependent kinase CDK5 blocks pancreatic cancer formation and progression through the suppression of Ras-Ral signaling. *Cancer Res* **70**: 4460–4469.
- Guerra C, Schuhmacher AJ, Canamero M, Grippo PJ, Verdaguier L, Perez-Gallego L, Dubus P, Sandgren EP, Barbacid M (2007) Chronic pancreatitis is essential for induction of pancreatic ductal adenocarcinoma by K-Ras oncogenes in adult mice. *Cancer Cell* **11**: 291–302.
- Heid I, Lubeseder-Martellato C, Sipos B, Mazur PK, Lesina M, Schmid RM, Siveke JT (2011) Early requirement of Rac1 in a mouse model of pancreatic cancer. *Gastroenterology* **141**: 719–730, 730 e1–7.
- Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, Ross S, Conrads TP, Veenstra TD, Hitt BA, Kawaguchi Y, Johann D, Liotta LA, Crawford HC, Putt ME, Jacks T, Wright CV, Hruban RH, Lowy AM, Tuveson DA (2003) Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. *Cancer Cell* **4**: 437–450.
- Jones S, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, Mankoo P, Carter H, Kamiyama H, Jimeno A, Hong SM, Fu B, Lin MT, Calhoun ES, Kamiyama M, Walter K, Nikolskaya T, Nikolsky Y, Hartigan J, Smith DR, Hidalgo M, Leach SD, Klein AP, Jaffee EM, Goggins M, Maitra A, Iacobuzio-Donahue C, Eshleman JR, Kern SE, Hruban RH, Karchin R, Papadopoulos N, Parmigiani G, Vogelstein B, Velculescu VE, Kinzler KW (2008) Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *Science* **321**: 1801–1806.
- Kanda M, Matthaei H, Wu J, Hong SM, Yu J, Borges M, Hruban RH, Maitra A, Kinzler K, Vogelstein B, Goggins M (2012) Presence of somatic mutations in most early-stage pancreatic intraepithelial neoplasia. *Gastroenterology* **142**: 730–733, e9.
- Karthe FA, Frese KK, DeNicola GM, Baccarini M, Tuveson DA (2011) C-Raf is required for the initiation of lung cancer by K-Ras(G12D). *Cancer Discov* **1**: 128–136.
- Lim KH, Baines AT, Fiordalisi JJ, Shipitsin M, Feig LA, Cox AD, Der CJ, Counter CM (2005) Activation of RalA is critical for Ras-induced tumorigenesis of human cells. *Cancer Cell* **7**: 533–545.
- Lim KH, O'Hayer K, Adam SJ, Kendall SD, Campbell PM, Der CJ, Counter CM (2006) Divergent roles for RalA and RalB in malignant growth of human pancreatic carcinoma cells. *Curr Biol* **16**: 2385–2394.
- Maurer T, Garrenton LS, Oh A, Pitts K, Anderson DJ, Skelton NJ, Fauber BP, Pan B, Malek S, Stokoe D, Ludlam MJ, Bowman KK, Wu J, Giannetti AM, Starovasnik MA, Mellman I, Jackson PK, Rudolph J, Wang W, Fang G (2012) Small-molecule ligands bind to a distinct pocket in Ras and inhibit SOS-mediated nucleotide exchange activity. *Proc Natl Acad Sci USA* **109**: 5299–5304.
- Moelling K, Schad K, Bosse M, Zimmermann S, Schwenker M (2002) Regulation of Raf-Akt Cross-talk. *J Biol Chem* **277**: 31099–31106.
- Molina-Arcas M, Hancock DC, Sheridan C, Kumar MS, Downward J (2013) Coordinate direct input of both KRAS and IGF1 receptor to activation of PI3 kinase in KRAS-mutant lung cancer. *Cancer Discov* **3**: 548–563.
- Moore MJ, Goldstein D, Hamm J, Figer A, Hecht JR, Gallinger S, Au HJ, Murawa P, Walde D, Wolff RA, Campos D, Lim R, Ding K, Clark G, Voskoglou-Nomikos T, Ptasynski M, Parulekar W (2007) Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol* **25**: 1960–1966.
- Morris JPt, Wang SC, Hebrok M (2010) KRAS, Hedgehog, Wnt and the twisted developmental biology of pancreatic ductal adenocarcinoma. *Nat Rev Cancer* **10**: 683–695.
- Navas C, Hernandez-Porras I, Schuhmacher AJ, Sibilia M, Guerra C, Barbacid M (2012) EGF receptor signaling is essential for k-ras oncogene-driven pancreatic ductal adenocarcinoma. *Cancer Cell* **22**: 318–330.
- Ostrem JM, Peters U, Sos ML, Wells JA, Shokat KM (2013) K-Ras(G12C) inhibitors allosterically control GTP affinity and effector interactions. *Nature* **503**(7477): 548–551.
- Prahallad A, Sun C, Huang S, Di Nicolantonio F, Salazar R, Zecchin D, Beijersbergen RL, Bardelli A, Bernards R (2012) Unresponsiveness of colon cancer to BRAF(V600E) inhibition through feedback activation of EGFR. *Nature* **483**: 100–103.
- Pratilas CA, Taylor BS, Ye Q, Viale A, Sander C, Solit DB, Rosen N (2009) V600E)BRAF is associated with disabled feedback inhibition of RAF-MEK signaling and elevated transcriptional output of the pathway. *Proc Natl Acad Sci USA* **106**: 4519–4524.
- Pylayeva-Gupta Y, Grabocka E, Bar-Sagi D (2011) RAS oncogenes: weaving a tumorigenic web. *Nat Rev Cancer* **11**: 761–774.
- Seidler B, Schmidt A, Mayr U, Nakhai H, Schmid RM, Schneider G, Saur D (2008) A Cre-loxP-based mouse model for conditional somatic gene expression and knockdown in vivo by using avian retroviral vectors. *Proc Natl Acad Sci USA* **105**: 10137–10142.
- Shimizu T, Tolcher AW, Papadopoulos KP, Beeram M, Rasco DW, Smith LS, Gunn S, Smetzer L, Mays TA, Kaiser B, Wick MJ, Alvarez C, Cavazos A, Mangold GL, Patnaik A (2012) The clinical effect of the dual-targeting strategy involving PI3K/AKT/mTOR and RAS/MEK/ERK pathways in patients with advanced cancer. *Clin Cancer Res* **18**: 2316–2325.
- Singh A, Greninger P, Rhodes D, Koopman L, Violette S, Bardeesy N, Settleman J (2009) A gene expression signature associated with 'K-Ras addiction' reveals regulators of EMT and tumor cell survival. *Cancer Cell* **15**: 489–500.
- Son J, Lyssiotis CA, Ying H, Wang X, Hua S, Ligorio M, Perera RM, Ferrone CR, Mullarky E, Shyh-Chang N, Kang Y, Fleming JB, Bardeesy N, Asara JM, Haigis MC, DePinho RA, Cantley LC, Kimmelman AC (2013) Glutamine supports pancreatic cancer growth through a KRAS-regulated metabolic pathway. *Nature* **496**: 101–105.
- Steele CW, Jamieson NB, Evans TR, McKay CJ, Sansom OJ, Morton JP, Carter CR (2013) Exploiting inflammation for therapeutic gain in pancreatic cancer. *Br J Cancer* **108**: 997–1003.
- Sun Q, Burke JP, Phan J, Burns MC, Olejniczak ET, Waterson AG, Lee T, Rossanese OW, Fesik SW (2012) Discovery of small molecules that bind to K-Ras and inhibit Sos-mediated activation. *Angew Chem Int Ed Engl* **51**: 6140–6143.
- Von Hoff DD, Ervin T, Arena FP, Chiorean EG, Infante J, Moore M, Seay T, Tjulandin SA, Ma WW, Saleh MN, Harris M, Reni M, Dowden S, Laheru D, Bahary N, Ramanathan RK, Tabernero J, Hidalgo M, Goldstein D, Van Cutsem E, Wei X, Iglesias J, Renschler MF (2013) Increased survival in pancreatic cancer with nab-paclitaxel plus gemcitabine. *N Engl J Med* **369**: 1691–1703.
- Warburg O (1956) On the origin of cancer cells. *Science* **123**: 309–314.
- Yadav D, Lowenfels AB (2013) The epidemiology of pancreatitis and pancreatic cancer. *Gastroenterology* **144**: 1252–1261.
- Ying H, Elpek KG, Vinjamoori A, Zimmerman SM, Chu GC, Yan H, Fletcher-Sanankone E, Zhang H, Liu Y, Wang W, Ren X, Zheng H, Kimmelman AC, Paik JH, Lim C, Perry SR, Jiang S, Malinn B, Protopopov A, Colla S, Xiao Y, Hezel AF, Bardeesy N, Turley SJ, Wang YA, Chin L, Thayer SP, DePinho RA (2011) PTEN is a major tumor suppressor in pancreatic ductal adenocarcinoma and regulates an NF-kappaB-cytokine network. *Cancer Discov* **1**: 158–169.
- Ying H, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sanankone E, Locasale JW, Son J, Zhang H, Colloff JL, Yan H, Wang W, Chen S, Viale A, Zheng H, Paik JH, Lim C, Guimaraes AR, Martin ES, Chang J, Hezel AF, Perry SR, Hu J, Gan B, Xiao Y, Asara JM, Weissleder R, Wang YA, Chin L, Cantley LC, DePinho RA (2012) Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* **149**: 656–670.
- Zimmermann G, Papke B, Ismail S, Vartak N, Chandra A, Hoffmann M, Hahn SA, Triola G, Wittinghofer A, Bastiaens PI, Waldmann H (2013) Small molecule inhibition of the KRAS-PDEdelta interaction impairs oncogenic KRAS signalling. *Nature* **497**: 638–642.
- Zimmermann S, Moelling K (1999) Phosphorylation and regulation of Raf by Akt (protein kinase B). *Science* **286**: 1741–1744.
- Zmajkovicova K, Jesenberger V, Catalanotti F, Baumgartner C, Reyes G, Baccarini M (2013) MEK1 is required for PTEN membrane recruitment, AKT regulation, and the maintenance of peripheral tolerance. *Mol Cell* **50**: 43–55.

This work is licensed under the Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>