

Keywords: nonsteroidal antiinflammatory drugs; prostate cancer; screening

Prostate cancer risk and nonsteroidal antiinflammatory drug use in the Finnish prostate cancer screening trial

T Veitonmäki^{*1,2,8}, T J Murtola^{1,3,8}, L Määttänen⁴, K Taari^{5,6}, U-H Stenman⁷, T L J Tammela^{1,3} and A Auvinen²

¹Department of Urology, Tampere University Hospital, Teiskontie 35, 33521 Tampere, Finland; ²School of Health Sciences, University of Tampere, Medisiininkatu 3, 33520 Tampere, Finland; ³School of Medicine, University of Tampere, Lääkärintie 1, 33520 Tampere, Finland; ⁴Finnish Cancer Registry, Pieni Roobertinkatu 9, 00130 Helsinki, Finland; ⁵Department of Urology, Helsinki University Hospital, Sairaalakatu 1, 01400 Vantaa, Finland; ⁶School of Medicine, University of Helsinki, Tukholmankatu 8 B, 00014 Helsinki, Finland and ⁷Department of Clinical Chemistry, Helsinki University Hospital, Haartmaninkatu 4, 00290 Helsinki, Finland

Background: The association between nonsteroidal antiinflammatory drugs (NSAIDs) and prostate cancer risk remains controversial. We examined the risk among NSAID users in 78 615 men in the Finnish Prostate Cancer Screening Trial.

Methods: We obtained information on NSAID prescription usage from Finnish nationwide prescription database and on over-the-counter use by a questionnaire. Prostate cancer cases were identified from the Finnish Cancer Registry.

Results: Prostate cancer risk was elevated among current NSAID prescription users irrespective of screening (hazard ratio (HR) = 1.45, confidence interval (95% CI) = 1.33–1.59 and HR = 1.71, 95% CI = 1.58–1.86 in the screening and control arm, respectively), but not for previous use of NSAIDs. The risk increase was similar among coxib and acetaminophen current users, and stronger for metastatic prostate cancer (HR = 2.41, 95% CI = 1.59–3.67 and HR = 3.44, 95% CI = 2.60–4.55 in the screening and control arm, respectively). Previous use of NSAIDs, aspirin use and over-the-counter NSAID usage were not associated with prostate cancer.

Conclusions: Differing association for current and previous use suggests that the risk increase is unlikely to be directly caused by the medication, but may be due to the conditions indicating NSAID prescription usage, such as symptoms of undiagnosed prostate cancer. To reduce inconsistency between the study outcomes, future epidemiological studies on NSAID use and prostate cancer risk should assess the indications for NSAID usage.

Prostate cancer is the most common malignancy and the second most common cause of cancer death among Western men (Ferlay *et al*, 2010; Siegel *et al*, 2011). Age, African-American ethnicity and family history of prostate cancer are the established risk factors (Steinberg *et al*, 1990; Crawford, 2009), while other aetiological factors remain debatable. The pathogenesis of prostate cancer involves environmental factors as the risk increases with the adoption of a 'westernised'

lifestyle. One such potential environmental factor might be the development of chronic inflammation in the prostate (De Marzo *et al*, 2007). Several lines of research have pointed to a role of inflammation in prostatic carcinogenesis and tumour progression (De Nunzio *et al*, 2011; Sfanos and De Marzo, 2012). Chronic inflammatory conditions are associated with many neoplasms such as colon, hepatic and bladder carcinoma (Coussens and Werb, 2002).

*Correspondence: Dr T Veitonmäki; E-mail: thea.veitonmaki@uta.fi

⁸These authors contributed equally to this work.

Received 16 March 2014; revised 8 June 2014; accepted 15 June 2014; published online 10 July 2014

© 2014 Cancer Research UK. All rights reserved 0007–0920/14

COX-2 is an inducible enzyme that facilitates inflammation by catalysing the conversion of arachidonic acid to prostaglandins (Hussain *et al*, 2003). COX-2 is overexpressed in prostate carcinoma (Gupta *et al*, 2000) and might have an important role in the proliferation of cancer cells (Yoshimura *et al*, 2000). COX-2 overexpression correlates with prostate cancer stage and mortality (Richardson *et al*, 2010; Shao *et al*, 2012). Nonsteroidal antiinflammatory drugs (NSAIDs), which inhibit the COX-2 pathway, could also reduce prostate cancer risk. NSAIDs might work also through a COX-2-independent pathway suppressing oncogenic β -catenin signalling, inhibiting angiogenesis, promoting tumour cell invasion and inducing apoptosis (Gurpinar *et al*, 2014). Aspirin and NSAID use has been associated with a decreased risk of several other cancers such as colorectal, oesophageal, gastric, breast and bladder cancer (Khuder and Mutgi, 2001; Corley *et al*, 2003; Rostom *et al*, 2007; Cuzick *et al*, 2009; Tian *et al*, 2010; Daugherty *et al*, 2011).

The association between NSAID use and prostate cancer risk remains controversial. Epidemiological evidence suggests that prostate cancer risk is reduced by 10% in regular aspirin users (Bosetti *et al*, 2012; Veitonmäki *et al*, 2013). Also other NSAIDs may have a protective association against prostate cancer (Mahmud *et al*, 2010). However, a clear relationship between risk reduction and frequency, dose or duration of use has not been shown (Mahmud *et al*, 2010; Bosetti *et al*, 2012; Veitonmäki *et al*, 2013). In our previous study, we examined the prostate cancer risk for prescription NSAID users in a nationwide population-based case-control study with a large and partly overlapping study population, but without information on screening (Veitonmäki *et al*, 2013). Compared with the nonusers, any NSAID use was associated with an elevated overall prostate cancer risk.

In this study, we examined the risk between NSAID and acetaminophen use and prostate cancer incidence in men attending the Finnish Prostate Cancer Screening Trial during 1996–2009.

MATERIALS AND METHODS

The Finnish Prostate Cancer Screening Trial is the largest component of the European Randomized Study of Prostate Cancer Screening (de Koning *et al*, 2002). In 1996–1999, all men aged 55, 59, 63 or 67 years and residing in the metropolitan areas of Helsinki and Tampere (a total of 80 484 men) were annually identified from the population register of Finland. After exclusion of prevalent prostate cancer cases, 80 144 men were randomly assigned either for PSA-based screening (31 866 men, the screening arm) or to be followed through national registries (48 278 men, the control arm). Men in the screening arm were recruited with mailed invitations to undergo a prostate-specific antigen (PSA) screening test at four-year intervals until the age of 71 years. Men with serum PSA greater than or equal to 4 ng ml^{-1} were referred to a urological clinic for diagnostic examinations, including digital rectal examination (DRE), transrectal ultrasound and prostate biopsy. Men with a PSA level of $3.0\text{--}3.9 \text{ ng ml}^{-1}$ were referred to an additional test, which was DRE during 1996–1998 and since 1999 a free/total PSA (F/T PSA) ratio with a cutoff point of 16%. Men with any of the above findings were considered screen-positive.

The first screening round was carried out during 1996–1999, the second 2000–2003 and the third in 2004–2007. Men who had emigrated from the study area or were diagnosed with prostate cancer were not re-invited. Information on vital status and emigration was obtained from the Population Register Centre. For information on prevalent prostate cancer cases at randomisation, cases diagnosed in the control arm and between the screening rounds, the study cohort was annually linked to the Finnish Cancer

Registry (Kilpeläinen *et al*, 2013). Information on cancer incidence, medication use and vital status was available until the end of 2009.

Finnish Cancer Registry is a population-based, nationwide register that collects data through mandatory notifications of all cancer diagnoses made by the Finnish health care units. It covers more than 99% of all cancer cases in Finland (Teppo *et al*, 1994). The register information includes the primary site of cancer, histology, date and method of diagnosis. Serum PSA values and tumour Gleason scores were obtained from the trial database.

Information on physician-prescribed antiinflammatory medication purchases during 1995–2009 was obtained from the comprehensive nationwide prescription database of the Social Insurance Institution (SII) of Finland. The SII is a governmental agency operating under the ministry of health and provides reimbursements for the cost of medicines prescribed by a physician with the exception of hospital inpatients. The reimbursement is available for all Finnish residents for each purchase of a physician-prescribed drug that has been approved as reimbursable by the SII. The prescription database records the amount, dose and date for each reimbursed purchase. Most prescription drugs in clinical use in Finland have been approved as reimbursable, thus recorded by the database. Purchases of prescription-free drugs are not recorded (Martikainen and Rajaniemi, 2002). The prescription NSAIDs available in Finland are shown in Supplementary Table 1. The medication data was available for 78 615 men (98.1% of the screening trial population). The study flow chart is shown in Figure 1.

To obtain information on over-the-counter usage, the participants of the third screening (years 2004–2007) round were sent a questionnaire on over-the-counter NSAID use with 92.6 % response rate of the screened men. The survey included questions about frequency, dose and duration of over-the-counter usage during this time since 1990 to the present. The survey included questions separately for ibuprofen, dexibuprofen, ketoprofen, aspirin and acetaminophen use.

Information on the comorbidities during hospitalisations in the study population was gained from the Care Registers for Social Welfare and Health Care (HILMO) by the national institutes for Health and Welfare. The registry gathers the information of any hospitalisation in Finnish health care units and information of the diagnoses are registered as ICD-10 codes.

The ethics committee of the Pirkanmaa health care district, Finland reviewed the study protocol (tracking numbers R03209 and R09159).

Statistical analysis. Cox proportional hazards regression was used to estimate hazard ratios (HR) and 95% confidence intervals (CIs) for prostate cancer by medication usage. Nonusers of any NSAIDs (analyses on overall NSAID use) or nonusers of the drug/drug group being analysed (analyses on aspirin, coxibs and acetaminophen) were used as a reference group. Each man in the study population contributed person-time from the date of the randomisation until the date of prostate cancer diagnosis, emigration, death or end of study (31 December 2009), whichever came first.

Cox regression model was adjusted for age, family history of prostate cancer, simultaneous use of other medications previously linked with prostate cancer risk (drugs used for benign prostatic hyperplasia, diabetes, hypercholesterolaemia and hypertension) and number of screening rounds attended. All, with the exception of family history, were statistically significant predictors of the outcome and included in the model. Analyses of the full trial cohort (both study arms together) were additionally adjusted for the trial arm. Analyses on aspirin, coxib or acetaminophen users were also adjusted for usage of other NSAIDs. The proportionality assumption was tested for time-independent variables by adding an interaction term of the exposure indicator with follow-up time into




Figure 1. Flow diagram of the study population.

the Cox regression model. In each case the interaction term was not statistically significant, thus supporting the assumption.

The time axis was years since randomisation. Age and NSAID use were included in the Cox regression model as time-dependent variables, with status updated annually. Men were considered as NSAID nonusers until the year of first purchase. Years with medication purchases were considered as exposed regardless of the dose (current users). Men who stopped using NSAIDs were categorised separately as previous users. No restrictions on frequency of changes in exposure status were imposed, and the exposure status changed as often as the drug purchases indicated.

Cumulative amount (daily doses), duration (years) and intensity (doses/years of usage) of medication use were analysed as time-dependent continuous variables. The defined daily doses (DDDs) recommended by the World Health Organization (WHO ATC/DDD index database) were used to quantify and standardise the amount of NSAID usage. For each year, the total milligram amount for each drug was calculated based on all purchases reimbursed that year. Yearly usage was divided by the DDD amount. The total cumulative numbers of DDDs and years of usage were obtained by adding together yearly DDDs or years with NSAID purchases from the entire follow-up. The intensity of NSAID use was calculated by dividing the yearly number of DDDs purchased with years of usage (number of DDDs/year). Previous NSAID users retained the level of cumulative usage they had reached before discontinuation.

For analyses of dose-dependence, trend tests were calculated by limiting the analysis to medication users only, and adding the quartiles of amount, duration or intensity of usage into the multivariable-adjusted model as a continuous time-dependent variable.

A total cumulative COX-2 inhibition index by NSAID usage was calculated for each person by multiplying the total amount of each NSAID used with the drug-specific COX-1/COX-2 inhibition ratio (Cryer and Feldman, 1998; Kulkarni *et al*, 2000). The analysis was then stratified by cumulative COX-2 inhibition.

To reduce confounding by differing likelihood for NSAID use, we used logistic regression to calculate propensity score estimating each participant's conditional probability (odds ratio) of NSAID usage post randomisation as a function of age, use of other medications previously linked with prostate cancer risk and pre-randomisation NSAID usage. Propensity from each variable was used to form total propensity score as described previously (Rosenbaum and Rubin, 1984). The analysis was then stratified by total propensity score.

To estimate effect modification by usage of medications for other indications, we stratified the analysis by ever use of BPH medication (5 α -reductase inhibitors and alpha-blockers), antidiabetic medication (oral drugs and insulins), cholesterol-lowering medication (statins, fibric acid derivatives, ezetimibe and bile acid-binding resins) and antihypertensive medication (diuretics,

beta-blockers, calcium-channel blockers and drugs targeting the renin-angiotensin system). Effect modification by underlying comorbidities was evaluated using Charlson comorbidity score index calculated based on comorbidities (obtained either from HILMO or by usage of other medications).

Cox regression analyses were performed using IBM SPSS statistics statistical software (version 20, Chicago, IL, USA). Adjusted means for PSA values were estimated using STATA version 12 (StataCorp LP, College Station, TX, USA).

RESULTS

Population characteristics. The median age at entry was 59 years (range 55–67 years) for both NSAID users and nonusers (Table 1). The prevalence of ever use of prescription NSAIDs during follow-up was 77.7% in the screening arm and 74.7% in the control arm, whereas the prevalence was 7.3% and 7.1% for aspirin use and 35.7% and 35.4%, respectively for NSAID usage before randomisation. The self-reported prevalence of over-the-counter use after 1990s was 70.3% for NSAIDs and 48.5% for aspirin.

Overall, the tumour characteristics did not differ by NSAID usage (Table 1). Ever users of prescription NSAIDs had more often used drugs for other indications in both screening trial arms. The proportions of screen-positive men were greater for NSAID users vs nonusers in all screening rounds (Supplementary Table 2).

Overall prostate cancer risk. When compared with NSAID nonusers, the overall prostate cancer risk was elevated among current NSAID prescription users (HR = 1.45, 95% CI = 1.33–1.59 and HR = 1.32, 95% CI = 1.10–1.59 in the screening and the control arm, respectively). Previous use or over-the-counter use was not associated with prostate cancer risk. A similar risk

association was observed also for acetaminophen, but not for aspirin. Both current and previous coxib use was associated with elevated prostate cancer risk (Tables 2a and b).

Prostate cancer risk by tumour stage and grade. Current, but not previous use of prescription NSAIDs, coxibs and acetaminophen associated with an elevated risk of metastatic prostate cancer in both trial arms (Tables 2a and b). Again, aspirin use was not significantly associated with metastatic prostate cancer risk, with the exception of elevated risk among prescription users in the control arm.

The increased prostate cancer risk associated with overall NSAID, coxib and acetaminophen use was not modified by tumour grade; the risk elevation with current NSAID use was similar for low-grade (Gleason 6 or less) and high-grade (Gleason 7–10) tumours in both arms of the screening trial (Tables 2a and b).

Prostate cancer risk by cumulative dose, duration and intensity of usage. Both the overall prostate cancer risk and the risk of metastatic disease were elevated in all quartiles of amount or duration of prescription NSAID and coxib use showing no obvious exposure-effect gradient (Table 3). Intensity of NSAID and coxib use was directly correlated with risk of metastatic prostate cancer after adjustment for the use of other drugs.

The association between aspirin prescription-use and prostate cancer remained statistically nonsignificant in all strata (Table 3).

Sensitivity analyses. The association between NSAID use and prostate cancer risk was not modified by cumulative COX-2 inhibition; the risk was elevated in each quartile of the cumulative COX-2 inhibition, both overall and metastatic disease (Supplementary Table 3).

Stratification by the use of other drug groups or NSAIDs before randomisation, Charlson comorbidity index or propensity for

Table 1. Population characteristics of men attending the Finnish prostate screening trial and the control arm

Characteristics of participants	Screening arm		Control arm	
	NSAID users	NSAID nonusers	NSAID users	NSAID nonusers
Number of participants	23 466	6 728	37 672	10 749
Age at randomisation; median (range)	59 (55–67)	59 (55–67)	59 (55–67)	59 (55–67)
Age at diagnosis; median (range)	67 (55–80)	66 (55–79)	68 (55–80)	67 (55–79)
Mean body mass index	26.9	26.0	—	—
Years of follow-up; median (range)	12 (0–14)	11 (0–14)	12 (0–14)	11 (0–14)
Prostate cancer diagnoses between randomisation and 31 December 2009	2 410	459	3 181	487
Gleason 6 or less; n (%) of diagnoses	1 546 (64.1)	292 (63.6)	1 548 (48.7)	236 (48.5)
Gleason 7–10; n (%)	833 (34.6)	158 (34.4)	1 522 (47.8)	223 (45.8)
Metastatic disease at Dx; n (%) ^a	93 (3.9)	18 (3.9)	252 (7.9)	45 (9.2)
Prevalence of medication use				
Benign prostatic hyperplasia drugs; n (%) ^b	7 931 (33.8)	1 047 (15.6)	12 861 (35.7)	2 160 (20.1)
Antidiabetic drugs; n (%) ^c	5 166 (22.0)	1 002 (14.9)	7 773 (21.6)	1 952 (18.2)
Cholesterol-lowering drugs; n (%) ^d	11 174 (47.6)	1 672 (24.9)	16 830 (46.7)	3 550 (33.0)
Antihypertensive drugs; n (%) ^e	16 963 (72.3)	3 490 (51.9)	25 742 (71.4)	6 736 (62.7)

Abbreviation: NSAID = nonsteroidal antiinflammatory drug.

^aMen with M1 tumour at diagnosis.

^bInformation on physician-prescribed purchases reimbursed by the Social Insurance Institution (SII) of Finland between 1995 and prostate cancer diagnosis, death or common closing date 31 December 2009, whichever comes first. Information obtained from comprehensive national prescription database. Includes 5 α -reductase inhibitors (finasteride and dutasteride) and alpha-blockers (tamsulosin and alfuzosin).

^cIncludes oral antidiabetic drugs (metformin, sulfonylureas, thiazolidinediones, dipeptidyl peptidase-4 inhibitors, meglitinides, α -glucosidase inhibitors and glugacon-like peptide agonists) and insulin.

^dCholesterol-lowering medication (statins, fibric acid derivatives, ezetimibe and bile acid-binding resins).

^eIncludes diuretics, beta-blockers, calcium-channel blockers, angiotensin-converting enzyme inhibitors and angiotensin receptor blockers.

Table 2a. Hazard ratio for prostate cancer by use of nonsteroidal antiinflammatory drugs and by prostate cancer stage and grade in the Finnish prostate cancer screening trial. a) Screening arm

Medication use after randomisation	Overall prostate cancer risk			Risk for localised cancer			Risk for metastatic cancer ^a			Risk for low-grade ^b prostate cancer			Risk for high-grade ^c prostate cancer		
	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	
All NSAIDs															
Prescription use	2354/515	Ref	2226/489	Ref	91/20	Ref	1511/327	Ref	813/178	Ref	Ref	Ref	Ref	Ref	
None															
Current		1.45 (1.33–1.59)		1.41 (1.29–1.55)		2.41 (1.59–3.67)		1.43 (1.28–1.60)		1.56 (1.33–1.82)		1.56 (1.33–1.82)		1.56 (1.33–1.82)	
Previous		0.94 (0.85–1.03)		0.96 (0.87–1.07)		0.50 (0.26–0.92)		1.03 (0.91–1.16)		1.01 (0.85–1.19)		1.01 (0.85–1.19)		1.01 (0.85–1.19)	
Over-the-counter use ^e	376/156	0.98 (0.81–1.18)	358/154	0.95 (0.79–1.15)	7/0	—	201/90	0.89 (0.69–1.14)	174/66	1.09 (0.82–1.45)					
Aspirin															
Prescription use	222/2647	Ref	207/2508	Ref	12/99	Ref	148/1690	Ref	74/917	Ref	Ref	Ref	Ref	Ref	
None															
Current		1.05 (0.82–1.35)		1.08 (0.83–1.39)		1.31 (0.41–4.16)		1.30 (0.97–1.75)		1.03 (0.64–1.65)		1.03 (0.64–1.65)		1.03 (0.64–1.65)	
Previous		0.80 (0.64–1.01)		0.78 (0.62–0.99)		1.57 (0.63–3.93)		0.78 (0.57–1.05)		1.31 (0.92–1.87)		1.31 (0.92–1.87)		1.31 (0.92–1.87)	
Over-the-counter use ^e	272/298	0.95 (0.81–1.12)	258/289	0.94 (0.79–1.11)	7/2	3.2 (0.66–15.67)	147/169	0.90 (0.72–1.13)	124/129	1.01 (0.79–1.29)					
Coxibs															
Prescription use	708/2161	Ref	618/2037	Ref	19/92	Ref	447/1391	Ref	256/735	Ref	Ref	Ref	Ref	Ref	
None															
Current		1.32 (1.10–1.59)		1.23 (1.02–1.49)		3.66 (1.92–6.98)		1.36 (1.09–1.71)		0.90 (0.66–1.23)		0.90 (0.66–1.23)		0.90 (0.66–1.23)	
Previous		1.16 (1.00–1.35)		1.12 (0.96–1.31)		0.94 (0.40–2.21)		1.07 (0.86–1.34)		1.06 (0.85–1.31)		1.06 (0.85–1.31)		1.06 (0.85–1.31)	
Acetaminophen															
Prescription use	801/2068	Ref	748/1967	Ref	35/76	Ref	479/1359	Ref	317/674	Ref	Ref	Ref	Ref	Ref	
None															
Current		1.51 (1.25–1.81)		1.42 (1.16–1.73)		3.24 (1.71–6.14)		1.27 (0.95–1.70)		2.00 (1.56–2.58)		2.00 (1.56–2.58)		2.00 (1.56–2.58)	
Previous		0.98 (0.82–1.18)		0.96 (0.79–1.17)		0.87 (0.32–2.39)		0.60 (0.47–0.77)		1.13 (0.83–1.54)		1.13 (0.83–1.54)		1.13 (0.83–1.54)	

Abbreviations: CI = confidence interval; HR = hazard ratio; NSAID = nonsteroidal antiinflammatory drug.

^aMen with stage T1–4N0–T1M1 tumours.

^bGleason score 6 or less.

^cGleason score 7 or higher.

^dHazard ratio for Cox regression analysis adjusted for age, use of cholesterol-lowering medication, antihypertensive medication, benign prostatic hyperplasia medication, antidiabetic medication and number of screening rounds attended. NSAID usage after randomisation included as time-dependent variable.

^eSelf-reported over-the-counter NSAID and aspirin use among the participants of the third screening round.

Table 2b. Hazard ratio for prostate cancer by use of non-steroidal anti-inflammatory drugs and by prostate cancer stage and grade in the Finnish prostate cancer screening trial. b) Control arm

Overall prostate cancer risk		Risk for localised cancer		Risk for metastatic cancer ^a		Risk for low-grade ^b prostate cancer		Risk for high-grade ^c prostate cancer		
Medication use after randomisation	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d
All NSAIDs										
Prescription use	3107/561	Ref	2762/499	Ref	250/47	Ref	1512/272	Ref	1485/260	Ref
None		1.71 (1.58–1.86)		1.61 (1.48–1.76)		3.44 (2.60–4.55)		1.62 (1.45–1.82)		1.77 (1.57–2.00)
Current		1.02 (0.94–1.12)		1.02 (0.93–1.12)		1.49 (1.07–2.07)		1.03 (0.92–1.17)		1.04 (0.92–1.18)
Previous										
Aspirin										
Prescription use	302/366	Ref	268/2993	Ref	21/276	Ref	139/1645	Ref	145/1600	Ref
None		1.07 (0.86–1.33)		1.00 (0.79–1.27)		2.49 (1.21–5.12)		1.57 (1.12–2.21)		0.99 (0.73–1.33)
Current		0.90 (0.75–1.08)		0.87 (0.71–1.05)		1.26 (0.66–2.41)		1.13 (0.85–1.50)		1.04 (0.81–1.33)
Previous										
Coxibs										
Prescription use	1017/2651	Ref	924/2337	Ref	65/232	Ref	510/1274	Ref	476/1279	Ref
None		1.34 (1.16–1.54)		1.27 (1.09–1.49)		2.58 (1.66–4.00)		1.31 (1.06–1.62)		1.33 (1.09–1.62)
Current		1.04 (0.93–1.17)		1.08 (0.95–1.22)		0.78 (0.44–1.38)		1.17 (0.97–1.40)		1.03 (0.88–1.20)
Previous										
Acetaminophen										
Prescription use	1136/2532	Ref	1012/2249	Ref	87/210	Ref	524/1260	Ref	574/1171	Ref
None		1.65 (1.44–1.90)		1.52 (1.31–1.78)		1.79 (1.12–2.86)		1.60 (1.27–2.01)		1.82 (1.51–2.18)
Current		1.00 (0.86–1.15)		1.02 (0.88–1.19)		2.85 (1.46–5.55)		1.02 (0.81–1.29)		1.24 (1.03–1.50)
Previous										
Overall prostate cancer risk		Risk for metastatic cancer ^a		Risk for low-grade ^b prostate cancer		Risk for high-grade ^c prostate cancer				
No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d	No of cases(exposed/non-exposed)	HR(95% CI) ^d			
All NSAIDs										
Prescription use	2295/1132	Ref	201/85	Ref	1108/602	Ref	1114/470			
None		1.71 (1.58–1.86)		3.44 (2.60–4.55)		1.62 (1.45–1.82)				
Current		1.02 (0.94–1.12)		1.49 (1.07–2.07)		1.03 (0.92–1.17)				
Previous										
Aspirin										
Prescription use	186/3271	Ref	18/268	Ref	82/1628	Ref	95/1489			
None		1.07 (0.86–1.33)		2.49 (1.21–5.12)		1.57 (1.12–2.21)				
Current		0.90 (0.75–1.08)		1.26 (0.66–2.41)		1.13 (0.85–1.50)				
Previous										

Coxibs		275/1201		226/1208		37/193		509/2472	
Prescription use									
None	Ref								
Current	1.33 (1.09–1.62)								
Previous	1.03 (0.88–1.20)								
Acetaminophen		206/1378		139/1571		31/255		354/3073	
Prescription use									
None	Ref								
Current	1.82 (1.51–2.18)								
Previous	1.24 (1.03–1.50)								

Abbreviations: CI = confidence interval; HR = hazard ratio; NSAID = nonsteroidal antiinflammatory drug.
^aMen with stage T1–4N0–1M1 tumours.
^bGleason score 6 or less.
^cGleason score 7 or higher.
^dHazard ratio for Cox regression analysis adjusted for age, use of cholesterol-lowering medication, antihypertensive medication, benign prostatic hyperplasia medication and antidiabetic medication. NSAID usage after randomisation included as time-dependent variable.

NSAID use did not modify the risk association with prostate cancer; current NSAID users had elevated prostate cancer risk, especially for metastatic disease whereas the risk in aspirin users did not differ from nonusers (Table 4).

DISCUSSION

Our study showed an increased HR for prostate cancer among the current prescription users of NSAIDs, coxibs and acetaminophen, but not in men who had been able to discontinue usage, the previous users. Over-the-counter NSAID usage was not associated with prostate cancer risk, either.

Unlike other NSAIDs, aspirin use whether by prescription or over-the-counter was not associated with prostate cancer risk.

Previous results on risk of prostate cancer and other cancer types among NSAID users have been conflicting; whereas some studies have shown an inverse association (Mahmud *et al*, 2010), also null associations (Leitzmann *et al*, 2002; Brasky *et al*, 2010) and increased risks have been reported in both case-control and population-based settings. (Langman *et al*, 2000; Sørensen *et al*, 2003; Murad *et al*, 2011). In our previous study (Veitonmäki *et al*, 2013) in a nationwide population-based setting, we found an elevated risk for prostate cancer in prescription NSAID users. A large case-control study (within the ProtecT trial) reported that prescription NSAID use was associated with prostate cancer (Murad *et al*, 2011), with weaker positive associations for aspirin and acetaminophen. In another case-control study increased risks of pancreatic cancer and prostatic cancer among current prescription NSAID users was observed, but a dose response was found only for pancreatic cancer (Langman *et al*, 2000). An increased risk for cancers of the prostate, kidney and for multiple myeloma was reported among prescription non-aspirin NSAID users in a population-based study (Sørensen *et al*, 2003). Some studies have reported NSAID use to associate to an increased risk for several other cancers (Cho *et al*, 2011; Ross *et al*, 2011; Vinogradova *et al*, 2011; Walter *et al*, 2011). Our findings could help explain some of the previous discrepancy in results, as the association between NSAID use and prostate cancer differed between prescription and over-the-counter use, and most previous studies have used either surveys or prescription databases and have not been able to address this difference. Future epidemiological studies on NSAIDs and prostate cancer risk should consider the two separately.

Our findings do not support a causal association between NSAID use and prostate cancer as the risk elevation was observed only for ongoing prescription use, was strongest for metastatic disease and did not differ by amount or duration of usage. This demonstrates that men starting use of prescription-NSAIDs are at higher risk of prostate cancer, especially metastatic disease already when commencing the use. A comparable risk elevation was observed also for acetaminophen use, with largely similar indications for usage, but a different mechanism of action. Thus the risk elevation is most likely caused by protopathic bias due to treatment of symptoms of undiagnosed prostate cancer, presumably mainly pain due to metastases. This is supported by greatest risk elevation for metastatic cancer being observed with short-term and high-intensity use. Nevertheless, a modest risk elevation by current NSAID and coxib use was observed also for localised and low-grade prostate cancer. This was evident in both study arms of the screening trial, thus not likely explained by more active screening among NSAID users. Again the risk elevation was not dose-dependent suggesting that the indications for NSAID usage are behind this association too.

Common indications for NSAIDs include acute minor infections and chronic inflammatory conditions. It is possible that the systemic inflammation involved in these conditions is involved in

Table 3. Hazard ratio for overall prostate cancer risk by amount and duration of prescription-use of non-steroidal anti-inflammatory drugs among men participating in the Finnish prostate cancer screening trial during 1996–2009

Quantity/ duration of medication use	All NSAIDs		Aspirin		Coxibs	
	HR(95% CI) ^a Overall PCA	HR(95% CI) Metastatic PCA	HR(95% CI) ^a Overall PCA	HR(95% CI) Metastatic PCA	HR(95% CI) ^a Overall PCA	HR(95% CI) Metastatic PCA
Cumulative quantity of medication use^b						
DDD quartiles						
1	1.24 (1.16–1.33)	1.35 (1.01–1.82)	0.92 (0.69–1.22)	1.05 (0.34–3.27)	1.07 (0.93–1.22)	0.83 (0.43–1.56)
2	1.35 (1.26–1.45)	1.78 (1.32–2.40)	0.99 (0.84–1.17)	1.77 (1.03–3.04)	1.21 (1.06–1.38)	1.17 (0.67–2.04)
3	1.29 (1.20–1.40)	2.42 (1.80–3.24)	0.88 (0.70–1.10)	0.89 (0.33–2.40)	1.17 (1.02–1.34)	1.54 (0.94–2.53)
4	1.20 (1.10–1.31)	1.72 (1.22–2.43)	0.93 (0.74–1.16)	1.11 (0.46–2.70)	1.04 (0.90–1.20)	1.42 (0.85–2.38)
P for trend	0.153	0.943	0.296	0.822	0.285	0.943
Duration of medication use^c						
Year quartiles ^d						
1	1.31 (1.22–1.39)	2.08 (1.63–2.65)	0.95 (0.82–1.11)	1.67 (1.01–2.77)	1.13 (1.04–1.23)	1.42 (1.02–1.96)
2	1.26 (1.18–1.36)	1.55 (1.16–2.06)	0.92 (0.74–1.15)	1.10 (0.45–2.66)	1.10 (0.97–1.24)	0.81 (0.45–1.46)
3	1.24 (1.13–1.36)	1.27 (0.84–1.92)	1.07 (0.81–1.39)	1.12 (0.36–3.49)		
4	1.19 (1.07–1.33)	0.97 (0.59–1.60)	0.82 (0.61–1.09)	0.67 (0.17–2.71)		
P for trend	0.893	0.151	0.280	0.628	0.146	0.980
Intensity of medication use (DDDs per year)^e						
Intensity quartile						
1	1.26 (1.17–1.35)	1.25 (0.92–1.70)	0.97 (0.77–1.23)	0.80 (0.25–2.47)	1.05 (0.92–1.20)	0.77 (0.41–1.45)
2	1.25 (1.16–1.36)	1.36 (0.95–1.93)	1.01 (0.84–1.22)	1.65 (0.88–3.12)	1.27 (1.11–1.45)	1.21 (0.68–2.17)
3	1.29 (1.20–1.40)	1.70 (1.24–2.33)	0.81 (0.64–1.04)	1.72 (0.81–3.67)	1.19 (1.03–1.36)	1.25 (0.71–2.20)
4	1.30 (1.20–1.40)	2.79 (2.13–3.65)	0.94 (0.77–1.14)	1.03 (0.46–2.32)	1.02 (0.89–1.17)	1.71 (1.09–2.68)
P for trend	0.870	< 0.001	0.344	0.691	0.631	0.053

Abbreviations: CI = confidence interval; HR = hazard ratio; NSAID = nonsteroidal antiinflammatory drug; PCA = prostate cancer.

^aHazard ratios of prostate cancer from Cox regression analysis adjusted for age, use of cholesterol-lowering medication, antihypertensive medication, benign prostatic hyperplasia medication and antidiabetic medication and the screening trial arm.

^bEstimated by including cumulative daily dose (DDD) quartiles for NSAID use: overall NSAID use 1–34 doses (1st quartile), 35–91 doses (2nd quartile), 92–265 doses (3rd quartile), over 265 doses (4th quartile); aspirin 1–3.5 doses (1st quartile), 3.6–10 doses (2nd quartile), 11–23 doses (3rd quartile), 24 or more doses (4th quartile); coxibs 1–21 doses (1st quartile), 22–49 doses (2nd quartile), 50–130 doses (3rd quartile), over 130 doses (4th quartile).

^cQuartiles for duration of NSAID use before diagnosis: overall NSAID use 1 year (1st quartile), over 1 to 3 years (2nd quartile), over 3 to 5 years (3rd quartile) and over 5 years (4th quartile). Use of aspirin before diagnosis 1 year (1st quartile), 2 years (2nd quartile), 3 years (3rd quartile), 4 or over years (4th quartile). Use of coxibs: 1 year and 2 years or longer.

^dDue to low number of long-term coxib users, duration of coxib use divided into two categories, cut-point set at the median.

^eQuartile cut-points: overall NSAID use: 2–20 DDDs per year (1st quartile), 21–34 DDDs per year (2nd quartile), 35–67 DDDs per year (3rd quartile) and 68 DDDs per year or more (4th quartile); Aspirin use: 0.25–3 DDDs per year (1st quartile), 4–5 DDDs per year (2nd quartile), 6–7 DDDs per year (3rd quartile), 7 DDDs per year or more (4th quartile); Coxib use: 7–20 DDDs per year (1st quartile), 21–35 DDDs per year (2nd quartile), 36–82 DDDs per year (3rd quartile) or 83 DDDs per year or more (4th quartile).

prostate carcinogenesis and prostate cancer progression. Prostatitis might be involved in the complex and multifactorial process of prostate carcinogenesis (MacLennan *et al*, 2006; Wagenlehner *et al*, 2007; Sfanos and De Marzo, 2012). Chronic prostatic inflammation may have a role in the development and progression of chronic prostatic disease, such as BPH and prostate carcinoma (De Nunzio *et al*, 2011). The role of inflammation in prostate cancer can possibly be explained through the cytokine pathway. Interleukin-6 (IL-6) is cytokine that is involved in numerous inflammatory processes (Hirano, 1992) and multiple lines of evidence point to a contributory role in prostate cancer initiation and progression (Twillie *et al*, 1995; Hobisch *et al*, 2000; Smith *et al*, 2001). IL-6 induces the expression of the serine peptidase inhibitor Kazal type 1 (SPINK1) that is overexpressed in prostate cancer (Stenman, 2011) and has been linked to a poorer outcome (Tomlins *et al*, 2008). Another possible biochemical explanation could be the nonsteroidal antiinflammatory drug (NSAID) activated gene-1. NAG-1 has a complex but poorly understood role in several human diseases including cancer. Laboratory studies suggest an antitumorigenic activity of NAG-1 that induces growth arrest or apoptosis, whereas clinical studies demonstrate that NAG-1

expression is upregulated in human prostate cancers, which may also correlate with tumour grade and progression. NAG-1 may have an antitumorigenic role at the early stages of carcinogenesis, but a pro-tumorigenic one during cancer progression (Wang *et al*, 2013). Chronic inflammation and several systematic rheumatic diseases might also be risk factors for malignancy (Coussens and Werb, 2002; Turesson and Matteson, 2013), though the findings regarding prostate cancer are conflicting.

In contrast to other types of NSAIDs, aspirin use was not associated with increased prostate cancer risk. Unlike other NSAIDs, prescription aspirin is mainly used for prevention of cardiovascular and cerebrovascular disorders, and less often as analgetic. The prescription aspirin use in our study population was mostly in combination with dipyridamole for secondary prevention of stroke or transient ischaemic attack (The European Stroke Association (ESO) Executive Committee and the ESO Writing Committee, 2008). Another common indication for aspirin use is cardiovascular disease prevention, for which it is used mainly over-the-counter. These users are more likely to be health-conscious men using small-dose aspirin for preventive reasons, combining aspirin use with healthier lifestyle in general compared with

Table 4. Multivariable-adjusted hazard ratios for overall and metastatic prostate cancer risk for NSAID prescription usage stratified by medication use, Charlson comorbidity index and propensity score

	BPH medication ^a		Cholesterol-lowering medication ^b		Antihypertensive medication ^c		Antidiabetic medication ^d		pre-randomisation NSAID ^e use	
	HR (95% CI) ^f		HR (95% CI)		HR (95% CI)		HR (95% CI)		HR(95%CI)	
Prescription drug use	Yes	No	Yes	No	Yes	No	Yes	No	yes	no
Overall prostate cancer risk										
NSAID										
Never	Ref	Ref	Ref	Ref	Ref	Ref	Ref	Ref	ref	ref
Current	1.37 (1.26–1.50)	1.76 (1.63–1.92)	1.39 (1.27–1.51)	1.80 (1.66–1.95)	1.47 (1.37–1.57)	1.98 (1.77–2.22)	1.38 (1.20–1.58)	1.66 (1.55–1.77)	1.38 (1.24–1.53)	1.78 (1.65–1.92)
Aspirin										
never	Ref	Ref	Ref	Ref	Ref	Ref	Ref	Ref	ref	ref
current	0.92 (0.73–1.16)	1.27 (1.01–1.60)	1.02 (0.84–1.24)	1.33 (0.97–1.82)	1.11 (0.94–1.32)	0.87 (0.48–1.58)	0.87 (0.63–1.20)	1.18(0.98–1.42)	1.09 (0.86–1.38)	1.05 (0.84–1.32)
Risk of metastatic^g prostate cancer										
NSAID										
Current	2.10 (1.44–3.07)	2.96 (2.22–3.96)	2.29 (1.49–3.54)	2.84 (2.16–3.73)	2.57 (1.96–3.36)	2.82 (1.80–4.42)	2.51 (1.51–4.18)	2.67 (2.06–3.46)	2.59 (1.56–4.31)	3.24 (2.47–4.25)
Aspirin										
Current	1.54 (0.67–3.52)	1.34 (0.55–3.28)	1.62 (0.78–3.34)	1.14 (0.37–3.57)	1.56 (0.85–2.87)	—	2.12 (0.92–4.92)	1.05 (0.43–2.56)	1.88 (0.81–4.36)	1.11 (0.45–2.71)
Charlson comorbidity index^h										
	0 HR (95% CI) ^f		1–2 HR (95% CI)		3 or more HR (95% CI)					
	Overall PCa risk	Risk for metastatic PCa ^g	Overall PCa risk	Risk for metastatic PCa ^g	Overall PCa risk	Risk for metastatic PCa ^g	Overall PCa risk	Risk for metastatic PCa ^g		
Prescription drug use	HR(95%CI) ^f	HR(95%CI) ^f	HR(95%CI) ^f	HR(95%CI) ^f	HR(95%CI) ^f	HR(95%CI) ^f	HR(95%CI) ^f	HR(95%CI) ^f		
NSAID										
Current	1.73 (1.60–1.89)	2.29 (2.13–4.03)	1.47 (1.32–1.64)	2.33 (1.56–3.47)	1.31 (1.09–1.57)	2.09 (1.12–3.89)				
ASA										
Current	1.23 (0.89–1.70)	1.43 (0.35–5.81)	1.17 (0.94–1.46)	1.15 (0.46–2.83)	0.85 (0.58–1.24)	1.51 (0.38–4.27)				
Propensity scoreⁱ										
	1st tertile		2nd tertile		3rd tertile					
	Overall PCa risk	Risk for metastatic PCa ^g	Overall PCa risk	Risk for metastatic PCa ^g	Overall pca risk	Risk for metastatic PCa ^g				
Prescription drug use	HR (95% CI) ^f	HR (95% CI) ^f	HR (95% CI) ^f	HR (95% CI) ^f	HR (95% CI) ^f	HR (95% CI) ^f				
NSAID										
Current	2.11 (1.86–2.40)	3.77 (2.58–5.53)	1.61 (1.47–1.77)	3.04 (2.08–4.45)	1.37 (1.23–1.53)	2.29 (1.38–3.80)				
ASA										
Current	1.17 (0.66–2.07)	1.20 (0.17–8.63)	0.99 (0.76–1.29)	0.92 (0.29–2.90)	1.15 (0.92–1.43)	2.05 (0.94–4.46)				

Abbreviations: ASA = aspirin; CI = confidence interval; HR = hazard ratio; NSAID = nonsteroidal antiinflammatory drug; PCa = prostate cancer.

^aBPH medication (5 α -reductase inhibitors and alpha-blockers).

^bAntihypertensive medication (diuretics, beta-blockers, calcium-channel blockers and drugs targeting the renin-angiotensin system).

^cCholesterol-lowering medication (statins, fibric acid derivatives, ezetimibe and bile acid-binding resins).

^dAntidiabetic medication (oral drugs and insulins).

^eNSAID usage status before the screening trial randomisation.

^fHazard ratios from Cox regression analysis adjusted for age, cholesterol-lowering medication, antihypertensive medication, benign prostatic hyperplasia medication and the screening trial arm.

^gMen with stage M1 tumour at diagnosis.

^hCharlson comorbidity index calculated and stratified into groups 0 points, 1–2 points, 3 points and over.

ⁱPropensity for NSAID usage after screening trial randomisation by baseline age, use of other drug groups during the follow-up and NSAID usage before randomisation.

nonusers. Our results from this study do not support prostate cancer preventive effect of aspirin even in long-term use, although the general elevated risk association among NSAID users may have masked a modest risk decreasing effect.

The strength of our study is detailed and objective information on prescription NSAID purchases from the SII prescription database. In addition, we had self-reported information on over-the-counter NSAID usage, although for a shorter period for participants of the third screening round. Thus the results on prescription and over-the-counter usage may not be directly comparable. We had the comprehensive population-based data from the Finnish Cancer registry and the Finnish prostate screening trial with detailed information on prostate cancer grade and stage, and were able to evaluate prostate cancer risk among NSAID users in a setting of systematic screening, which reduces surveillance bias. Further, we could assess the impact of comorbidities and usage of other medication on the results. Thus we were able to minimise the possibility for information and misclassification bias.

Age, ethnic origin and hereditary factors are well-known risk factors for prostate cancer (Steinberg *et al*, 1990; Crawford, 2009). More than 98% of our study population is of Finnish ancestry (Statistics Finland) and the potential for confounding by ethnicity was minimal. Adjusting the analysis for family history did not change our results. We did not have the information on possible lifestyle risk factors such as dietary aspects, alcohol and tobacco consumption or physical activity. This could have caused confounding in either direction, depending on their association with prostate cancer risk. However, their role as prostate cancer risk factors is unclear (Patel and Klein, 2009).

The main limitation of our analysis is missing information on indications of NSAID use, making it difficult to analyse the reasons behind NSAID use. Our study shows that conditions requiring NSAID prescription use may affect prostate cancer risk. Future studies should address the indications for NSAID usage.

The prescription database gives detailed and objective exposure information eliminating recall bias. However, the analysis is based on the assumption that the amount of NSAIDs dispensed is a good approximation of the actual consumption. This may not be true for NSAIDs that are often prescribed to be taken as needed and prescribed drugs may be used only partially or not at all. This might have caused exposure misclassification creating bias towards the null. However, it does not limit our inference of increased prostate cancer risk.

In conclusion, our study showed an elevated risk of prostate cancer in current NSAID prescription users, but not in previous or over-the-counter users. The result is likely to be mostly due to protopathic bias as NSAIDs are used to treat symptoms of undiagnosed metastatic prostate cancer. However, also risk of localised prostate cancer was elevated with current prescription use without dose-dependence, suggesting that conditions indicating such usage could be a prostate cancer risk factor. We found no protecting association for aspirin or NSAID use for prostate cancer. Future studies on NSAIDs and prostate cancer should separate over-the-counter or physician-prescribed use and consider indications for usage to reduce the discrepancy between studies.

ACKNOWLEDGEMENTS

This study has been supported by grants from Astellas, Finnish Cancer Society, Academy of Finland (Grant 132385 and 260931) and the Competitive State Research Financing of Tampere University Hospital (Grant No. 9P053). TV has received a lecture fee from Abbvie and GSK, travel support from Ferring, Sanofi and Astellas and nonrestricted grants from Astellas and Amgen. TJM

has received lecture fee from Novartis, has participated in scientific congresses at the expense of Ferring Pharmaceuticals and Astellas and has received nonrestricted research prize from Astellas. TLJT has served as a board member for Astellas, Amgen and Pfizerand, and has received consulting fees from Orion Pharma, lecture fees from Astellas and Amgen, payment for developing educational presentations from GlaxoSmithKline and travel support from Amgen. KT has received lecture fees from GlaxoSmithKline, honoraria for planning national meetings for GlaxoSmithKline, Astellas and travel support from Astellas, Ipsen, Janssen-Cilag. He has received a nonrestricted grant from Amgen, is a payed consultant/member of advisory board for Astellas, GlaxoSmithKline, Amgen, Pfizer, Sanofi, Janssen-Cilag and has affiliation/employment for Medivation. U-HS holds a patent for the assay of free PSA and has received royalties from the patent. Funding sources have not affected the acquisition, analysis or interpretation of data, or any other aspect of scientific purity.

REFERENCES

- Bosetti C, Rosato V, Gallus S, La Vecchia C (2012) Aspirin and urologic cancer risk: an update. *Nat Rev Urol* **9**(2): 102–110.
- Brasky TM, Velicer CM, Kristal AR, Peters U, Potter JD, White E (2010) Nonsteroidal anti-inflammatory drugs and prostate cancer risk in the VITaminsAnd Lifestyle (VITAL) cohort. *Cancer Epidemiol Biomarkers Prev* **19**(12): 3185–3188.
- Cho E, Curhan G, Hankinson SE, Kantoff P, Atkins MB, Stampfer M, Choueiri TK (2011) Prospective evaluation of analgesic use and risk of renal cell cancer. *Arch Intern Med* **171**(16): 1487–1493.
- Crawford ED (2009) Understanding the epidemiology, natural history, and key pathways involved in prostate cancer. *Urology* **73**(5 Suppl): S4–S10.
- Corley DA, Kerlikowske K, Verma R, Buffler P (2003) Protective association of aspirin/NSAIDs and esophageal cancer: a systematic review and meta-analysis. *Gastroenterology* **124**(1): 47–56.
- Coussens L, Werb Z (2002) Inflammation and cancer. *Nature* **420**(6917): 860–867.
- Cryer B, Feldman M (1998) Cyclooxygenase-1 and cyclooxygenase-2 selectivity of widely used nonsteroidal anti-inflammatory drugs. *Am J Med* **104**: 413–421.
- Cuzick J, Otto F, Baron JA, Brown PH, Burn J, Greenwald P, Jankowski J, La Vecchia C, Meyskens F, Senn HJ, Thun M (2009) Aspirin and nonsteroidal anti-inflammatory drugs for cancer prevention: an international consensus statement. *Lancet Oncol* **10**(5): 501–507.
- Daugherty SE, Pfeiffer RM, Sigurdson AJ, Hayes RB, Leitzmann M, Schatzkin A, Hollenbeck AR, Silverman DT (2011) Nonsteroidal antiinflammatory drugs and bladder cancer: a pooled analysis. *Am J Epidemiol* **173**(7): 721–730.
- de Koning HJ, Auvinen A, Berenguer Sanchez A, Calais da Silva F, Ciatto S, Denis L, Gohagan JK, Hakama M, Hugosson J, Kranse R, Nelen V, Prorok PC, Schröder FH. European Randomized Screening for Prostate Cancer (ERSPC) TrialInternational Randomized Cancer Screening Trials Evaluation Group (2002) Large-scale randomized prostate cancer screening trials: program performances in the European Randomized Screening for Prostate Cancer trial and the Prostate, Lung, Colorectal and Ovary cancer trial. *Int J Cancer* **97**(2): 237–244.
- De Marzo AM, Platz EA, Sutcliffe S, Xu J, Grönberg H, Drake CG, Nakai Y, Isaacs WB, Nelson WG (2007) Inflammation in prostate carcinogenesis. *Nat. Rev. Cancer* **7**: 256–269.
- De Nunzio C, Kramer G, Marberger M, Montironi R, Nelson W, Schröder F, Sciarra A, Tubaro A (2011) The controversial relationship between benign prostatic hyperplasia and prostate cancer: the role of inflammation. *Eur Urol* **60**(1): 106–117.
- Ferlay J., Parkin DM, Steliarova-Foucher E (2010) Estimates of cancer incidence and mortality in Europe in 2008. *Eur J Cancer* **46**(4): 765–781.
- Gupta S, Srivastava M, Ahmad N, Bostwick DG, Mukhtar H (2000) Over-expression of cyclooxygenase-2 in human prostate adenocarcinoma. *Prostate* **42**(1): 73–78.
- Gurpinar E, Grizzle WE, Piazza GA (2014) NSAIDs inhibit tumorigenesis, but how? *Clin Cancer Res* **20**(5): 1104–1113.
- Hirano T (1992) The biology of interleukin-6. *Chem Immunol* **51**: 153–180.

- Hobisch A, Rogatsch H, Hittmair A, Fuchs D, Bartsch Jr G, Klocker H, Bartsch G, Culig Z (2000) Immunohistochemical localization of interleukin-6 and its receptor in benign, premalignant and malignant prostate tissue. *J. Pathol* **191**: 239–244.
- Hussain T, Gupta S, Mukhtar H (2003) Cyclooxygenase-2 and prostate carcinogenesis. *Cancer Lett* **191**(2): 125–135.
- Khuder SA, Mutgi AB (2001) Breast cancer and NSAID use: a meta-analysis. *Br J Cancer* **84**: 1188–1192.
- Kilpeläinen TP, Tammela TL, Malila N, Hakama M, Santti H, Määttänen L, Stenman UH, Kujala P, Auvinen A (2013) Prostate cancer mortality in the Finnish randomized screening trial. *J Natl Cancer Inst* **105**(10): 719–725.
- Kulkarni SK, Jain NK, Singh A (2000) Cyclooxygenase isoenzymes and newer therapeutic potential for selective COX-2 inhibitors. *Methods Find Exp Clin Pharmacol* **22**(5): 291–298.
- Langman MJ, Cheng KK, Gilman EA, Lancashire RJ (2000) Effect of anti-inflammatory drugs on overall risk of common cancer: case-control study in general practice research database. *BMJ* **320**(7250): 1642–1646.
- Leitzmann MF, Stampfer MJ, Ma J, Chan JM, Colditz GA, Willett WC, Giovannucci E (2002) Aspirin use in relation to risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev* **11**(10 Pt 1): 1108–1111.
- MacLennan GT, Eisenberg R, Fleshman RL, Taylor JM, Fu P, Resnick MI, Gupta S (2006) The influence of chronic inflammation in prostatic carcinogenesis: a 5-year followup study. *J Urol* **176**(3): 1012–1016.
- Mahmud SM, Franco EL, Aprikian AG (2010) Use of nonsteroidal anti-inflammatory drugs and prostate cancer risk: a meta-analysis. *Int J Cancer* **127**(7): 1680–1691.
- Martikainen J, Rajaniemi S (2002) *Drug reimbursement systems in EU member states, Iceland and Norway*. The Social Insurance Institution, Finland, Social security and health reports 54: Helsinki, Finland. <http://www.kela.fi/in/internet/english.nsf/NET/100203115310PN>.
- Murad AS, Down L, Davey Smith G, Donovan JL, Athene Lane J, Hamdy FC, Neal DE, Martin RM (2011) Associations of aspirin, non-steroidal anti-inflammatory drug and acetaminophen use with PSA-detected prostate cancer: findings from a large, population-based, case-control study (the ProtecT study). *Int J Cancer* **128**(6): 1442–1448.
- Patel A, Klein E (2009) Risk factors for prostate cancer. *Nat Clin Pract Urol* **6**: 87–95.
- Richardson E, Uglehus RD, Due J, Busch C, Busund LT (2010) COX-2 is overexpressed in primary prostate cancer with metastatic potential and may predict survival. A comparison study between COX-2, TGF-beta, IL-10 and Ki67. *Cancer Epidemiol* **34**(3): 316–322.
- Rosenbaum PR, Rubin DB (1984) Reducing bias in observational studies using subclassification on the propensity score. *J Am Stat Assoc* **79**: 516–524.
- Ross JA, Blair CK, Cerhan JR, Soler JT, Soler JT, Hirsch BA, Roessler MA, Higgins RR, Nguyen PL (2011) Nonsteroidal anti-inflammatory drug and acetaminophen use and risk of adult myeloid leukemia. *CancerEpidemiol Biomarkers Prev* **20**(8): 1741–1750.
- Rostom A, Dubé C, Lewin G, Tsertsvadze A, Barrowman N, Code C, Sampson M, Moher D. U.S. Preventive Services Task Force (2007) Nonsteroidal anti-inflammatory drugs and cyclooxygenase-2 inhibitors for primary prevention of colorectal cancer: a systematic review prepared for the U.S. Preventive Services Task Force. *Ann Intern Med* **146**(5): 376–389.
- Sfanos KS, De Marzo AM (2012) Prostate cancer and inflammation: the evidence. *Histopathology* **60**(1): 199–215.
- Shao N, Feng N, Wang Y, Mi Y, Li T, Hua L (2012) Systematic review and meta-analysis of COX-2 expression and polymorphisms in prostate cancer. *Mol Biol Rep* **39**(12): 10997–11004.
- Siegel R, Ward E, Brawley O, Jemal A (2011) Cancer statistics, 2011: the impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin* **61**(4): 212–236.
- Smith PC, Hobisch A, Lin D-L, Culig Z, Keller ET (2001) Interleukin-6 and prostate cancer progression. *Cytokine Growth Factor Rev* **12**: 33–40.
- Statistics Finland. Finland in figures: Population structure [online] available at <http://www.stat.fi/til/vaerak/index.html> [30 July 2011].
- Steinberg GD, Carter BS, Beatty TH, Childs B, Walsh PC (1990) Family history and the risk of prostate cancer. *Prostate* **17**(4): 337–347.
- Stenman UH (2011) SPINK1: a new therapeutic target in cancer? *Clin Chem* **57**(11): 1474–1475.
- Sørensen HT, Friis S, Nørgård B, Mellemkjaer L, Blot WJ, McLaughlin JK, Ekbohm A, Baron JA (2003) Risk of cancer in a large cohort of nonaspirin NSAID users: a population-based study. *Br J Cancer* **88**: 1687–1692.
- Tian W, Zhao Y, Liu S, Li X (2010) Meta-analysis on the relationship between nonsteroidal anti-inflammatory drug use and gastric cancer. *Eur J Cancer Prev* **19**(4): 288–298.
- Teppo L, Pukkala E, Lehtonen M (1994) Data quality and quality control of a population-based cancer registry. *Acta Oncol* **33**(4): 365–369.
- The European Stroke Association (ESO) Executive Committee and the ESO Writing Committee (2008) Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. *Cerebrovasc Dis* **25**: 457–507.
- Tomlins SA, Rhodes DR, Yu J, Mehra R, Perner S, Demichelis F, Helgeson BE, Laxman B, Morris DS, Cao Q, Cao X, Andrén O, Fall K, Johnson L, Wei JT, Shah RB, Al-Ahmadie H, Eastham JA, Eggeer SE, Fine SW, Hotakainen K, Stenman UH, Tsodikov A, Gerald WL, Lilja H, Reuter VE, Kantoff PW, Scardino PT, Rubin MA, Bjartell AS, Chinnaiyan AM (2008) The role of SPINK1 in ETS rearrangement-negative prostate cancers. *Cancer Cell* **13**(6): 519–528.
- Turesson C, Matteson EL (2013) Malignancy as a comorbidity in rheumatic diseases. *Rheumatology* **52**(1): 5–14.
- Twillie DA, Eisenberger MA, Carducci MA, Hsieh WS, Kim WY, Simons JW (1995) Interleukin-6: a candidate mediator of human prostate cancer morbidity. *Urology* **45**: 542–549.
- Veitonmäki T, Tammela TL, Auvinen A, Murtola T (2013) Use of aspirin, but not other non-steroidal anti-inflammatory drugs is associated with decreased prostate cancer risk at the population level. *Eur J Cancer* **49**(4): 938–945.
- Vinogradova Y, Coupland C, Hippisley-Cox J (2011) Exposure to cyclooxygenase-2 inhibitors and risk of cancer: nested case-control studies. *Br J Cancer* **105**(3): 452–459.
- Wagenlehner FM, Elkahwaji JE, Algaba F, Bjerklund-Johansen T, Naber KG, Hartung R, Weidner W (2007) The role of inflammation and infection in the pathogenesis of prostate carcinoma. *BJU Int* **100**(4): 733–737.
- Walter RB, Milano F, Brasky TM, White E (2011) Long-term use of acetaminophen, aspirin, and other nonsteroidal anti-inflammatory drugs and risk of hematologic malignancies: results from the prospective Vitamins and Lifestyle (VITAL) study. *J Clin Oncol* **29**(17): 2424–2431.
- Wang X, Baek SJ, Eling TE (2013) The diverse roles of nonsteroidal anti-inflammatory drug activated gene (NAG-1/GDF15) in cancer. *Biochem Pharmacol* **85**(5): 597–606.
- World Health Organization. ATC/DDD index database [online] available at http://www.whocc.no/ddd/definition_and_general_considera/ [30 July 2011].
- Yoshimura R, Sano H, Masuda C, Kawamura M, Tsubouchi Y, Chargui J, Yoshimura N, Hla T, Wada S (2000) Expression of cyclooxygenase-2 in prostate carcinoma. *Cancer* **89**(3): 589–596.

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License.

Supplementary Information accompanies this paper on British Journal of Cancer website (<http://www.nature.com/bjc>)