

Keywords: glioma; tricyclic antidepressants; selective serotonin reuptake inhibitors; chemoprevention; pharmacoepidemiology; Denmark

Use of tricyclic antidepressants and risk of glioma: a nationwide case–control study

Anton Pottegård^{*1}, Luis Alberto García Rodríguez², Lotte Rasmussen¹, Per Damkier^{1,3}, Søren Friis⁴ and David Gaist^{5,6}

¹Clinical Pharmacology, Department of Public Health, University of Southern Denmark, Odense, Denmark; ²Centro Español Investigación Farmacoepidemiológica (CEIFE), Madrid, Spain; ³Department of Clinical Chemistry and Pharmacology, Odense University Hospital, Odense, Denmark; ⁴Danish Cancer Society Research Center, Danish Cancer Society, Copenhagen, Denmark; ⁵Department of Neurology, Odense University Hospital, Odense, Denmark and ⁶Department of Clinical Research, Faculty of Health Sciences, University of Southern Denmark, Odense, Denmark

Background: A protective effect of tricyclic antidepressants (TCAs) against gliomas has been suggested by a small number of studies. We investigated this putative association in a nationwide setting.

Methods: Using a case–control design, we identified all patients with histologically verified glioma (cases) in Denmark between 2000 and 2012 and matched these 1:20 to population controls. Conditional logistic regression was used to estimate adjusted odds ratios (ORs) for glioma associated with long-term (≥ 3 years) use of TCAs. Similar analyses were performed for selective serotonin reuptake inhibitors (SSRIs).

Results: We identified 3767 glioma cases and 75 340 population controls. Long-term use of TCAs was inversely associated with risk of glioma (OR 0.72, 95% CI: 0.41–1.25). Long-term SSRI use was not associated with glioma risk (OR 0.93, 95% CI: 0.75–1.16).

Conclusions: Our study indicated that long-term use of TCAs may be associated with a reduced risk of glioma, however, the statistical precision was limited. A similar pattern was not observed for use of SSRIs.

Gliomas constitute about 80% of malignant brain tumours and cause significant morbidity and mortality (Ostrom *et al*, 2014). Although some progress has been made in identifying risk factors for glioma (Chen *et al*, 2011; Goodenberger and Jenkins, 2012; Ostrom *et al*, 2014), the aetiology remains largely unknown. Investigations of the effect of commonly used drugs on glioma risk (Walker *et al*, 2011; Ferris *et al*, 2012; Gaist *et al*, 2013a, b) may improve our understanding of glioma aetiology.

Laboratory studies have indicated that tricyclic antidepressants (TCAs) exert antineoplastic effects (Levkovitz *et al*, 2005; Jahchan *et al*, 2013), although the results are equivocal. Thus far, only one epidemiological study has specifically addressed the risk of glioma among users of antidepressants (Walker *et al*, 2011), reporting a statistically significantly reduced risk of glioma associated with repeat TCA use (odds ratio (OR) 0.59, 95% CI: 0.42–0.81).

To provide further epidemiological data on the potential protective effect of TCA use against gliomas, we conducted a nationwide case–control study of the association between long-term use of TCA and risk of glioma. For comparison, we performed similar analyses for selective serotonin reuptake inhibitors (SSRIs).

MATERIALS AND METHODS

In a nationwide case–control study, we compared the use of TCAs among individuals with incident glioma (cases) with use among cancer-free persons (controls) to estimate ORs for glioma associated with overall or long-term TCA use.

Data sources. We used data from five Danish nationwide registries: the Danish Cancer Registry (Gjerstorff, 2011), the National

*Correspondence: Dr A Pottegård; E-mail: apottegaard@health.sdu.dk

Received 4 December 2015; revised 18 March 2016; accepted 30 March 2016; published online 26 April 2016

© 2016 Cancer Research UK. All rights reserved 0007–0920/16



Prescription Registry (Kildemoes *et al*, 2011), the National Patient Register (Lyng *et al*, 2011), a register in Statistics Denmark holding information on educational level (Baadsgaard and Quitzau, 2011), and the Civil Registration System (Pedersen, 2011). Detailed information on the included registries is provided in the Supplementary Appendix A, while codes for identification of cases, drug use, and other characteristics are provided in the Supplementary Appendix B.

Cases and controls. From the Danish Cancer Registry, we identified all individuals in Denmark with a first-time diagnosis of glioma between 1 January 2000 and 31 December 2012. The date of cancer diagnosis defined the index date. Case diagnoses were restricted to histologically verified gliomas. Exclusion criteria were age < 18 years or > 85 years at diagnosis, previous cancer (except non-melanoma skin cancer), and residency outside Denmark within 10 years prior to index date, ensuring at least 10 years of follow-up and a minimum of 5 years of prescription coverage (see Supplementary Appendix A).

We matched each case to 20 population controls on sex and birth year by risk-set sampling, applying the same selection criteria as for cases. Controls were assigned an index date identical to that of their corresponding case. With the nested case-control design and risk-set sampling of control participants, the calculated ORs provide unbiased estimates of the corresponding incidence rate ratios in the underlying source population.

Exposure definition. Our primary exposure was use of TCA (Anatomical Therapeutic Classification (ATC) code N06AA, see Supplementary Appendix C). We disregarded prescriptions redeemed within 1 year prior to the index date, to reduce the possibility of reverse causation (Jørgensen *et al*). Ever use of TCA was defined as having filled ≥ 1 prescriptions for any TCA prior to index date. Ever use was further classified into current/recent use (≥ 1 prescriptions within 1–1.99 years prior to the index date) and past use (≥ 1 prescriptions > 2 years prior to the index date).

Long-term use of TCA was defined as ≥ 3 years of cumulative duration of use prior to the index date. The duration of each prescription in days was defined as the total number of defined daily doses per prescription.

Main analyses. We used conditional logistic regression to estimate ORs for glioma associated with long-term (≥ 3 years) use of TCA. We further performed analyses according to the cumulative duration of TCA use in four categories: < 1, 1–2, 3–4, and ≥ 5 years. In all analyses, the reference group was never-users of TCA.

Using prescription, patient, and sociodemographic data (see Supplementary Appendices A and B), disregarding the period 1 year prior to the index date, we adjusted for several potential confounders known or suspected to be associated with risk of gliomas, including (i) use (≥ 2 prescriptions) of statins, anti-diabetics, low-dose aspirin, non-aspirin non-steroidal anti-inflammatory drugs, inhibitors of the renin-angiotensin system, antihistamines, anti-asthma drugs, oral contraceptives, and hormonal replacement therapy; (ii) history of diabetes, stroke, allergy, and asthma; and (iii) highest achieved education.

Supplementary analyses. We performed a number of pre-planned supplementary analyses.

First, we repeated the main analyses with SSRI use as the primary exposure. Further, we performed dose-response analyses for all single antidepressant agents (including non-TCA/SSRIs) used by > 100 controls (see Supplementary Appendix C). In these analyses, never use of SSRIs or the individual agent was used as the reference.

Second, we stratified the analyses by type of glioma, that is, glioblastoma multiforme, astrocytoma grade II and III, oligodendroglioma grade II and III, and ‘other types’.

Third, we examined the influence of consistency of TCA use on glioma risk by stratifying long-term TCA use according to the presence or absence of prescription-free periods of ≥ 1 year within 5 years prior to the index date.

Fourth, we examined associations for glioma with TCA use within subgroups defined by sex, age, or previous polyneuropathy. The latter was done to evaluate the potential influence of confounding by indication on the risk estimates.

Finally, we lagged drug and covariate exposure with 2 or 0 years, respectively, instead of the 1-year lag used in the main analyses.

Other. All analyses were performed using Stata Release 14.0 (StataCorp, College Station, TX, USA). The study was approved by the Danish Data Protection Agency. According to Danish law, studies based solely on register data do not require approval from an ethics review board (Thygesen *et al*, 2011).

Table 1. Characteristics of cases and their matched controls

	Cases (n = 3767)	Controls (n = 75 340)
Age, median (IQR)	60 (49–68)	60 (49–68)
Sex		
Male	2217 (58.9%)	44 340 (58.9%)
Female	1550 (41.1%)	31 000 (41.1%)
Glioma subtype		
Glioblastoma multiforme	2353 (62.5%)	NA
Astrocytoma grade II and III	594 (15.8%)	NA
Oligodendroglioma grade II and III	476 (12.6%)	NA
Others	344 (9.1%)	NA
Use of TCA^a		
Never use	3630 (96.4%)	72 152 (95.8%)
Ever use	137 (3.6%)	3188 (4.2%)
Current/recent use	36 (0.96%)	973 (1.3%)
Past use	101 (2.7%)	2215 (2.9%)
Long-term use (≥ 3 years)	13 (0.35%)	373 (0.50%)
Continuous long-term use	11 (0.29%)	292 (0.39%)
Use of SSRI^a		
Never use	3336 (88.6%)	66 118 (87.8%)
Ever use	431 (11.4%)	9222 (12.2%)
Current/recent use	207 (5.5%)	4043 (5.4%)
Past use	224 (5.9%)	5179 (6.9%)
Long-term use (≥ 3 years)	89 (2.4%)	1932 (2.6%)
Continuous long-term use	75 (2.0%)	1534 (2.0%)
Drugs		
Statins	513 (13.6%)	11 290 (15.0%)
Anti-diabetics	294 (7.8%)	6971 (9.3%)
Low-dose aspirin	513 (13.6%)	11 856 (15.7%)
NSAIDs	1735 (46.1%)	35 533 (47.2%)
Inhibitors of the renin-angiotensin system ^b	682 (18.1%)	14 442 (19.2%)
Antihistamines	491 (13.0%)	10 097 (13.4%)
Anti-asthma drugs	556 (14.8%)	12 044 (16.0%)
Oral contraceptives	449 (11.9%)	9013 (12.0%)
Hormone replacement therapy	514 (13.6%)	10 984 (14.6%)
Diagnoses		
Diabetes	118 (3.1%)	3056 (4.1%)
Stroke	65 (1.7%)	1452 (1.9%)
Allergy or asthma	122 (3.2%)	2827 (3.8%)
Highest achieved education		
Short (7–10 years)	1302 (34.6%)	25 886 (34.4%)
Medium (11–13 years)	1539 (40.9%)	30 365 (40.3%)
Long (> 13 years)	813 (21.6%)	16 001 (21.2%)
Missing	113 (3.0%)	3088 (4.1%)
Abbreviations: IQR = interquartile range; NA, not applicable; NSAID = non-steroidal anti-inflammatory drugs; SSRI = selective serotonin reuptake inhibitor; TCA = tricyclic antidepressant.		
^a > 1 year prior to the index date.		
^b Angiotensin-converting enzyme inhibitors and angiotensin-receptor blockers.		

RESULTS

We identified 4522 incident cases of gliomas during 2000–2012. Exclusions left 3767 eligible cases that were matched to 75 340 controls (Table 1; and flow chart in the Supplementary Results I).

Long-term (≥ 3 years) use of TCA was associated with a statistically non-significant reduced risk of glioma with an adjusted OR of 0.72 (95% CI: 0.41–1.25). There was a tendency towards stronger inverse associations with glioma risk with longer duration of TCA use, although these findings had limited statistical precision (Table 2) and a test for trend did not achieve statistical significance. The corresponding analyses for SSRIs revealed ORs close to unity in almost all strata (Table 2). No clear patterns were seen for individual antidepressants (Supplementary Results II), however, numbers were small.

Subgroup analyses showed some variations, however, the statistical precision was limited. Reduced ORs were observed when the analyses were restricted to men, ages between 65–85 years, or glioblastoma multiforme (OR 0.56, 95% CI: 0.26–1.19; Table 3). The corresponding subgroup analyses for SSRIs showed no material variation in risk estimates, except among subjects aged 65–85 years and with glioma subtype (Supplementary Results III). Varying the lag-time did not affect the OR estimates (data not shown).

As we found the lowest ORs for glioma with TCA use close to the index date, we performed an exploratory *post hoc* analysis examining the association between ‘current use’ of TCA, defined as use within the last year prior to index date, and glioma risk finding an OR of 1.06 (95% CI: 0.80–1.40). In addition, we performed an analysis of long-term use among such current users, which returned an OR of 0.81 (95% CI: 0.44–1.48).

Table 2. Association between the use of TCA and SSRI and risk of glioma according to various exposure definitions

	Cases	Controls	Crude OR ^a	Adjusted OR ^b
TCA				
Never use	3630	72 152	1.00 (ref)	1.00 (ref)
Ever use	137	3188	0.85 (0.72–1.02)	0.89 (0.75–1.06)
Current/recent use	36	973	0.73 (0.52–1.03)	0.76 (0.55–1.07)
Past use	101	2215	0.91 (0.74–1.11)	0.95 (0.77–1.16)
Long-term use	13	373	0.69 (0.40–1.21)	0.72 (0.41–1.25)
Continuous long-term use	11	292	0.75 (0.41–1.36)	0.76 (0.42–1.40)
Cumulative duration				
< 1 year	109	2334	0.93 (0.76–1.13)	0.97 (0.80–1.18)
1–2 years	15	481	0.63 (0.37–1.05)	0.65 (0.39–1.09)
3–4 years	5	174	0.57 (0.23–1.39)	0.59 (0.24–1.44)
> 5 years	8	199	0.81 (0.40–1.64)	0.83 (0.41–1.68)
SSRI				
Never use	3336	66 118	1.00 (ref.)	1.00 (ref.)
Ever use	431	9222	0.92 (0.83–1.03)	0.95 (0.86–1.06)
Current/recent use	207	4043	1.01 (0.87–1.17)	1.04 (0.90–1.20)
Past use	224	5179	0.86 (0.75–0.99)	0.88 (0.76–1.01)
Long-term use	89	1932	0.90 (0.72–1.12)	0.93 (0.75–1.16)
Continuous long-term use	75	1534	0.95 (0.75–1.20)	0.98 (0.77–1.24)
Cumulative duration				
< 1 year	242	5266	0.91 (0.80–1.04)	0.93 (0.81–1.07)
1–2 years	100	2024	0.98 (0.80–1.20)	1.00 (0.81–1.23)
3–4 years	43	905	0.93 (0.68–1.27)	0.96 (0.70–1.30)
> 5 years	46	1027	0.86 (0.64–1.17)	0.89 (0.66–1.20)

Abbreviations: OR = odds ratio; SSRI = selective serotonin reuptake inhibitor; TCA, tricyclic antidepressant. Long-term use is defined as ≥ 3 years of cumulative use. In all analyses, prescriptions within the year prior to the index date were disregarded.

^aAdjusted for age and gender (by design; risk-set matching and conditional analysis).

^bFurther adjusted for (i) use (≥ 2 prescriptions) of statins, anti-diabetics, low-dose aspirin, non-aspirin NSAIDs, inhibitors of the renin-angiotensin system, antihistamines, anti-asthma drugs, oral contraceptives, and hormonal replacement therapy; (ii) previous diagnoses of diabetes, stroke, allergy, and asthma; and (iii) highest achieved education.

Table 3. Associations between long-term use of TCA (≥ 3 years) and risk of glioma, specified by subgroups

	Cases	Controls	Crude OR ^a	Adjusted OR ^b
Sex				
Men	4/2164	149/42 963	0.53 (0.20–1.43)	0.54 (0.20–1.46)
Women	9/1466	224/29 189	0.81 (0.41–1.57)	0.84 (0.43–1.64)
Age				
18–49 years	—/928	34/18 478	—	—
50–64 years	6/1443	146/28 716	0.83 (0.37–1.87)	0.83 (0.37–1.89)
65–85 years	5/1259	193/24 958	0.51 (0.21–1.25)	0.54 (0.22–1.32)
Histological subgroup				
Glioblastoma multiforme	7/2262	255/44 905	0.54 (0.26–1.15)	0.56 (0.26–1.19)
Comorbidity				
No polyneuropathy	13/3614	363/71 873	0.72 (0.41–1.25)	0.74 (0.42–1.28)

Abbreviations: OR = odds ratio. In all analyses prescriptions within the year prior to the index date were disregarded.

^aAdjusted for age and gender (by design; risk-set matching and conditional analysis).

^bFurther adjusted for (i) use (≥ 2 prescriptions) of statins, anti-diabetics, low-dose aspirin, non-aspirin NSAIDs, inhibitors of the renin-angiotensin system, antihistamines, anti-asthma drugs, oral contraceptives, and hormonal replacement therapy; (ii) previous diagnoses of diabetes, stroke, allergy, and asthma; and (iii) highest achieved education.

DISCUSSION

In this nationwide study, we found an inverse, albeit statistically non-significant, association between TCA use and risk of glioma. However, the statistical precision of our analyses was generally limited.

The main strength of our study is the use of high-quality nationwide registries, whereas the main limitation was the low number of exposed cases limiting the statistical precision and complicating interpretation of subgroup analyses.

The only previous observational study of TCA use and glioma risk reported a statistically significant reduction in glioma risk associated with the use of TCAs with an OR of 0.59 (95% CI, 0.42–0.81) while also showing a statistically significant dose-dependency (Walker *et al*, 2011). The authors also reported a null association between use of SSRI and glioma risk (OR 0.96; 95% CI: 0.61–1.53) (Walker *et al*, 2011). Although with less statistical precision, our results are largely compatible with those of the study by Walker *et al* (2011).

Biologically plausible explanations supporting these findings appear insufficiently accounted for. Some *in vitro* studies suggest that antidepressants inhibit the growth of human glioma cell lines (Ishiyuchi *et al*, 2007; Liu *et al*, 2015). The primary mechanism appears to be related to the overexpression of AMPAR receptors in glioma tissues with TCAs acting as ligands resulting in apoptosis. Autophagy has been suggested as another mechanism of action (Jeon *et al*, 2011). Study conditions have been heterogeneous, but overall, fluoxetine, paroxetine, sertraline, and clomipramine have demonstrated an enhancing effect on apoptosis in human glioma cell lines (Levkovitz *et al*, 2005; Tzadok *et al*, 2010; Liu *et al*, 2015). No such effect could be demonstrated for imipramine or mianserine (Levkovitz *et al*, 2005). Other antidepressants have not been subject to comparable studies. As such, we would hypothesise that signals, if any, be present for fluoxetine, paroxetine, sertraline, and clomipramine, but not for imipramine or mianserine. No such pattern was observed (see Supplementary Results II), however, the statistical precision was low in the analyses of individual agents.

Although our findings support the hypothesised protective effect of TCAs on the development of glioma, further substantiation from association studies and elucidation of the biological rationale is required.

ACKNOWLEDGEMENTS

Morten Olesen is acknowledged for help with data management.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Baadsgaard M, Quitzau J (2011) Danish registers on personal income and transfer payments. *Scand J Public Health* **39**: 103–105.

- Chen C, Xu T, Chen J, Zhou J, Yan Y, Lu Y, Wu S (2011) Allergy and risk of glioma: a meta-analysis. *Eur J Neurol Off J Eur Fed Neurol Soc* **18**: 387–395.
- Ferris JS, McCoy L, Neugut AI, Wrensch M, Lai R (2012) HMG CoA reductase inhibitors, NSAIDs and risk of glioma. *Int J Cancer J Int Cancer* **131**: E1031–E1037.
- Gaist D, Andersen L, Hallas J, Sørensen HT, Schrøder HD, Friis S (2013a) Use of statins and risk of glioma: a nationwide case-control study in Denmark. *Br J Cancer* **108**: 715–720.
- Gaist D, García-Rodríguez LA, Sørensen HT, Hallas J, Friis S (2013b) Use of low-dose aspirin and non-aspirin nonsteroidal anti-inflammatory drugs and risk of glioma: a case-control study. *Br J Cancer* **108**: 1189–1194.
- Gjerstorff ML (2011) The Danish Cancer Registry. *Scand J Public Health* **39**: 42–45.
- Goodenberger ML, Jenkins RB (2012) Genetics of adult glioma. *Cancer Genet* **205**: 613–621.
- Ishiyuchi S, Yoshida Y, Sugawara K, Aihara M, Ohtani T, Watanabe T, Saito N, Tsuzuki K, Okado H, Miwa A, Nakazato Y, Ozawa S (2007) Ca²⁺-permeable AMPA receptors regulate growth of human glioblastoma via Akt activation. *J Neurosci* **27**: 7987–8001.
- Jahchan NS, Dudley JT, Mazur PK, Flores N, Yang D, Palmerton A, Zmoos A-F, Vaka D, Tran KQT, Zhou M, Krasinska K, Riess JW, Neal JW, Khatri P, Park KS, Butte AJ, Sage J (2013) A drug repositioning approach identifies tricyclic antidepressants as inhibitors of small cell lung cancer and other neuroendocrine tumors. *Cancer Discov* **3**: 1364–1377.
- Jeon S-H, Kim SH, Kim Y, Kim YS, Lim Y, Lee YH, Shin SY (2011) The tricyclic antidepressant imipramine induces autophagic cell death in U-87MG glioma cells. *Biochem Biophys Res Commun* **413**: 311–317.
- Jørgensen T, Herrstedt J, Friis S, Hallas J (2012) Polypharmacy and drug use in elderly Danish cancer patients during 1996 to 2006. *J Geriatr Oncol* **3**: 33–40.
- Kildemoes HW, Sørensen HT, Hallas J (2011) The Danish National Prescription Registry. *Scand J Public Health* **39**: 38–41.
- Levkovitz Y, Gil-Ad I, Zeldich E, Dayag M, Weizman A (2005) Differential induction of apoptosis by antidepressants in glioma and neuroblastoma cell lines: evidence for p-c-Jun, cytochrome c, and caspase-3 involvement. *J Mol Neurosci* **27**: 29–42.
- Liu K-H, Yang S-T, Lin Y-K, Lin J-W, Lee Y-H, Wang J-Y, Hu C-J, Lin E-Y, Chen S-M, Then C-K, Shen S-C (2015) Fluoxetine, an antidepressant, suppresses glioblastoma by evoking AMPAR-mediated calcium-dependent apoptosis. *Oncotarget* **6**: 5088–5101.
- Lynge E, Sandegaard JL, Rebolj M (2011) The Danish National Patient Register. *Scand J Public Health* **39**: 30–33.
- Ostrom QT, Bauchet L, Davis FG, Deltour I, Fisher JL, Langer CE, Pekmezci M, Schwartzbaum JA, Turner MC, Walsh KM, Wrensch MR, Barnholtz-Sloan JS (2014) The epidemiology of glioma in adults: a 'state of the science' review. *Neuro Oncol* **16**: 896–913.
- Pedersen CB (2011) The Danish Civil Registration System. *Scand J Public Health* **39**: 22–25.
- Thygesen LC, Daasnes C, Thaulow I, Brønnum-Hansen H (2011) Introduction to Danish (nationwide) registers on health and social issues: structure, access, legislation, and archiving. *Scand J Public Health* **39**: 12–16.
- Tzadok S, Beery E, Israeli M, Uziel O, Lahav M, Fenig E, Gil-Ad I, Weizman A, Nordenberg J (2010) *In vitro* novel combinations of psychotropics and anti-cancer modalities in U87 human glioblastoma cells. *Int J Oncol* **37**: 1043–1051.
- Walker AJ, Card T, Bates TE, Muir K (2011) Tricyclic antidepressants and the incidence of certain cancers: a study using the GPRD. *Br J Cancer* **104**: 193–197.



This work is licensed under the Creative Commons Attribution-Non-Commercial-Share Alike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Supplementary Information accompanies this paper on British Journal of Cancer website (<http://www.nature.com/bjc>)