Keywords: panitumumab; metastatic colorectal cancer; RAS; KRAS; NRAS; BRAF; amphiregulin; response

Impact of tumour *RAS/BRAF* status in a first-line study of panitumumab + FOLFIRI in patients with metastatic colorectal cancer

Meinolf Karthaus^{*1}, Ralf-Dieter Hofheinz², Laurent Mineur³, Henry Letocha⁴, Richard Greil⁵, Josef Thaler⁶, Eva Fernebro⁷, Kelly S Oliner⁸, Michael Boedigheimer⁹, Brian Twomey⁹, Ying Zhang⁹, Gaston Demonty¹⁰ and Claus-Henning Köhne¹¹

¹Klinikum Neuperlach/Klinikum Harlaching, Oskar-Maria-Graf-Ring 51, D81737 Munich, Germany; ²Universitätsmedizin Mannheim, Theodor-Kutzer Ufer 1-3, 68167 Mannheim, Germany; ³Institut Sainte-Catherine GI and Liver Cancer Unit, 84 000 Avignon, France; ⁴Oncology Clinic, Västmanland's Hospital, 721 89 Västerås, Sweden; ⁵Illrd Medical Department, Paracelsus Medical University Salzburg and CCCIT Salzburg Cancer Research Institute, Müllner Hauptstrasse 45, 5020 Salzburg, Austria; ⁶Klinikum Wels-Grieskirchen, Grieskirchner Straße 42, A-4600 Wels, Austria; ⁷Central Hospital, Strandvägen 8, 35185 Växjö, Sweden; ⁸Formerly of Amgen Inc., 1 Amgen Center Dr MS 30E-2-C, Thousand Oaks, CA 91320, USA; ⁹Amgen Inc., 1 Amgen Center Dr MS 30E-2-C, Thousand Oaks, CA 91320, USA; ¹⁰Amgen GmbH, Dammstrasse 23, 6301 Zug, Switzerland and ¹¹Onkologie Klinikum Oldenburg, Rahel-Straus-Str. 10, 26133 Oldenburg, Germany

Background: To investigate tumour biomarker status and efficacy of first-line panitumumab + FOLFIRI for metastatic colorectal carcinoma (mCRC).

Methods: 154 patients received first-line panitumumab + FOLFIRI every 14 days. Primary end point was objective response rate (ORR). Data were analysed by tumour RAS (KRAS/NRAS) and BRAF status, and baseline amphiregulin (AREG) expression.

Results: Objective responses occurred more frequently in RAS wild type (WT) (59%) vs RAS mutant (MT) (41%) mCRC and in RAS WT/BRAF WT (68%) vs RAS or BRAF MT (37%) disease. Median response duration was longer in RAS WT (13.0 months) vs RAS MT (5.8 months) (hazard ratio (HR): 0.16). Median progression-free survival was longer in RAS WT vs MT (11.2 vs 7.3 months; HR, 0.37) and was also longer in RAS WT/BRAF WT vs RAS or BRAF MT (13.2 vs 6.9 months; HR, 0.25). Incidence of adverse events was similar regardless of RAS/BRAF status, and no new safety signals were noted. Among patients with RAS WT tumours, ORR was 67% with high AREG expression and 38% with low AREG expression.

Conclusions: First-line panitumumab + FOLFIRI was associated with favourable efficacy in patients with RAS WT and RAS WT/BRAF WT vs MT mCRC tumours and was well tolerated.

FOLFIRI (folinic acid, infusional 5-fluorouracil, and irinotecan) is a recommended first- and second-line chemotherapy backbone in the treatment of metastatic colorectal cancer (mCRC) (National Comprehensive Cancer Network, 2015). Following a recent review by the European Medicines Agency (EMA), both of the epidermal growth factor receptor (EGFR) inhibitors – panitumumab and

cetuximab – are now approved in Europe for first-line use in combination with FOLFIRI or FOLFOX (folinic acid, infusional 5-fluorouracil, and oxaliplatin) in patients with RAS wild-type (WT) mCRC.

In the first-line setting, EGFR inhibitors were originally indicated in patients with KRAS WT mCRC in combination with

*Correspondence: Professor M Karthaus; E-mail: meinolf.karthaus@klinikum-muenchen.de

Received 9 May 2016; revised 21 July 2016; accepted 19 September 2016; published online 20 October 2016

© 2016 Cancer Research UK. All rights reserved 0007–0920/16

either FOLFOX (panitumumab or cetuximab) or FOLFIRI (cetuximab). Subsequently, mutations were identified in *KRAS* and *NRAS* in 17% of patients with non-mutated *KRAS* exon 2 in the phase III PRIME trial of panitumumab + FOLFOX vs FOLFOX alone (Douillard *et al*, 2013). As a result, EGFR inhibitor use was refined to include only those patients with RAS WT disease. At the same time, studies also identified *BRAF* as an important negative prognostic – though not predictive – marker for survival in patients with mCRC, regardless of treatment (Phipps *et al*, 2012; Douillard *et al*, 2013; Peeters *et al*, 2013, 2014a, c). Tumour expression of the biomarker amphiregulin (AREG) has also been correlated with survival during anti-EGFR therapy (Jacobs *et al*, 2009; Loupakakis *et al*, 2014).

The improved risk–benefit profile of EGFR inhibitor treatment in patients selected by RAS status was later verified in trials of first- and second-line FOLFOX and FOLFIRI in combination with panitumumab or cetuximab (Douillard *et al*, 2013; Heinemann *et al*, 2014a, b; Venook *et al*, 2014; Bokemeyer *et al*, 2015; Van Cutsem *et al*, 2015). Importantly, patients with RAS mutant (MT) tumours showed no improvement in efficacy with the addition of EGFR inhibitor compared with chemotherapy alone. Indeed, there is some evidence that EGFR inhibitors combined with FOLFOX in such patients are detrimental compared with FOLFOX alone (Douillard *et al*, 2013; Bokemeyer *et al*, 2015), and EGFR inhibitors should therefore not be given to patients with RAS MT tumours.

Panitumumab plus FOLFIRI demonstrated superiority over FOLFIRI alone in a second-line phase II trial that demonstrated improved progression-free survival (PFS) in patients with RAS WT mCRC, although overall survival (OS) was not significantly different in this study (Peeters *et al*, 2014b). The only published study of first-line panitumumab plus FOLFIRI in mCRC is a single-arm trial that indicated the efficacy and predictable safety profile of the combination in *KRAS* WT patients (Köhne *et al*, 2012), although preliminary data have been presented from a phase II study in which panitumumab plus FOLFOX4 or FOLFIRI was evaluated in patients with *KRAS* WT colorectal cancer and liver-limited disease (Abad *et al*, 2014). Here, we present a retrospective analysis of this first-line trial of panitumumab plus FOLFIRI, reporting efficacy and safety data for first-line FOLFIRI plus panitumumab according to tumour RAS/BRAF status and AREG levels in patients with mCRC.

PATIENTS AND METHODS

Study design. This is a retrospective analysis of data from a phase II, single-arm study (NCT 00508404); full details of the study design have been presented previously (Köhne *et al*, 2012). In brief, first-line panitumumab (6 mg kg^{−1}) + FOLFIRI were administered every 14 days until progression, unacceptable toxicity or withdrawal of consent. If FOLFIRI was withdrawn or suspended as a result of toxicity, panitumumab could be continued, and vice versa.

Patients. Patients were adults with histologically or cytologically confirmed, radiologically measurable mCRC and an Eastern Cooperative Oncology Group (ECOG) performance status of 0–2. Patients could be enrolled only if all disease sites were evaluated within 28 days before enrolment, and tissue from the primary or metastatic site was available. Those who had received prior systemic therapy (including anti-EGFR therapy) for mCRC (except adjuvant fluoropyrimidine-based chemotherapy given ≥ 6 months before enrolment) were excluded. Full inclusion and exclusion criteria have been reported previously (Köhne *et al*, 2012).

The study protocol was approved by the relevant independent ethics committees. The study was conducted in accordance with the regulations and guidelines of the International Conference on Harmonisation of Good Clinical Practice. All patients provided signed, informed consent before any study-related procedures were performed.

End points and analyses. The primary end point was the objective response rate (ORR) assessed using modified Response Evaluation Criteria in Solid Tumors (mRECIST v1.0) (Therasse *et al*, 2000). Secondary end points included disease control rate (DCR), duration of response (DoR), depth of response (DpR; defined as the percentage of tumour shrinkage at nadir or progression), PFS, and time to progression (TTP). Resection rates were also evaluated, as was early tumour shrinkage (ETS), defined as the percentage of patients with ≥ 30% or ≥ 20% tumour shrinkage at week 8 (exploratory analysis). Safety was evaluated in terms of the incidence and severity of adverse events (AEs), using the National Cancer Institute Common Toxicity Criteria version 3.0.

Data were analysed descriptively by tumour RAS/BRAF status. Tumour specimens were assayed for mutations in *KRAS* exons 3–4, *NRAS* exons 2–4 and *BRAF* exon 15 by bidirectional Sanger sequencing. Mutations in *KRAS* exon 2 were analysed by CE-marked DxS kit.

Baseline tumour AREG levels were analysed in the RAS WT and MT populations. Total RNA was extracted from formalin-fixed paraffin-embedded tissue samples and AREG expression levels were analysed by qualified reverse transcription quantitative polymerase chain reaction (RT-qPCR) assays (see Supplementary Material for details). A cutoff point for AREG status was prespecified based on analysis of data from an earlier clinical trial (STEPP) (Lacouture *et al*, 2010). Cox proportional hazards (PH) models were used to evaluate AREG expression levels as a continuous covariate. Decision curves were used to estimate the PFS hazard ratio (HR) with increasing levels of baseline AREG expression. A Gaussian Process (GP) model was used to fit the PH model (Joensuu *et al*, 2012) using the GPstuff toolkit in MATLAB (Vanhatalo *et al*, 2013) – details are included in the Supplementary Material.

RESULTS

Ascertainment of tumour *KRAS*/*BRAF* mutation status. In total, 154 patients were enrolled in the study (Köhne *et al*, 2012). One patient withdrew consent and three patients had no DNA and/or tumour tissue available. The RAS mutational analysis therefore included 150 patients, of whom 143 received at least one dose of panitumumab.

Overall, 38% of patients had *KRAS* exon 2 mutations, and 10% had RAS mutations beyond *KRAS* exon 2 (*KRAS* exon 3 and 4 mutations each in 3% of tumours; *NRAS* exon 2 and 3 mutations each in 2% of tumours; no tumour was found to carry an *NRAS* exon 4 mutation). Complete RAS data were available for 143 of 150 patients, of whom 69 (45%) had RAS WT tumours (i.e., WT for exons 2, 3 and 4 of both *KRAS* and *NRAS*). *BRAF* mutations were present in nine patients (6%), all of whom had tumours that were WT for RAS.

Patients. Baseline characteristics were generally well balanced between patients with RAS WT vs RAS MT mCRC, and between patients with RAS WT/*BRAF* WT and RAS or *BRAF* MT status (Table 1). More patients with RAS WT and RAS WT/*BRAF* WT mCRC had liver-limited metastases, whereas more RAS MT and RAS or *BRAF* MT patients had metastases at other sites only. The sum of the longest diameters of measurable target lesions was also slightly larger in patients with RAS WT and RAS WT/*BRAF* WT mCRC compared with the corresponding MT populations. The median follow-up time in the study was 34.0 weeks (range, 5–223 weeks).

Efficacy. Overall, 141 patients were evaluable for response assessment. Objective responses occurred more frequently in patients with RAS WT (59%) vs RAS MT (41%) mCRC, and in patients with RAS WT/BRAF WT (68%) vs RAS or BRAF MT (37%) tumours (Table 2). Disease control rates were similar between patients with RAS WT (91%) and MT (92%) mCRC and were slightly higher in the RAS WT/BRAF WT (95%) group than in those with RAS or BRAF MT (89%) mCRC (Table 2).

The DoR analysis included only those patients who experienced a partial or complete response: 40 patients with mCRC WT for both RAS and BRAF, and 30 patients with MT RAS and WT BRAF mCRC (no patient with BRAF MT disease responded to treatment). Median DoR was longer in RAS WT (13.0 months) vs RAS MT (5.8 months) disease (HR, 0.16 (95% confidence interval (CI): 0.07, 0.37)) (Figure 1).

Median DpR ($n = 141$) was also significantly greater in patients with RAS WT (59.3% (Q1, Q3: 26.4, 77.0%)) than in those with

RAS MT (35.7% (18.8, 62.0%); $P = 0.0181$) disease. Early tumour shrinkage $\geq 30\%$ was reported in 49% of patients with RAS WT disease and 37% of those with RAS MT disease (odds ratio, 1.6), while ETS $\geq 20\%$ was reported in 74 and 50% of patients, respectively (odds ratio: 2.8). Early tumour shrinkage was associated with longer PFS in the RAS WT vs MT groups ($\geq 30\%$ cutoff: median 14.3 vs 7.8 months, HR, 0.29; $\geq 20\%$ cutoff: 13.3 vs 7.3 months, HR, 0.34). Furthermore, when patient groups were combined ($n = 135$), patients achieving ETS had significantly longer PFS than those not achieving the relevant ETS criteria (ETS $\geq 30\%$ vs $< 30\%$: median 10.9 vs 7.2 months, HR, 0.45, $P = 0.0003$; ETS $\geq 20\%$ vs $< 20\%$: 9.1 vs 6.9 months, HR, 0.48, $P = 0.0005$).

Metastasis resection rates ($n = 143$) were numerically higher in patients with RAS WT ($n = 9$; 13% (95% CI: 6.1, 23.3)) vs RAS MT mCRC ($n = 7$; 9% (3.9, 18.5)) and were also higher in patients with RAS WT/BRAF WT ($n = 9$; 15% (7.1, 26.6)) vs those with RAS or BRAF MT status ($n = 7$; 8% (3.5, 16.6)). Rates of complete

Table 1. Baseline demographics and disease characteristics

	Panitumumab + FOLFIRI ($n = 143$)			
	RAS WT ($n = 69$)	RAS MT ($n = 74$)	RAS WT/BRAF WT ($n = 60$)	RAS or BRAF MT ($n = 83$)
Male sex, n (%)	55 (80)	42 (57)	50 (83)	47 (57)
White ethnicity, n (%)	66 (96)	73 (99)	57 (95)	82 (99)
Age (years), median (range)	62.2 (38–84)	63.7 (37–80)	64.5 (38–84)	64.0 (37–80)
ECOG PS, n (%)				
0/1	66 (96)	69 (93)	57 (95)	78 (94)
2	3 (4)	5 (7) ^a	3 (5)	5 (6) ^a
Primary tumour, n (%)				
Colon	40 (58)	48 (65)	34 (57)	54 (65)
Rectum	29 (42)	26 (35)	26 (43)	29 (35)
Time since mCRC diagnosis (months), ^b median (range)	1.1 (0–29)	1.2 (0–44)	1.0 (0–29)	1.2 (0–44)
Number of metastatic sites, n (%)				
1	30 (43)	31 (42)	27 (45)	34 (41)
2	22 (32)	25 (34)	18 (30)	29 (35)
≥ 3	17 (25)	18 (24)	15 (25)	20 (24)
Sites of metastases, n (%)				
Liver only	26 (38)	20 (27)	24 (40)	22 (27)
Liver + other	33 (48)	35 (47)	29 (48)	39 (47)
Other only	10 (14)	19 (26)	7 (12)	22 (27)
Sum of longest diameters of measurable lesions (mm), ^c median (range)	135.5 (20–381)	107.0 (20–371)	136.0 (20–381)	104.0 (20–371)

Abbreviations: ECOG PS = Eastern Cooperative Oncology Group performance status; mCRC = metastatic colorectal cancer; MT = mutant; WT = wild type.

^aOne patient in the RAS mutant group had an ECOG PS of 3.

^bDate of enrolment minus date of primary diagnosis or metastatic disease.

^cTarget lesions only.

Table 2. Best response, objective response and disease control rates

	Panitumumab + FOLFIRI ($n = 141$)			
	RAS WT ($n = 68$)	RAS MT ($n = 73$)	RAS WT/RAF WT ($n = 59$)	RAS or BRAF MT ($n = 82$)
Best response, n (%)				
Complete response	2 (3)	1 (1)	2 (3)	1 (1)
Partial response	38 (56)	29 (40)	38 (64)	29 (35)
Stable disease	22 (32)	37 (51)	16 (27)	43 (52)
Disease progression	5 (7)	3 (4)	2 (3)	6 (7)
Unevaluable/not done	1 (1)	3 (4)	1 (2)	3 (4)
Objective response, n (%) [95% CI]	40 (59) [46.2, 70.6]	30 (41) [29.7, 53.2]	40 (68) [54.4, 79.4]	30 (37) [26.2, 48.0]
Unadjusted odds ratio (95% CI)	2.0 (1.0, 4.2)		3.7 (1.7, 7.9)	
Disease control, n (%) [95% CI]	62 (91) [81.8, 96.7]	67 (92) [83.0, 96.9]	56 (95) [85.9, 98.9]	73 (89) [80.2, 94.9]
Unadjusted odds ratio (95% CI)	0.9 (0.2, 3.7)		2.3 (0.5, 13.8)	

Abbreviations: CI = confidence intervals; MT = mutant; WT = wild type.

Figure 1. Duration of response by tumour RAS status.

resection were similar across the four groups (RAS WT, $n = 4$, 6% (95% CI: 1.6, 14.2); RAS MT, $n = 5$, 7% (2.2, 15.1); RAS WT/BRAF WT, $n = 4$, 7% (1.9, 16.2); RAS or BRAF MT, $n = 5$, 6% (2.0, 13.5)).

Median PFS ($n = 143$) was longer in patients with RAS WT vs MT status (11.2 vs 7.3 months, respectively; HR, 0.37 (95% CI: 0.24, 0.58)) (Figure 2A) and was also longer in patients with RAS WT/BRAF WT vs RAS or BRAF MT mCRC (13.2 vs 6.9 months, respectively; HR, 0.25 (95% CI: 0.15, 0.41)) (Figure 2B). Median TTP ($n = 143$) was longer in patients with RAS WT (13.2 (95% CI: 7.8, 17.0) months) vs MT tumours (7.3 (95% CI: 6.1, 7.6) months) and in those with RAS WT/BRAF WT (13.3 (95% CI: 9.0, 17.0) months) vs RAS or BRAF MT tumours (7.2 (95% CI: 5.7, 7.4) months).

Median overall survival ($n = 143$) was not reached in any of the four RAS/BRAF groups, although the number of deaths was numerically lower in the RAS WT ($n = 5$; 7%) and RAS WT/BRAF WT ($n = 4$; 7%) groups vs the RAS MT ($n = 11$; 15%) and RAS or BRAF MT ($n = 12$; 14%) groups (Cox proportional HR, RAS WT vs RAS MT = 0.42 (95% CI: 0.15, 1.23); RAS WT/BRAF WT vs RAS or BRAF MT = 0.37 (95% CI: 0.12, 1.15)).

Safety. For patients with RAS WT tumours ($n = 69$), the median (range) cumulative dose of panitumumab delivered (adjusted for weight) was 72.0 (6.0–395.5) mg kg^{-1} . The median (range) relative dose intensity (RDI) was 89% (35–101). For patients with RAS MT tumours ($n = 74$), the median (range) adjusted cumulative panitumumab dose was 60.4 (5.8–150.0) mg kg^{-1} , with a median (range) RDI of 90% (45–103). In the RAS WT/BRAF WT and MT groups, the median (range) adjusted cumulative panitumumab dose was 74.8 (6.0, 395.5) and 58.0 (5.8, 150.0) mg kg^{-1} , respectively, while the median (range) RDI was 89% (35–101) and 90% (45–103), respectively.

The incidence of AEs was similar regardless of tumour RAS/BRAF status (Table 3), and no new safety signals were noted in these analyses. AEs leading to study discontinuation occurred in 24–32% of patients across RAS/BRAF groups (Table 3), and treatment-related AEs leading to study discontinuation occurred in 16–27% of patients across RAS/BRAF groups (Table 3).

Skin toxicity of any grade occurred in 68 of 69 patients (99%) with RAS WT tumours, 71 of 74 patients (96%) with RAS MT tumours, 59 of 60 patients (98%) with RAS WT/BRAF WT tumours and 80 of 83 patients (96%) with RAS or BRAF MT tumours.

Amphiregulin expression. Of the 154 patients in the study, 100 had evaluable RAS status and AREG levels (RAS WT, $n = 50$; RAS MT, $n = 50$) (Supplementary Table S1). Tumours with high AREG levels more commonly had WT RAS (31 of 50; 62%) while those

with low AREG levels more commonly had MT RAS (36 of 50; 72%) (Supplementary Table S2). Among RAS WT patients, the ORR (95% CI) was 67% (51, 82) in those with high AREG expression and 38% (18, 58) in those with low AREG expression (difference: 29% (2, 53)) (Supplementary Table S3). ORRs in the RAS MT group were 31% (10, 53) and 45% (29, 60), respectively (difference, –14% (–39, 15)). The HR for PFS was more favourable in the high AREG group (RAS WT/RAS MT: 0.30 (95% CI: 0.12, 0.75)) compared with the low AREG group (0.49 (95% CI: 0.21, 1.1)) (Supplementary Figure S1).

The Cox PH model showed a significant RAS-by-AREG interaction ($P = 0.03$) (Supplementary Table S4). There was a steep transition between responders and non-responders as AREG expression decreased, which was mainly the result of changes in the RAS WT group (Supplementary Figure S2).

DISCUSSION

This analysis represents the first reported RAS data beyond *KRAS* exon 2 for panitumumab + FOLFIRI in the first-line treatment of mCRC, an indication that has recently been approved by the EMA. The results show consistently favourable efficacy for first-line panitumumab + FOLFIRI treatment in patients with RAS WT/BRAF WT tumours compared with MT mCRC tumours. This is consistent with the primary data for the *KRAS* analysis of this study (Köhne *et al*, 2012), in which response rates for patients with *KRAS* WT ($n = 86$) and *KRAS* MT ($n = 68$) tumours were 56 and 38% and median PFS was 8.9 and 7.2 months, respectively. In the current analysis, extended RAS testing identified 69 patients with RAS WT tumours, and small increases in response rate (59%) and median PFS (11.2 months) were seen in these patients compared with the primary analysis population. The results are also consistent with previous RAS analysis data for panitumumab plus FOLFIRI in the treatment of mCRC (Cohn *et al*, 2011; Mitchell *et al*, 2011; Peeters *et al*, 2014b; Abad *et al*, 2014). Thus, patients whose tumours harbour RAS mutations beyond *KRAS* exon 2 are unlikely to benefit from addition of panitumumab to FOLFIRI. Consistent results have also been reported in studies with another anti-EGFR monoclonal antibody, cetuximab (Van Cutsem *et al*, 2015; Heinemann *et al*, 2014a, b), highlighting the importance of up-front tumour RAS testing in patients being considered for EGFR inhibitor therapy. As OS was neither an end point nor followed in this trial, efficacy of panitumumab + FOLFIRI was consistent with that reported for RAS WT populations in first-line

Figure 2. Progression-free survival by tumour (A) RAS and (B) RAS/BRAF status.

Table 3. Summary of adverse events				
	Panitumumab + FOLFIRI (n = 143)			
	RAS WT (n = 69)	RAS MT (n = 74)	RAS WT/RAF WT (n = 60)	RAS or BRAF MT (n = 83)
Any AE, n (%)	69 (100)	74 (100)	60 (100)	83 (100)
Worst grade ≥ 3	59 (86)	57 (77)	52 (87)	64 (77)
Serious AE	40 (58)	38 (51)	32 (53)	46 (55)
AEs leading to discontinuation ^a	21 (30)	18 (24)	19 (32)	20 (24)
Any treatment-related AE, n (%)	69 (100)	74 (100)	60 (100)	83 (100)
Worst grade ≥ 3	50 (72)	50 (68)	44 (73)	56 (67)
Serious AE	20 (29)	18 (24)	14 (23)	24 (29)
AEs leading to discontinuation ^a	16 (23)	13 (18)	16 (27)	13 (16)

Abbreviations: AE = adverse event; MT = mutant; WT = wild type.

^aPermanent discontinuation of any study drug.

studies of panitumumab + FOLFOX4 (Douillard *et al*, 2013; Schwartzberg *et al*, 2014; Douillard *et al*, 2015), as well as in studies of cetuximab in combination with FOLFIRI or FOLFOX4 (Van Cutsem *et al*, 2015; Bokemeyer *et al*, 2011; Ciardiello *et al*, 2014). For example, in the CRYSTAL study of FOLFIRI plus cetuximab, the PFS in patients with WT RAS (11.4 months) and MT RAS (7.4 months) (Van Cutsem *et al*, 2015) was almost identical to that in the present study (11.2 and 7.3 months, respectively). The results of the present study are also consistent with previous data showing that *BRAF* mutations are associated with poor prognosis in mCRC regardless of first-line treatment (Phipps *et al*, 2012; Morris *et al*, 2014). Discussion is ongoing, however, regarding the potential usefulness of EGFR inhibitors in patients with *RAS* WT/*BRAF* MT mCRC. Two recent meta-analyses, have reached differing conclusions regarding the predictive role of *BRAF* mutations in patients receiving EGFR inhibitor therapy. While Pietrantonio *et al* (2015) focus on the lack of significant ORR, PFS or OS benefits on addition of EGFR inhibitors to chemotherapy, Rowland *et al* (2015), who included trials of first- and second-line treatment, conclude that there is insufficient evidence to demonstrate a different treatment benefit between patients with *RAS* WT/*BRAF* WT and *RAS* WT/*BRAF* MT disease, and therefore insufficient data to exclude patients with *RAS* WT/*BRAF* MT from EGFR inhibitor therapy. Regardless, such

patients should therefore be considered to be at high risk of rapid progression and should be managed accordingly.

No new safety signals were seen with the combination of panitumumab + FOLFIRI in RAS WT/BRAF WT population; AEs were similar to those seen in the KRAS exon 2 WT population of this study (Thaler *et al*, 2012) and in previous studies using this combination in patients with mCRC (Cohn *et al*, 2011; Mitchell *et al*, 2011; Peeters *et al*, 2014b). Overall 28% of patients withdrew from study treatment because of AEs. Consistent with existing data on EGFR inhibitors, there was a high incidence of skin toxicity. While there was no protocol-mandated proactive management of skin toxicity in the present study, it is now recommended for patients receiving EGFR inhibitors (Boone *et al*, 2007; Melosky *et al*, 2009). An earlier analysis of data from the present study showed a higher incidence of skin toxicity in patients with KRAS WT tumours than in those with KRAS MT tumours (Thaler *et al*, 2012), which may reflect the higher mean cumulative panitumumab dose and longer duration of treatment (i.e., panitumumab cycles) received by the KRAS WT group. However, exposure-adjusted AE rates showed integument-related toxicity overall to be higher in the KRAS MT vs WT population. Thus there is no evidence of an association between tumour KRAS/RAS status and toxicity. Furthermore, despite the high incidence of skin toxicity, generic quality of life (QoL) instruments have shown no impact of EGFR inhibitors plus FOLFIRI on overall QoL (Melosky *et al*, 2009; Bennett *et al*, 2011; Thaler *et al*, 2012). Although proactive management of skin toxicity may have reduced the impact for patients, it may also be that the QoL tools used in the study provided too general an assessment to determine the true impact of this AE. Future trials of EGFR inhibitors should include skin-toxicity specific QoL assessment tools to support better understanding of the true impact of this AE on patient wellbeing.

Among patients with RAS WT mCRC, high AREG expression was associated with response to panitumumab + FOLFIRI. Consistent with other studies (Jacobs *et al*, 2009; Baker *et al*, 2011; Pentheroudakis *et al*, 2013; Loupakakis *et al*, 2014; Jonker *et al*, 2014; Stahler *et al*, 2016), there was an interaction between RAS and AREG levels. A higher percentage of patients with RAS WT mCRC had high AREG levels compared with those with RAS MT mCRC, suggesting that AREG levels are associated with EGFR signalling. Further studies are needed to determine whether there is an AREG expression level below which there is little or no response to panitumumab + FOLFIRI treatment. Biomarker studies remain critical to our understanding of targeted agents in mCRC, and warrant further investigation.

One of the strengths of this comprehensive study was the high percentage of patients from the original cohort who were available for RAS/BRAF testing – which was conducted centrally for all specimens – allowing the identification of additional risks for lack of response to treatment. It should be noted that the analyses were retrospective and exploratory in nature, although, as noted above, the results were consistent with previous analyses of efficacy by RAS mutation status in patients with mCRC.

In conclusion, first-line panitumumab + FOLFIRI was associated with consistently favourable efficacy in patients with RAS WT/BRAF WT vs MT mCRC tumours and was well tolerated, despite the expected high incidence of skin toxicity. The combination of first-line panitumumab + FOLFIRI also gave efficacy similar to that seen in the RAS WT populations in other first-line studies of EGFR-targeted agents plus FOLFIRI or FOLFOX. As per the licensed indication for panitumumab, therefore, patients with RAS mutations should not receive panitumumab treatment. Across all lines of therapy, determination of tumour RAS status improves identification of patients unlikely to respond to treatment with panitumumab compared with evaluation of KRAS exon 2 alone. The combination of panitumumab with

FOLFIRI can be considered as an important treatment option for first-line patients with RAS WT/BRAF WT mCRC.

ACKNOWLEDGEMENTS

This work was supported by Amgen (Europe) GmbH. Medical writing support (funded by Amgen (Europe) GmbH) was provided by Dan Booth PhD (Bioscript Medical Ltd).

CONFLICT OF INTEREST

MK is an advisor for, and has received honoraria from, Amgen Ltd. R-DH has received honoraria from Amgen Ltd. LM is an advisor for, and has received honoraria and research funding from, Amgen. HL has acted as a consultant/advisor for Amgen Ltd & Roche Ltd. RG has received research support from Amgen. JT has received honoraria and research funding from Amgen Ltd. KSO is a former employee of Amgen Inc. MB, BT, YZ, GD are employees of Amgen Inc. C-HK has acted as a consultant/advisor to Amgen Ltd, Merck KG Darmstadt & Roche Ltd. EF declares no conflict of interest.

REFERENCES

- Abad A, Massuti B, Gravalos C, Escudero P, Guillen C, Manzano JL, Gomez MA, Safont MJ, Plazas JG, Sastre J, Pericay C, Dueñas R, López-López C, Losa F, Ayerbes MV, González E, Yuste A, Carrato A, Aranda E (2014) Phase II trial of panitumumab plus FOLFOX4 or FOLFIRI in subjects with KRAS wild-type colorectal cancer and liver-limited disease: the PLANET study. *J Clin Oncol* **32**(5 Suppl): Abstract 3560.
- Baker JB, Dutta D, Watson D, Maddala T, Munneke BM, Shak S, Rowinsky EK, Xu LA, Harbison CT, Clark EA, Mauro DJ, Khambata-Ford S (2011) Tumour gene expression predicts response to cetuximab in patients with KRAS wild-type metastatic colorectal cancer. *Br J Cancer* **104**: 488–495.
- Bennett L, Zhao Z, Barber B, Zhou X, Peeters M, Zhang J, Xu F, Wizeorek J, Douillard JY (2011) Health-related quality of life in patients with metastatic colorectal cancer treated with panitumumab in first- or second-line treatment. *Br J Cancer* **105**: 1495–1502.
- Bokemeyer C, Bondarenko I, Hartmann JT, de Braud F, Schuch G, Zubel A, Celik I, Schlichting M, Koralewski P (2011) Efficacy according to biomarker status of cetuximab plus FOLFOX-4 as first-line treatment for metastatic colorectal cancer: the OPUS study. *Ann Oncol* **22**: 1535–1546.
- Bokemeyer C, Köhne CH, Ciardiello F, Lenz HJ, Heinemann V, Klinkhardt U, Beier F, Duecker K, van Krieken JH, Tejpar S (2015) FOLFOX4 plus cetuximab treatment and RAS mutations in colorectal cancer. *Eur J Cancer* **51**: 1243–1252.
- Boone SL, Rademaker A, Liu D, Pfeiffer C, Mauro DJ, Lacouture ME (2007) Impact and management of skin toxicity associated with anti-epidermal growth factor receptor therapy: survey results. *Oncology* **72**: 152–159.
- Ciardiello F, Lenz H-J, Köhne CH, Heinemann V, Tejpar S, Esser R, Beier F, Stroh C, Duecker K, Van Cutsem E (2014) Effect of KRAS and NRAS mutational status on first-line treatment with FOLFIRI plus cetuximab in patients with metastatic colorectal cancer (mCRC): new results from the CRYSTAL trial. *J Clin Oncol* **32**(Suppl 3): Abstract #LBA443.
- Cohn AL, Shumaker GC, Khandelwal P, Smith DA, Neubauer MA, Mehta N, Richards D, Watkins DL, Zhang K, Yassine MR (2011) An open-label, single-arm, phase 2 trial of panitumumab plus FOLFIRI as second-line therapy in patients with metastatic colorectal cancer. *Clin Colorectal Cancer* **10**: 171–177.
- Douillard JY, Douillard JY, Oliner KS, Siena S, Tabernero J, Burkes R, Barugel M, Humblet Y, Bodoky G, Cunningham D, Jassam J, Rivera F, Kocáková I, Ruff P, Błasińska-Morawiec M, Smakal M, Canon JL, Rother M, Williams R, Rong A, Wizeorek J, Sidhu R, Patterson SD (2013) Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer. *N Engl J Med* **369**: 1023–1034.

- Douillard JY, Siena S, Peeters M, Koukakis R, Terwey JH, Tabernero J (2015) Impact of early tumour shrinkage and resection on outcomes in patients with wild-type RAS metastatic colorectal cancer. *Eur J Cancer* **51**: 1231–1242.
- Heinemann V, Modest D, Fischer von Weikersthal L, Decker T, Kiani A, Vehling-Kaiser U, Al-Batran S-E, Heintges T, Lerchenmüller C, Kahl C, Seipelt G, Kullmann F, Stauch M, Scheithauer W, Held S, Giessen C, Möhler M, Jagenburg A, Stintzing S (2014a) Independent radiological evaluation of objective response by early tumor shrinkage, and depth of response in FIRE-3 (AIO KRK-0306). *Ann Oncol* **25**(Suppl 2): ii117 Abstract O-0030.
- Heinemann V, von Weikersthal LF, Decker T, Kiani A, Vehling-Kaiser U, Al-Batran SE, Heintges T, Lerchenmüller C, Kahl C, Seipelt G, Kullmann F, Stauch M, Scheithauer W, Hielscher J, Scholz M, Müller S, Link H, Niederle N, Rost A, Höffkes HG, Moehler M, Lindig RU, Modest DP, Rossius L, Kirchner T, Jung A, Stintzing S (2014b) FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomised, open-label, phase 3 trial. *Lancet Oncol* **15**: 1065–1075.
- Jacobs B, De Roock W, Piessevaux H, Van Oirbeek R, Biesmans B, De Schutter J, Fieuws S, Vandesompele J, Peeters M, Van Laethem JL, Humblet Y, Pénault-Llorca F, De Hertogh G, Laurent-Puig P, Van Cutsem E, Tejpar S (2009) Amphiregulin and epiregulin mRNA expression in primary tumors predicts outcome in metastatic colorectal cancer treated with cetuximab. *J Clin Oncol* **27**: 5068–5074.
- Joensuu H, Vehtari A, Riihimäki J, Nishida T, Steigen SE, Brabec P, Plank L, Nilsson B, Cirilli C, Braconi C, Bordoni A, Magnusson MK, Linke Z, Sufliarsky J, Federico M, Jonasson JG, Dei Tos AP, Rutkowski P (2012) Risk of recurrence of gastrointestinal stromal tumour after surgery: an analysis of pooled population-based cohorts. *Lancet Oncol* **13**: 265–274.
- Jonker DJ, Karapetis CS, Harbison C, O'Callaghan CJ, Tu D, Simes RJ, Malone DP, Langer C, Tebbutt N, Price TJ, Shapiro J, Siu LL, Wong RP, Bjarnason G, Moore MJ, Zalberg JR, Khambata-Ford S (2014) Epiregulin gene expression as a biomarker of benefit from cetuximab in the treatment of advanced colorectal cancer. *Br J Cancer* **110**: 648–655.
- Köhne CH, Hofheinz R, Mineur L, Letocha H, Greil R, Thaler J, Fernebro E, Gamelin E, Decosta L, Karthaus M (2012) First-line panitumumab plus irinotecan/5-fluorouracil/leucovorin treatment in patients with metastatic colorectal cancer. *J Cancer Res Clin Oncol* **138**: 65–72.
- Lacouture ME, Mitchell EP, Piperdi B, Pillai MV, Shearer H, Iannotti N, Xu F, Yassine M (2010) Skin toxicity evaluation protocol with panitumumab (STEP), a phase II, open-label, randomized trial evaluating the impact of a pre-emptive skin treatment regimen on skin toxicities and quality of life in patients with metastatic colorectal cancer. *J Clin Oncol* **28**: 1351–1357.
- Loupakis F, Cremolini C, Fioravanti A, Orlandi P, Salvatore L, Masi G, Schirripa M, Di Desidero T, Antoniotti C, Canu B, Faviana P, Sensi E, Lupi C, Fontanini G, Basolo F, Di Paolo A, Danesi R, Falcone A, Bocci G (2014) EGFR ligands as pharmacodynamic biomarkers in metastatic colorectal cancer patients treated with cetuximab and irinotecan. *Target Oncol* **9**: 205–214.
- Melosky B, Burkes R, Rayson D, Alcindor T, Shear N, Lacouture M (2009) Management of skin rash during EGFR-targeted monoclonal antibody treatment for gastrointestinal malignancies: Canadian recommendations. *Curr Oncol* **16**: 16–26.
- Mitchell EP, Piperdi B, Lacouture ME, Shearer H, Iannotti N, Pillai MV, Xu F, Yassine M (2011) The efficacy and safety of panitumumab administered concomitantly with FOLFIRI or Irinotecan in second-line therapy for metastatic colorectal cancer: the secondary analysis from STEP (Skin Toxicity Evaluation Protocol With Panitumumab) by KRAS status. *Clin Colorectal Cancer* **10**: 333–339.
- Morris V, Overman MJ, Jiang ZQ, Garrett C, Agarwal S, Eng C, Kee B, Fogelman D, Dasari A, Wolff R, Maru D, Kopetz S (2014) Progression-free survival remains poor over sequential lines of systemic therapy in patients with BRAF-mutated colorectal cancer. *Clin Colorectal Cancer* **13**: 164–171.
- National Comprehensive Cancer Network (2015) *NCCN Clinical Practice Guidelines in Oncology. Colon Cancer*. V2.2015. Jamestown, PA: National Comprehensive Cancer Network.
- Peeters M, Oliner KS, Parker A, Siena S, Van Cutsem E, Huang J, Humblet Y, Van Laethem JL, André T, Wizek J, Reese D, Patterson SD (2013) Massively parallel tumor multigene sequencing to evaluate response to panitumumab in a randomized phase III study of metastatic colorectal cancer. *Clin Cancer Res* **19**: 1902–1912.
- Peeters M, Oliner KS, Price TJ, Cervantes A, Sobrero AF, Ducreux M, Hotko Y, Andre T, Chan E, Lordick F, Punt CJA, Strickland A, Wilson G, Ciuleanu TE, Roman L, Van Cutsem E, Yu H, Jung AS, Sidhu R, Patterson SD (2014a) Updated analysis of KRAS/NRAS and BRAF mutations in study 20050181 of panitumumab (pmab) plus FOLFIRI for second-line treatment (tx) of metastatic colorectal cancer (mCRC). *J Clin Oncol* **32**(Suppl 5): Abstract #3568.
- Peeters M, Oliner KS, Price TJ, Cervantes A, Sobrero AF, Ducreux M, Hotko Y, Andre T, Chan E, Lordick F, Punt CJA, Strickland A, Wilson G, Ciuleanu T-E, Roman L, Van Cutsem E, Tian Y, Jung AS, Sidhu R, Patterson SD (2014c) Analysis of KRAS/NRAS mutations in phase 3 study 20050181 of panitumumab (pmab) plus FOLFIRI versus FOLFIRI for second-line treatment (tx) of metastatic colorectal cancer (mCRC). *J Clin Oncol* **32**(Suppl 3): Abstract #LBA387.
- Peeters M, Price TJ, Cervantes A, Sobrero A, Ducreux MP, André T, Lordick F, Punt CJA, Koukakis R, Terwey J, van Cutsem E (2014b) Tumour shrinkage and response outcomes during second-line panitumumab (pmab) + FOLFIRI vs FOLFIRI treatment. *Ann Oncol* **25**(Suppl 4): Abstract #7739 (546P).
- Pentheroudakis G, Kotoula V, De Roock W, Kouvatsas G, Papakostas P, Makatsoris T, Papamichael D, Xanthakis I, Sgouros J, Televantou D, Kafiri G, Tsamandas AC, Razis E, Galani E, Bafaloukos D, Efstratiou I, Bompalaki I, Pectasides D, Pavlidis N, Tejpar S, Fountzilas G (2013) Biomarkers of benefit from cetuximab-based therapy in metastatic colorectal cancer: interaction of EGFR ligand expression with RAS/RAF, PIK3CA genotypes. *BMC Cancer* **13**: 49.
- Phipps AI, Buchanan DD, Makar KW, Burnett-Hartman AN, Coghill AE, Passarelli MN, Baron JA, Ahnen DJ, Win AK, Potter JD, Newcomb PA (2012) BRAF mutation status and survival after colorectal cancer diagnosis according to patient and tumor characteristics. *Cancer Epidemiol Biomarkers Prev* **21**: 1792–1798.
- Pietrantonio F, Cremolini C, Petrelli F, Di Bartolomeo M, Loupakis F, Maggi C, Antoniotti C, de Braud F, Falcone A, Iacovelli R (2015) First-line anti-EGFR monoclonal antibodies in panRAS wild-type metastatic colorectal cancer: A systematic review and meta-analysis. *Crit Rev Oncol Hematol* **96**: 156–166.
- Rowland A, Dias MM, Wiese MD, Kichenadasse G, McKinnon RA, Karapetis CS, Sorich MJ. Meta-analysis of BRAF mutation as a predictive biomarker of benefit from anti-EGFR monoclonal antibody therapy for RAS wild-type metastatic colorectal cancer (2015) *Br J Cancer* **112**: 1888–1894.
- Schwartzberg LS, Rivera F, Karthaus M, Fasola G, Canon JL, Hecht JR, Yu H, Oliner KS, Go WY (2014) PEAK: a randomized, multicenter phase II study of panitumumab plus modified fluorouracil, leucovorin, and oxaliplatin (mFOLFOX6) or bevacizumab plus mFOLFOX6 in patients with previously untreated, unresectable, wild-type KRAS exon 2 metastatic colorectal cancer. *J Clin Oncol* **32**: 2240–2247.
- Stahler A, Heinemann V, Giessen-Jung C, Crispin A, Schallhorn A, Stintzing S, Fischer von Weikersthal L, Vehling-Kaiser U, Stauch M, Quietzsch D, Held S, von Einem JC, Holch J, Neumann J, Kirchner T, Jung A, Modest DP (2016) Influence of mRNA expression of epiregulin and amphiregulin on outcome of patients with metastatic colorectal cancer treated with 5-FU/LV plus irinotecan or irinotecan plus oxaliplatin as first-line treatment (FIRE 1-trial). *Int J Cancer* **138**: 739–746.
- Thaler J, Karthaus M, Mineur L, Greil R, Letocha H, Hofheinz R, Fernebro E, Gamelin E, Baños A, Köhne CH (2012) Skin toxicity and quality of life in patients with metastatic colorectal cancer during first-line panitumumab plus FOLFIRI treatment in a single-arm phase II study. *BMC Cancer* **12**: 438.
- Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van Glabbeke M, van Oosterom AT, Christian MC, Gwyther SG (2000) New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* **92**: 205–216.
- Van Cutsem E, Lenz HJ, Köhne CH, Heinemann V, Tejpar S, Melezinek I, Beier F, Stroh C, Rougier P, van Krieken JH, Ciardiello F (2015) Fluorouracil, leucovorin, and irinotecan plus cetuximab treatment and RAS mutations in colorectal cancer. *J Clin Oncol* **33**: 692–700.
- Vanhatalo J, Riihimäki J, Hartikainen J, Jylänki P, Tolvanen V, Vehtari A (2013) GPstuff: Bayesian modeling with Gaussian processes. *J Mach Learn Res* **14**: 1179.

Venook AP, Niedzwiecki D, Lenz H-J, Innocenti F, Mahoney MR, O'Neil BH, Shaw JE, Polite BN, Hochster HS, Atkins JN, Goldberg RM, Mayer RJ, Schilsky RL, Bertagnolli MM, Blanke CD. Cancer and Leukemia Group B (Alliance), SWOG, and ECOG (2014) CALGB/SWOG 80405: Phase III trial of irinotecan/5-FU/leucovorin (FOLFIRI) or oxaliplatin/5-FU/leucovorin (mFOLFOX6) with bevacizumab (BV) or cetuximab (CET) for patients (pts) with *KRAS* wild-type (wt) untreated metastatic

adenocarcinoma of the colon or rectum (MCRC). *J Clin Oncol* 32(5 Suppl): Abstract LBA3.

This work is licensed under the Creative Commons Attribution 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Supplementary Information accompanies this paper on British Journal of Cancer website (<http://www.nature.com/bjc>)