

Keywords: hepatocholangiocarcinoma; chemotherapy; prognostic factors

Gemcitabine plus platinum-based chemotherapy for first-line treatment of hepatocholangiocarcinoma: an AGEO French multicentre retrospective study

Maëva Salimon¹, Caroline Prioux-Klotz², David Tougeron³, Vincent Hautefeuille⁴, Morgane Caulet⁵, Jérôme Gournay¹, Tamara Matysiak-Budnik¹, Jaafar Bennouna¹, Manuela Tiako Meyo⁶, Thierry Lecomte⁵, Aziz Zaanani⁶ and Yann Touchefeu^{*1}

¹Institut des Maladies de l'appareil Digestif, Nantes University Hospital, 1 place Alexis Ricordeau, Nantes 44000, France; ²Department of Gastroenterology and Oncology, Cochin University Hospital, Paris 75014, France; ³Department of Gastroenterology, Poitiers University Hospital, Poitiers 86600, France; ⁴Amiens University Hospital, Department of Gastroenterology and Oncology, Amiens 80054, France; ⁵Tours University Hospital, Department of Gastroenterology and Oncology, Tours 37000, France and ⁶Department of Gastroenterology and Digestive Oncology, Sorbonne Paris Cité, Paris Descartes University, Georges Pompidou European Hospital, Paris 75015, France

Background: Hepatocholangiocarcinoma (cHCC-ICC) is a rare liver tumour for which no data on chemosensitivity exist. The aims of this multicentre study were to evaluate overall survival (OS), progression-free survival (PFS), and prognostic factors in cHCC-ICC treated by gemcitabine plus platinum as first-line.

Methods: Unresectable cHCC-ICC treated by gemcitabine plus platinum-based chemotherapy between 2008 and 2017 were retrospectively analysed. Diagnosis was based on histology or, in case of ICC or HCC histology, on discordant computerised tomography scan enhancement patterns associated with discordant serum tumour marker elevation suggesting the alternative tumour. OS and PFS were evaluated by Kaplan–Meier method and prognostic factors by Log-rank test and Cox model.

Results: Among 30 patients included, cHCC-ICC was histologically proven in 22 (73.3%). 18 (60%) received gemcitabine plus oxaliplatin (GEMOX), 9 (30%) GEMOX plus bevacizumab, and 3 (10%) gemcitabine plus cisplatin. RECIST criteria were reported in 28 patients: 8 (28.6%) showed partial response, 14 (50%) stable disease, and 6 (21.4%) tumour progression at first evaluation. Median PFS and OS were 9.0 and 16.2 months, respectively. Serum bilirubin $\geq 30 \mu\text{mol l}^{-1}$ ($P=0.001$) and positive serology for HBV and/or HCV ($P=0.014$) were independent poor prognostic factors for OS.

Conclusions: Gemcitabine plus platinum-based chemotherapy is effective as first-line for advanced cHCC-ICC.

Hepatocholangiocarcinoma (cHCC-ICC) is a rare primary hepatic tumour with both, hepatocellular carcinoma (HCC) and cholangiocarcinoma (ICC) histological, features. Its first histological classification dates from 1949 by Allen and Lisa (Allen and Lisa,

1949) and it is now described in the 2010 WHO classification (Theise *et al*, 2010). The oncogenesis of this tumour remains unclear with hypotheses of the existence of progenitors capable of double differentiation or dedifferentiation of matures hepatocytes

*Correspondence: Dr Y Touchefeu; E-mail: yann.touchefeu@chu-nantes.fr

Received 11 July 2017; revised 17 October 2017; accepted 20 October 2017; published online 23 November 2017

© 2018 Cancer Research UK. All rights reserved 0007–0920/18

(Kim *et al*, 2004). Moreover, in murine models, existence of several intra-hepatic cell clones with differential potential into hepatocyte and biliary differentiation at the origin of mixed tumours has been described (Piscaglia *et al*, 2009).

The prevalence of cHCC-ICC varies from 1 to 5% of primary liver cancer in Asia and Western countries in different surgical series (Lee *et al*, 2006; Wachtel *et al*, 2008; Bergquist *et al*, 2016). This tumour is more frequent in males, and it is sometimes associated with chronic viral hepatitis B and C, especially in Asian countries. Cirrhosis is associated in ~30% of cases (Lee *et al*, 2006; Chok *et al*, 2009; Bergquist *et al*, 2016). About 30% of cHCC-ICC are diagnosed at an advanced stage with synchronous metastases (Wachtel *et al*, 2008; Wang *et al*, 2010; Bergquist *et al*, 2016). The diagnosis may be difficult, and may be based on the analysis of surgical resection specimen or liver biopsy. However, the mixed feature of cHCC-ICC can be misdiagnosed by a biopsy with the identification of the HCC or the ICC histological component only (Tagushi *et al*, 1996). Some typical cHCC-ICC radiological images have been described on contrast enhanced computerised tomography scan (CT-scan) and magnetic resonance imaging, combining, both, arterial phase enhancement/portal venous washout typical of HCC and progressive fibrous stroma central enhancement typical of ICC. Elevation of tumour serum markers can suggest the diagnosis of HCC or ICC, namely, alpha-fetoprotein (α FP) for HCC, and carbohydrate antigen 19-9 (CA 19-9) and carcinoembryonic antigen for ICC. The combination of typical imaging of either HCC or ICC with the elevation of serum tumour markers suggesting the alternative tumour can also help to identify cHCC-ICC (Jarnagin *et al*, 2002; Tang *et al*, 2006; Maximin *et al*, 2014; Li *et al*, 2016).

There has been no randomised trial investigating the specific management of cHCC-ICC. Few studies have reported the clinical outcomes after resection, liver transplantation, or transarterial chemoembolization (TACE) for localised cHCC-ICC (Tagushi *et al*, 1996; Lee *et al*, 2006; Kim *et al*, 2010; Panjala *et al*, 2010; Wang *et al*, 2010; Sapisochin *et al*, 2011; Groeschl *et al*, 2013; Park *et al*, 2013; Garancini *et al*, 2014; Vilchez *et al*, 2016). Concerning advanced tumours, Sorafenib is the standard of care for HCC with median overall survival (OS) ranging from 6.5 to 10.7 months (Llovet *et al*, 2008; Cheng *et al*, 2009), whereas the combination of gemcitabine and platinum (cisplatin or oxaliplatin) is the standard first-line chemotherapy for ICC with median OS of 11.7 months with GEMCIS regimen in the ABC-02 trial (Valle *et al*, 2010). Few data regarding gemcitabine and platinum-based chemotherapy for HCC have been published. In a large retrospective including 204 patients, median OS was 9 months in patients treated with GEMOX regimen (Zaanan *et al*, 2013). Bevacizumab, a vascular endothelial growth factor-antibody, in association with gemcitabine and oxaliplatin, has also been investigated in both HCC and advanced biliary-tract cancers but with no demonstrated benefits (Zhu *et al*, 2006, 2010).

To our knowledge, there is no published data investigating the systemic treatment of advanced cHCC-ICC. The main objective of this retrospective, multicentre study was to evaluate the efficacy of a gemcitabine plus platinum-based chemotherapy in first-line treatment of unresectable, advanced cHCC-ICC.

PATIENTS AND METHODS

Patients. Patients with unresectable cHCC-ICC treated with gemcitabine plus platinum-based first-line therapy between January 2008 and February 2017 in six French centres, were retrospectively included. Inclusion criteria were patients older than 18 years and gemcitabine plus cisplatin or oxaliplatin (GEMCIS or GEMOX regimen) as first-line systemic therapy, with ECOG score ≤ 2 before initiation of chemotherapy.

We included the patients with histological diagnosis of cHCC-ICC, or in case of ICC or HCC typical histology, with discordant CT-scan enhancement findings associated with serum tumour marker elevations, (typical ICC histology but HCC enhancement pattern: early arterial enhancement with early 'washout' corresponding at an hypoattenuating in the portal venous phase, with elevated α FP; typical HCC histology but ICC enhancement pattern: heterogeneous minor peripheral enhancement with gradual enhancement centrally, with elevated CA19-9). Data were last updated in June 2017.

Treatment. The treatment mostly consisted of intra-venous administration of gemcitabine at a standard dose of 1000 mg m⁻² within 30 min and oxaliplatin at a standard dose of 85–100 mg m⁻² within 2 h, given every 2 weeks (GEMOX) or gemcitabine at a standard dose of 1000–1250 mg m⁻² within 30 min and cisplatin at a standard dose of 25 mg m⁻² within 1 h on days 1 and 8, every 3 weeks (GEMCIS) with possible dose adjustments. When bevacizumab was associated with GEMOX regimen, the treatment was administrated at dose of 5 mg kg⁻¹ every 2 weeks. Treatments were administrated until disease progression.

Outcome measures. Tumour response rates were assessed by RECIST criteria. OS was defined from the first day of chemotherapy until death from any reason. Progression-free survival (PFS) was defined from the first day of chemotherapy until disease progression or death, whichever occurred first. Patients who were alive and did not experience any of these events were censored at the date of the last follow-up.

Statistics. OS and PFS were calculated according to the Kaplan-Meier method. Prognostic factors of OS and PFS were analysed in univariate analysis with the Log-rank test. Variables with a *P*-value < 0.05 or clinically relevant with a *P* ≤ 0.20 in univariate analysis were included in the multivariate analysis, performed with the Cox proportional hazard model, with a significance level of *P* < 0.05 . Analyses were performed using the software Graph Pad Prism 6 and XLStat 2017.

Consent and ethics statement. It was a retrospective study including patients managed with standard care only. The majority of patient was dead or lost to follow-up at the time of data collection. A consent form was not required for this study. The study has been performed according to the Declaration of Helsinki and its latter amendments.

RESULTS

Patient and tumour characteristics. Thirty patients were included (Flow chart, Figure 1). The main patients' characteristics are presented in Table 1. Most of the patients were males (66.7%), 27 (90%) of good general status (ECOG 0–1), 8 patients had cirrhosis (26.7%), 3 patients had positive HBV (10%) and 3 had positive HCV serology (10%), with one co-infection. Most of the patients (53.3%) had synchronous metastases. Two patients (6.67%) initially treated by surgery (one local surgery and one liver transplantation) were included after disease progression. The median follow-up for all the patients was 12.75 months (range 1–51.5 months).

Four patients had a biliary obstruction treated by drainage before chemotherapy. In other patients, elevated serum bilirubin was related to the liver dysfunction.

One patient was excluded because of an ECOG score = 3. This patient died after 4 days after the first cycle of GEMOX owing to a liver failure secondary to drug toxicity.




Figure 1. Flow-chart.

Treatment. Eighteen (60%) and 9 (30%) patients received GEMOX chemotherapy alone or in combination with bevacizumab, respectively. Three patients (10%) received GEMCIS chemotherapy. Overall, the median number of cycles for the first line of chemotherapy was 10 (range 3–28).

In the second line chemotherapy, 14 patients (46.7%) were treated with FOLFIRI ($n=6$, 42.8%), capecitabine ($n=3$, 21.4%), LV5FU2 ($n=2$, 14.3%), or other regimens (LV5FU2-cisplatin, cyclophosphamide, or sunitinib) for the three remaining patients. Five patients (16.7%) received a third line of treatment: FOLFIRI regimen ($n=1$), FOLFOX regimen ($n=1$), GEMOX regimen ($n=1$), sorafenib ($n=1$), and sunitinib ($n=1$).

Efficacy

Tumour response. RECIST assessment was performed in 28 patients (93.3%) with measurable disease. At the first evaluation, a partial response was observed in 8 (28.6%), stable disease in 14 (50%), and progression in 6 (21.4%) patients. None of the patients had tumour resection after first-line chemotherapy.

OS. Median OS was 16.2 months. The 1-year and 2-year OS rates were 66% and 26.1%, respectively (Figure 2A).

Median OS of the 22 patients with histological diagnosis of cHCC-ICC was identical, 16.2 months (Figure 2B).

PFS. Median PFS was 9.0 months. The 1-year and 2-year PFS rates were 24.2% and 9.7%, respectively (Figure 2A).

Median PFS of the 22 patients with histological diagnosis of cHCC-ICC was 8.1 months, without significant difference compared with the overall population (Figure 2B).

Prognostic factors

PFS. In univariate analysis, synchronous metastases and serum bilirubin level $\geq 30 \mu\text{mol l}^{-1}$ were the two significant factors associated with PFS: HR = 2.44; 95% CI (1.19–11.03); $P=0.023$ and HR = 2.57; 95% CI (1.32–11.53); $P=0.019$, respectively. According to the results of the univariate analysis, four factors were included in the multivariate analysis (gender, synchronous metastases, serum bilirubin level $\geq 30 \mu\text{mol l}^{-1}$ and positive serology for HBV and/or HCV). Synchronous metastases (HR = 4.0, 95% CI (1.40–11.40), $P=0.009$), serum bilirubin level

Table 1. Patient characteristics (30 patients)

	N	%
Age (years (range))	64.5 (41–88)	
Sex		
Male	20	66.7
Female	10	33.3
Baseline performance status		
0–1	27	90
2	3	10
Cirrhosis	8	26.7
Child–Pugh Score A	7	90
Child–Pugh Score B	1	10
Positive serology		
HBV	3	10
HCV	3	10
Chronic alcoholism	5	16.7
Obesity (BMI >30)	6	20
Diabetes	8	26.7
Primary sclerosing cholangitis	0	0
Histological diagnosis (cHCC-ICC)	22	73.3
Discordant imaging and/or serum tumour markers		
HCC histology and discordant imaging and elevated CA 19-9	2	6.7
ICC histology and discordant imaging and elevated αFP	6	20
Wash-in/wash-out criteria		
CT-scan	18	60
MRI	12	40
Hepatic lesion(s)		
Unique	13	43.3
Multiple	17	56.7
Initial surgery treatment	2	6.7
(included transplant)	1	3.3
Tumour markers (median, (range); % > N)		
AFP	5.3 $\mu\text{g l}^{-1}$ (2.5–2000)	50
CA19-9	83 IU ml^{-1} (5.9–20 000)	57.8
ACE	3 $\mu\text{g l}^{-1}$ (0.7–275)	23.5
Total bilirubin level at diagnosis (median, (range))	12.53 $\mu\text{mol l}^{-1}$ (4–531)	
Extra-hepatic synchronous metastases		
At least one site	12	40
Lung	5	41.7
Peritoneum	4	33.3
Bone	3	25
Pancreas	1	8.3
Oesophagus	1	8.3
Adrenal gland	1	8.3

$\geq 30 \mu\text{mol l}^{-1}$ (HR = 4.59, 95% CI (1.43–14.78), $P=0.011$) and positive serology for HBV and/or HCV (HR = 7.27, 95% CI (1.71–30.94), $P=0.007$) were significant independent poor prognostic factors for PFS. Results are detailed in Table 2.

OS. In univariate analysis, serum bilirubin level $\geq 30 \mu\text{mol l}^{-1}$ was the only significant poor prognostic factor for OS (HR = 3.66; 95% CI (2.50–34.03); $P=0.002$). According to the results of the univariate analysis, four factors were included in the multivariate analysis (gender, synchronous metastases, serum bilirubin level $\geq 30 \mu\text{mol l}^{-1}$ and positive serology for HBV and/or HCV). Serum bilirubin level $\geq 30 \mu\text{mol l}^{-1}$ (HR = 10.23, 95% CI (2.51–41.70), $P=0.001$) and positive serology for HBV and/or HCV (HR = 6.89, 95% CI (1.47–32.20), $P=0.014$) were significant independent poor prognostic factors for OS. Results are detailed in Table 3.




Figure 2. Kaplan-Meier survival curves. (A) Overall survival (OS) and progression-free survival (PFS) of the 30 patients with unresectable cHCC-ICC. (B) overall survival (OS) and progression-free survival (PFS) of the 22 patients with histological proven and unresectable cHCC-ICC.

There was no difference in terms of OS and PFS between patients with high serum level of CA 19-9 versus elevated level of α FP.

DISCUSSION

To our knowledge, this is the first published report of systemic treatment for advanced cHCC-ICC. In patients with advanced cHCC-ICC treated with gemcitabine combined with cisplatin or oxaliplatin, the median PFS and OS were 9.0 and 16.2 months, respectively. In addition, the 28.6% response rate and 50% disease control rate suggest a chemosensitivity of these tumours and are close to the results obtain with these drugs in ICC. Serum bilirubin level $\geq 30 \mu\text{mol/l}$ and positive serology for HBV and/or HCV were independent prognostic factors for OS and PFS, whereas the presence of synchronous metastases was independent prognostic factor for PFS.

There are many retrospective data on outcomes after surgical resection of cHCC-ICC, median OS after curative ranging from 4 to 48 months (Chok *et al*, 2009). The outcomes of liver transplantation are less favorable than for HCC (Panjala *et al*, 2010; Sapisochin *et al*, 2011; Groeschl *et al*, 2013; Park *et al*, 2013; Garancini *et al*, 2014; Vilchez *et al*, 2016). Some studies have investigated TACE for cHCC-ICC. In a retrospective study investigating TACE in 50 patients, median OS was 12.3 months with 70% of patients considered as responders, 85% of them having hypervascularized tumours likely to have a predominant HCC component (Kim *et al*, 2010). Sorafenib is the standard of care for advanced HCC, with time to progression ranging from 2.8 to 5.5 months and OS ranging from 6.5 to 11.7 months in phase III trials (Llovet *et al*, 2008; Cheng *et al*, 2009). Combination of gemcitabine and cisplatin is the standard first-line chemotherapy for advanced ICC, with median PFS and OS of 8.0 and 11.7 months in the ABC-02 trial (Valle *et al*, 2010). The observed 9.0 months PFS and 16.2 months OS in our study are in accordance with the results obtained

Table 2. Prognostic factors for progression-free survival (PFS): hazard-ratio (HR), CI 95%, and P-value in univariate and multivariate analysis

Variables	Univariate analysis			Multivariate analysis		
	HR	CI 95%	P-value	HR	CI 95%	P-value
Female sex	0.43	(0.19–1.00)	0.05	0.59	(0.20–1.75)	0.35
Synchronous metastases	2.44	(1.19–10.03)	0.023	4.00	(1.40–11.40)	0.009
Serum bilirubin level $\geq 30 \mu\text{mol l}^{-1}$	2.57	(1.32–11.53)	0.019	4.59	(1.43–14.78)	0.011
Positive serology for HBV and/or HCV	2.32	(0.89–12.6)	0.08	7.27	(1.71–30.94)	0.007
Age ≥ 75 years	0.49	(0.22–1.14)	0.11			
Underlying cirrhosis	1.02	(0.42–2.50)	0.96			
Chronic alcoholism	1.83	(0.58–8.47)	0.21			
Obesity	0.63	(0.25–1.68)	0.38			
Diabetes	0.92	(0.37–2.29)	0.86			
CA 19-9 $\geq 80 \text{IU ml}^{-1}$	1.14	(0.49–2.66)	0.76			

Statistically significant results are shown in bold ($P < 0.05$).

in ICC, although no conclusion can be drawn from this retrospective study.

Our patients' population showed similar characteristics that those described for cHCC-ICC. Median age was 64.5 years, similar to that from the literature (62–65 years), with a higher prevalence of male sex with a sex ratio of 2 (in literature: 65.5–70.8% of men) (Wachtel *et al*, 2008; Bergquist *et al*, 2016; Connell *et al*, 2016). There were 26.7% of associated cirrhosis, as frequently described, two patients had positive HBV serology and two others HCV serology, and one had a co-infection, with two patients without underlying cirrhosis. In the literature, the prevalence of HBV or HCV serology in cHCC-ICC population is heterogenous ranging from 17 to 58% and from 0 to 75%, respectively, with higher rates in Asian population (Jarnagin *et al*, 2002; Lee *et al*, 2006; Tang *et al*, 2006; Chok *et al*, 2009; Fowler *et al*, 2013; Bergquist *et al*, 2016).

There were no differences of outcomes depending on the treatment, GEMOX, GEMCIS, GEMOX-bevacizumab, but no conclusion can be made due to our study design and its small sample size. Only three patients were treated with GEMCIS, making comparisons between GEMOX or GEMCIS regimen not relevant and only nine patients had bevacizumab in association. There are few published data concerning combinations of gemcitabine and oxaliplatin for the treatment of HCC. Recently, a large retrospective multicentre study included 204 patients with advanced HCC treated with GEMOX, reported a 22% response rate, 4.5 months median PFS and 11.0 months median OS (Zaanan *et al*, 2013). In the treatment of advanced HCC, the combination of gemcitabine, oxaliplatin, and bevacizumab has been reported in a phase II study, with 5.3 months median PFS and 9.6 months median OS (Zhu *et al*, 2006). A systematic review investigated gemcitabine/cisplatin and gemcitabine/oxaliplatin combinations in the treatment of biliary tract cancers in 1470 patients. This study suggests that the combination of gemcitabine and cisplatin could have a short survival advantage when cisplatin is administered according to the standard protocol on days 1 and 8 compared with

Table 3. Prognostic factors for overall survival (OS): hazard-ratio (HR), CI 95%, and P-value in univariate and multivariate analysis.

Variables	Univariate analysis			Multivariate analysis		
	HR	CI 95%	P-value	HR	CI 95%	P-value
Female sex	0.50	(0.21–1.37)	0.20	0.78	(0.24–2.56)	0.69
Synchronous metastases	1.51	(0.63–4.31)	0.34	2.21	(0.74–6.62)	0.16
Serum bilirubin level $\geq 30 \mu\text{mol l}^{-1}$	3.66	(2.50–34.03)	0.002	10.23	(2.51–41.70)	0.001
Positive serology for HBV and/or HCV	2.18	(0.69–12.40)	0.15	6.89	(1.47–32.20)	0.014
Age ≥ 75 years	0.62	(0.24–1.58)	0.34			
Underlying cirrhosis	1.22	(0.45–3.39)	0.68			
Chronic alcoholism	2.13	(0.59–15.84)	0.21			
Obesity	0.71	(0.24–2.20)	0.58			
Diabetes	0.81	(0.29–2.18)	0.66			
CA 19–9 $\geq 80 \text{ IU ml}^{-1}$	1.62	(0.64–4.89)	0.29			

Statistically significant results are shown in bold ($P < 0.05$).

GEMOX regimen administered every 2 weeks. However, this regimen was associated with increased side effects and toxicities: asthenia, diarrhoea, haematological toxicity, and notably hepatotoxicity, which may not be devoid of consequence in cirrhotic patients (Fiteni *et al*, 2014). The combination of gemcitabine, oxaliplatin, and bevacizumab has also been investigated in advanced biliary-tract cancers with 7 months median PFS and 21.7 months median OS (Zhu *et al*, 2010). According to these data and our results, the combination of gemcitabine and platinum-based chemotherapy could be a promising regimen for the treatment of cHCC-ICC. There are no data reporting the combination of gemcitabine plus platinum chemotherapy with sorafenib, the strategy that could also be investigated, with a potential activity on both components of cHCC-ICC.

Elevated serum bilirubin was identified as an independent significant poor prognostic factor for OS and PFS. This is an identified poor prognostic factor for both, ICC and HCC (Paik *et al*, 2009; Vienne *et al*, 2010). The small size of our population does not allow us to differentiate the etiology of the jaundice (liver dysfunction or biliary obstruction) as independent prognostic factors. Positive serology for HBV and/or HCV was another independent significant poor prognostic factor for OS and PFS. There are few data from prospective studies concerning the prognostic impact of HBV and/or HCV in advanced HCC or ICC, but HBV infection was a poor prognostic factor in patients with HCC in the Asia-Pacific study (Cheng *et al*, 2009).

Our study has some limitations. First, it is a retrospective study, with limited data about the toxicity of the treatments. There are some missing data (biological markers) and four patients were lost to follow-up (13.3%). We included the patients with histologically documented cHCC-ICC, but also those with HCC or ICC histology but discordant contrast-enhanced CT scan findings associated with discordant tumour marker levels. These criteria to identify cHCC-ICC have been previously proposed (Maximin *et al*, 2014) and are important to consider as biopsies often lead to a misdiagnosis. In a series of 23 resected cHCC-ICC, all the tumours were misdiagnosed by the pre-operative histology, 20 considered as HCC, three as ICC (Tagushi *et al*, 1996). The combination of elevated serum tumour markers and enhancement patterns on imaging should strongly suggest the diagnosis of cHCC-ICC in the following circumstances: imaging features of both ICC and HCC,

regardless of marker levels; elevation of both αFP and CA 19–9, regardless of imaging appearance; or discordance between imaging and tumour marker elevation (typical HCC enhancement pattern with elevated CA 19–9 or typical ICC enhancement pattern with elevated αFP) (Maximin *et al*, 2014). The combination of histological, radiological and biological criteria is important to identify cHCC-ICC patients.

In conclusion, these results suggest the efficacy of gemcitabine plus platinum-based chemotherapy in the first-line treatment for unresectable cHCC-ICC with good performance status. Prospective trials are needed but challenging because of scarcity of these tumours.

ACKNOWLEDGEMENTS

We sincerely acknowledge the contribution of our colleagues from all participating hospitals for helping us to collect their data, especially Vanessa Le Berre (clinical research associate in Poitiers).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Allen RA, Lisa JR (1949) Combined liver cell and bile duct carcinoma. *Am J Pathol* **25**(4): 647–655.
- Bergquist JR, Groeschl RT, Ivanics T, Shubert CR, Habermann EB, Kendrick ML, Farnell MB, Nagorney DM, Truty MJ, Smoot RL (2016) Mixed hepatocellular and cholangiocarcinoma: a rare tumor with a mix of parent phenotypic characteristics. *HPB* **18**: 886–889.
- Cheng AL, Kang YK, Chen Z, Tsao CJ, Qin S, Kim JS, Luo R, Feng J, Ye S, Yang TS, Xu J, Sun Y, Liang H, Liu J, Wang J, Tak WY, Pan H, Burock K, Zou J, Voliotis D, Guan Z (2009) Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol* **10**: 25–34.
- Chok KS, Ng KK, Cheung TT, Yuen WK, Poon RT, Lo CM, Fan ST (2009) An update on long term outcome of curative hepatic resection for hepatocholangiocarcinoma. *World J Surg* **33**: 1916–1921.
- Connell LC, Harding JJ, Shia J, Abou-Alfa GK (2016) Combined intrahepatic cholangiocarcinoma and hepatocellular carcinoma. *Chin Clin Oncol* **5**: 66.
- Fiteni F, Nguyen T, Vernerey D, Paillard MJ, Kim S, Demarchi M, Fein F, Borg C, Bonnetain F, Pivot X (2014) Cisplatin/gemcitabine or oxaliplatin/gemcitabine in the treatment of advanced biliary tract cancer: a systematic review. *Cancer Med* **3**(6): 1502–1511.
- Fowler KJ, Sheybani A, Parker 3rd RA, Doherty S, M Brunt E, Chapman WC, Menias CO (2013) Combined hepatocellular and cholangiocarcinoma (biphenotypic) tumors: imaging features and diagnostic accuracy of contrast enhanced CT and MRI. *AJR Am J Roentgenol* **201**: 332.
- Garancini M, Goffredo P, Pagni F, Romano F, Roman S, Sosa JA, Giardini V (2014) Combined hepatocellular-cholangiocarcinoma: a population-level analysis of an uncommon primary liver tumor. *Liver Transplant* **20**: 952–959.
- Groeschl RT, Turaga KK, Gamblin TC (2013) Transplantation versus resection for patients with combined hepatocellular carcinomacholangiocarcinoma. *J Surg Oncol* **107**: 608–612.
- Jarnagin WR, Weber S, Tickoo SK, Koea JB, Obiekwe S, Fong Y, DeMatteo RP, Blumgart LH, Klimstra D (2002) Combined hepatocellular and cholangiocarcinoma: demographic, clinical, and prognostic factors. *Cancer* **94**: 2040–2046.
- Kim H, Park C, Han KH, Choi J, Kim YB, Kim JK, Park YN (2004) Primary liver carcinoma of intermediate (hepatocyte–cholangiocyte) phenotype. *J Hepatol* **40**: 298–304.
- Kim JH, Yoon HK, Ko GY, Gwon DI, Jang CS, Song HY, Shin JH, Sung KB (2010) Nonresectable combined hepatocellular carcinoma and

- cholangiocarcinoma: analysis of the response and prognostic factors after transcatheter arterial chemoembolization. *Radiology* **255**: 270–277.
- Lee WS, Lee KW, Heo JS, Kim SJ, Choi SH, Kim YI, Joh JW (2006) Comparison of combined hepatocellular and cholangiocarcinoma with hepatocellular carcinoma and intrahepatic cholangiocarcinoma. *Surg Today* **36**: 892–897.
- Li R, Yang D, Tang CL, Cai P, Ma KS, Ding SY, Zhang XH, Guo DY, Yan XC (2016) Combined hepatocellular carcinoma and cholangiocarcinoma (biphenotypic) tumors: clinical characteristics, imaging features of contrast-enhanced ultrasound and computed tomography. *BMC Cancer* **16**: 158.
- Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, de Oliveira AC, Santoro A, Raoul JL, Forner A, Schwartz M, Porta C, Zeuzem S, Bolondi L, Greden TF, Galle PR, Seitz JF, Borbath I, Häussinger D, Giannaris T, Shan M, Moscovici M, Voliotis D, Bruix J, SHARP Investigators Study Group (2008) Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* **359**(4): 378–390.
- Maximin S, Ganeshan DM, Shanbhogue AK, Dighe MK, Yeh MM, Kolokythas O, Bhargava P, Lalwani N (2014) Current update on combined hepatocellular- cholangiocarcinoma. *Eur J Radiol Open* **1**: 40–48.
- Paik WH, Park YS, Hwang JH, Lee SH, Yoon CJ, Kang SG, Lee JK, Ryu JK, Kim YT, Yoon YB (2009) Palliative treatment with self-expandable metallic stents in patients with advanced type III or IV hilar cholangiocarcinoma: a percutaneous versus endoscopic approach. *Gastrointest Endosc* **69**(1): 55–62.
- Panjala C, Senecal DL, Bridges MD, Kim GP, Nakhleh RE, Nguyen JH, Harnois DM (2010) The diagnostic conundrum and liver transplantation outcome for combined hepatocellular-cholangiocarcinoma. *Am J Transplant* **10**: 1263–1267.
- Park YH, Hwang S, Ahn CS, Kim KH, Moon DB, Ha TY, Song GW, Jung DH, Park GC, Namgoong JM, Park CS, Park HW, Kang SH, Jung BH, Lee SG (2013) Long-term outcome of liver transplantation for combined hepatocellular carcinoma and cholangiocarcinoma. *Transplant Proc* **45**: 3038–3040.
- Piscaglia AC, Shupe TD, Pani G, Tesori V, Gasbarrini A, Petersen BE (2009) Establishment of cancer cell lines from rat hepatocholangiocarcinoma and assessment of the role of granulocyte-colony stimulating factor and hepatocyte growth factor in their growth, motility and survival. *J Hepatol* **51**: 77–92.
- Sapisochin G, Fidelman N, Roberts JP, Yao FY (2011) Mixed hepatocellular cholangiocarcinoma and intrahepatic cholangiocarcinoma in patients undergoing transplantation for hepatocellular carcinoma. *Liver Transplant* **17**(8): 934–942.
- Tagushi J, Nakashima O, Tanaka M, Hisaka T, Takazawa T, Kojiro M (1996) A clinicopathological study on combined hepatocellular and cholangiocarcinoma. *J Gastroenterol Hepatol* **11**: 758–764.
- Tang D, Nagano H, Nakamura M, Wada H, Marubashi S, Miyamoto A, Takeda Y, Umeshita K, Dono K, Monden M (2006) Clinical and pathological features of Allen's type C classification of resected combined hepatocellular and cholangiocarcinoma: a comparative study with hepatocellular carcinoma and cholangiocellular carcinoma. *J Gastrointest Surg* **10**: 987–998.
- Theise ND, Nakashima O, Park YN, Nakanuma Y (2010) Combined hepatocellular-cholangiocarcinoma. In: *Who classification of tumours of the digestive system*, Bosman FT, Carneiro F, Hruban RH, Theise ND (eds) 4th edn. 225–227. IARC: Lyon, France.
- Valle J, Wasan H, Palmer DH, Cunningham D, Anthony A, Maraveyas A, Madhusudan S, Iveson T, Hughes S, Pereira SP, Roughton M, Bridgewater J, ABC-02 Trial Investigators (2010) Cisplatin plus gemcitabine versus gemcitabine for biliary tract cancer. *N Engl J Med* **362**(14): 1273–1281.
- Vienne A, Hobeika E, Gouya H, Lapidus N, Fritsch J, Choury AD, Chryssostalis A, Gaudric M, Pelletier G, Buffet C, Chaussade S, Prat F (2010) Prediction of drainage effectiveness during endoscopic stenting of malignant hilar strictures: the role of liver volume assessment. *Gastrointest Endosc* **72**(4): 728–735.
- Vilchez V, Shah MB, Daily MF, Pena L, Tzeng CW, Davenport D, Hosein PJ, Gedaly R, Maynard E (2016) Long-term outcome of patients undergoing liver transplantation for mixed hepatocellular carcinoma and cholangiocarcinoma: an analysis of the UNOS database. *HPB* **18**: 29–34.
- Wachtel MS, Zhang Y, Xu T, Chiriva-Internati M, Frezza EE (2008) Combined hepatocellular cholangiocarcinomas; analysis of a large database. *Clin Med Pathol* **1**: 43–47.
- Wang J, Wang F, Kessinger A (2010) Outcome of combined hepatocellular and cholangiocarcinoma of the liver. *J Oncol* **1**–7.
- Zaanan A, Williet N, Hebbar M, Dabakuyo TS, Fartoux L, Mansourbakht T, Dubreuil O, Rosmorduc O, Cattani S, Bonnetain F, Boige V, Taïeb J (2013) Gemcitabine plus oxaliplatin in advanced hepatocellular carcinoma: a large multicenter AGE0 study. *J Hepatol* **58**(1): 81–88.
- Zhu AX, Blaszkowsky LS, Ryan DP, Clark JW, Muzikansky A, Horgan K, Sheehan S, Hale KE, Enzinger PC, Bhargava P, Stuart K (2006) Phase II study of gemcitabine and oxaliplatin in combination with bevacizumab in patients with advanced hepatocellular carcinoma. *J Clin Oncol* **24**(12): 1898–1903.
- Zhu AX, Meyerhardt JA, Blaszkowsky LS, Kambadakone AR, Muzikansky A, Zheng H, Clark JW, Abrams TA, Chan JA, Enzinger PC, Bhargava P, Kwak EL, Allen JN, Jain SR, Stuart K, Horgan K, Sheehan S, Fuchs CS, Ryan DP, Sahani DV (2010) Efficacy and safety of gemcitabine, oxaliplatin, and bevacizumab in advanced biliary-tract cancers and correlation of changes in 18-fluorodeoxyglucose PET with clinical outcome: a phase 2 study. *Lancet Oncol* **11**(1): 48–54.

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 4.0 Unported License.