Keywords: germ-cell tumours; systemic immune-inflammation index; PD-L1; prognostic factor; immunity

Systemic immune-inflammation index in germ-cell tumours

Michal Chovanec^{1,2,3}, Zuzana Cierna⁴, Viera Miskovska⁵, Katarina Machalekova⁶, Katarina Kalavska^{2,3,7}, Katarina Rejlekova^{1,2}, Daniela Svetlovska^{3,8}, Dusan Macak⁹, Stanislav Spanik^{5,10}, Karol Kajo⁶, Pavel Babal^{4,11}, Ugo De Giorgi¹², Michal Mego^{*1,2,3} and Jozef Mardiak^{1,2}

¹Second Department of Oncology, Faculty of Medicine, Comenius University, Klenova 1, Bratislava 833 10, Slovakia; ²Department of Oncology, National Cancer Institute, Klenova 1, Bratislava 833 10, Slovakia; ³Translational Research Unit, Faculty of Medicine, Comenius University, Klenova 1, Bratislava 833 10, Slovakia; ⁴Department of Pathology, Faculty of Medicine, Comenius University, Sasinkova 4, Bratislava 811 08, Slovakia; ⁵First Department of Oncology, Faculty of Medicine, Comenius University, Kollarska 12, Bratislava 812 50, Slovakia; ⁶Department of Pathology, St Elisabeth Cancer Institute, Heydukova 10, Bratislava 812 50, Slovakia; ⁷Department of Molecular Oncology, Cancer Research Institute, Slovak Academy of Sciences, Dubravska cesta 9, Bratislava 845 05, Slovakia; ⁸Department of Clinical Trials, National Cancer Institute, Klenova 1, Bratislava 833 10, Slovakia; ⁹Department of Pathology, National Cancer Institute, Klenova 1, Bratislava 833 10, Slovakia; ¹⁰Department of Oncology, St Elisabeth Cancer Institute, Heydukova 10, Bratislava 812 50, Slovakia; ¹¹Faculty Hospital with Policlinics Skalica, a.s., Koreszkova 936/7, Skalica 909 01, Slovakia and ¹²Department of Oncology, Istituto Scientifico Q1 Romagnolo Q2 per lo Studio e la Cura dei Tumori (IRST) IRCCS, Via Piero Maroncelli 40, Meldola 470 14, Italy

Background: We evaluated systemic immune-inflammation index (SII) and its association with patient outcome in germ-cell tumours (GCTs).

Methods: Two independent cohorts of patients were analysed; the discovery set ($n = 171$) from a single institution and the validation set ($n = 181$) previously included in a study evaluating PD-L1 in GCTs. The SII was calculated using platelet (P), neutrophil (N) and lymphocyte (L) counts before chemotherapy and correlated with survival using regression analyses and Kaplan–Meier method.

Results: In the discovery cohort, the SII was associated with poor risk clinical features. Patients with low SII had significantly longer progression-free survival (HR = 0.22, 95% CI 0.12–0.41, $P < 0.001$) and overall survival (OS) (HR = 0.16, 95% CI 0.08–0.32, $P < 0.001$) compared to high SII. This index was independent of International Germ Cell Cancer Collaborative Group criteria in multivariable Cox regression analysis for OS and was validated in an independent cohort. When combining PD-L1 expression on tumour infiltrating lymphocytes (TILs) and SII, we identified three distinctive prognostic groups.

Conclusions: High SII was associated with poor outcome in GCTs. Combination of PD-L1 positive TILs and SII could further refine prognosis in GCTs.

Immune mechanisms have a significant role in antitumour response and cancer development (Mantovani *et al*, 2008). Testicular germ cell tumours (GCTs) have an exceptional sensitivity to platinum-based chemotherapy (Einhorn, 1979, 1990; Kondagunta *et al*, 2005; Mardiak *et al*, 2005; Mead *et al*, 2005). However, patients who fail front-line and salvage

*Correspondence: Dr M Mego; E-mail: misomego@gmail.com

Note: this study was presented on 3–7 June 2017 at the American Society of Clinical Oncology (ASCO) annual meeting, Chicago, IL, USA, as publication-only abstract no. e16042.

Received 20 October 2017; revised 22 November 2017; accepted 23 November 2017; published online 27 February 2018

© 2018 Cancer Research UK. All rights reserved 0007–0920/18

chemotherapy are rendered incurable (Motzer, 2000; Einhorn *et al*, 2007). Scientific uncertainty regarding mechanisms of resistance to chemotherapy leads to further research in an effort to understand treatment failure in this subset of GCT patients (Romano *et al*, 2016; Sestakova *et al*, 2016; Albany *et al*, 2017). Numerous studies confirmed the efficacy of the check-point inhibition in various types of malignancies in recent years (Topalian *et al*, 2012; Herbst *et al*, 2014; Powles *et al*, 2014; Ansell *et al*, 2015; Garon *et al*, 2015). Furthermore, the expression of programmed death receptor 1 ligand (PD-L1) on tumour cells and tumour infiltrating lymphocytes (TILs) has shown a significant prognostic power in our series of patients (Cierna *et al*, 2016; Chovanec *et al*, 2017). The inflammatory tumour microenvironment (TME) has many roles in tumour progression and metastasis. Simple blood test such as complete blood count (CBC) can identify immune-inflammatory elements (neutrophils, lymphocytes and platelets) that might shed light on the inflammatory TME (Mantovani *et al*, 2008; Lippitz, 2013). Immune inflammatory cells have proven to be prognostic in several types of cancer, including urothelial, colorectal, renal cell cancer and mesothelioma (Kishi *et al*, 2009; Kao *et al*, 2010; Santoni *et al*, 2013; Hu *et al*, 2015; Rossi *et al*, 2015). Neutrophil to lymphocyte ratio was previously developed as a prognostic tool and was widely tested in other solid tumours (Xiao *et al*, 2013; Zheng *et al*, 2013; Balta *et al*, 2014). Additional value of platelets was suggested, as platelets were shown to protect circulating tumour cells from shear stress during circulation, induce epithelial-mesenchymal transition and promote tumour extravasation to metastatic sites (Labelle *et al*, 2011; Placke *et al*, 2012; Schumacher *et al*, 2013). Neutrophils, lymphocytes, and platelets have been recently used in a joined tool, a systemic immune-inflammation index (SII), to provide prognostic information in patients with malignant tumours (Hu *et al*, 2014; Lolli *et al*, 2016a). It was established that SII provides a more powerful tool combining three independent prognostic factors compared to platelets, neutrophil to lymphocytes- or neutrophil to platelets-based tools in cancer (Hong *et al*, 2015; Wang *et al*, 2017; Yu *et al*, 2017). High SII reflects pro-inflammatory activity, which was linked to progression, metastasis and poor outcome in cancer (Seruga *et al*, 2008; Cools-Lartigue *et al*, 2013). SII was also associated with higher counts of circulating tumour cells (Hu *et al*, 2014; Zheng *et al*, 2017). In this study, we evaluated the prognostic value of SII in GCTs and correlated with PD-L1 expression on tumour cells and TILs.

PATIENTS AND METHODS

This retrospective translational study included a discovery set (DS) of 171 patients with GCTs treated from 1999 to 2015 in the National Cancer Institute in Slovakia, with available CBC before systemic platinum-based chemotherapy and sufficient follow-up clinical data. A validation set (VS) consisted of 181 patients included in our previous translational trial of 240 patients evaluating a prognostic significance of PD-L1 on TILs in GCTs, for whom baseline CBC data and sufficient follow-up clinical data were available (Chovanec *et al*, 2017). In contrast, the current study aimed to evaluate the prognostic significance of SII and explore its' associations with PD-L1 expression in GCTs. Clinical data were recorded and compared with SII and PD-L1 expression on TILs. The Institutional Review Board approved this study and a waiver of consent form for patients was granted.

Systemic immune-inflammation index. The SII is an index based on platelets (*P*), neutrophils (*N*) and lymphocytes (*L*) counts. It was calculated using the following formula: $SII = P \times N/L$ as defined previously (Hu *et al*, 2014). The median value was used as

the cutoff value of SII, which was then dichotomised into low (below median) and high (above median) categories.

Tumour pathology, tissue microarray construction and immunohistochemical staining for PD-L1. The present study assessed the association of SII with the PD-L1 expression, which was evaluated in 181 of 240 patients in our previous translational study (Chovanec *et al*, 2017). Tumour specimens collected before the administration of systemic therapy were reviewed by two pathologists associated with the study and PD-L1 expression was evaluated in tumour and on TILs as described previously (Chovanec *et al*, 2017). Germ-cell tumours were identified according to the World Health Organisation criteria (Moch *et al*, 2016). Tissue microarray construction and immunohistochemical staining with rabbit anti-PD-L1 monoclonal antibody was described in detail previously (Chovanec *et al*, 2017). Tumour infiltrating lymphocytes were identified with haematoxylin-eosin staining according to the typical morphology. Tumour cells and TILs with PD-L1 expressions were scored by a weighted histoscore (HS), which accounts for both the extent of cell staining (*S*) and the staining intensity (*I*) (Kirkegaard *et al*, 2006). Positively staining cells were estimated on a scale from 0 to 100%. Subsequently, a score from 0 to 3 (0 = no staining; 1 = weak; 2 = intermediate; and 3 = strong staining) was assigned to describe the average intensity of positively staining cells. The HS was then calculated by following formula: $S \times I$; to yield a scale from 0 to 300. On the basis of the HS, a PD-L1 expression was graded as low (0–150) or high (160–300) as we described previously (Mego *et al*, 2013). The highest PD-L1 expression among GCT subtypes was chosen for mixed GCTs.

Statistical analysis. A Shapiro–Wilk test have shown a significant difference from the normal distribution of PD-L1 HS, therefore non-parametric tests were used for analyses. Differences in distributions of PD-L1 expression between two groups of patients were analysed using the Mann–Whitney *U* test. For analyses of associations between the SII and PD-L1 expression, a one-way analysis of variance was used when using PD-L1 as a continuous variable. We used Fisher's exact test to assess the associations between SII and PD-L1 when used as categorical variables.

A median follow-up time was identified as a median time duration since the diagnosis to the time of the last follow-up. Progression-free survival (PFS) was calculated from the date of orchiectomy or tumour biopsy to the date of progression or death or the date of the last adequate follow-up. Overall survival (OS) was calculated from the date of diagnosis to the date of death or last follow-up. We performed a Kaplan–Meier analysis to estimate PFS and OS using a product limit method and we subsequently compared the results by the log-rank test. A multivariable analysis was performed using Cox proportional hazards model for PFS and OS to assess the differences in prognosis on the basis of SII and PD-L1 expression and a prognosis according to International Germ Cell Cancer Collaborative Group (IGCCCG) criteria (International Germ Cell Cancer Collaborative Group, 1997). All reported *P*-values were two-sided. A *P*-value <0.05 was considered as significant. Statistical analyses were performed using NCSS 10 software (NCSS, 2015, LLC. Kaysville, UT, USA, ncss.com/software/ncss).

RESULTS

Patients' characteristics from discovery and VSs are shown in Table 1. Majority of patients in the discovery and the VS had a non-seminoma histology. A testicular tumour was the most common primary site and more than half of patients were in a good risk category according to the IGCCCG criteria. The median follow-up in the DS was 49 months (0–170) for all patients and 49

months (7–179) for patients still alive. During the follow-up, 42 (25%) patients experienced a disease progression and 34 (20%) patients have died. The estimated 5-year PFS and OS was 75% (95% confidence interval (CI) 68–81%) and 78% (95% CI 73–86%), respectively. The median time of follow-up in the VS was 85 months (0–189) for all patients and 90 months (22–189) for patients still alive. During the follow-up, 42 (25%) of these patients experienced a disease progression and 34 (20%) have died. The estimated 5-year PFS and OS was 89% (95% CI 84–9%) and 91% (95% CI 87–95%), respectively. We attribute the difference in 5-year survival between the DS and the VS to the higher proportion of IGCCCG good-risk patients and lower proportion of IGCCCG poor-risk patients in the VS (Table 1).

Association between the SII and patient/tumour characteristics. We found strongly significant correlations between the SII and poor patients' characteristics in both cohorts (Table 2). Poor and intermediate risk IGCCCG categories and multiple metastatic sites were associated with the high SII in both groups (all $P < 0.001$). Bulky retroperitoneal disease, liver or other non-visceral pulmonary metastases (NPVM) were also significantly associated with the high SII (all $P < 0.001$). Moreover, the high SII was also associated with high tumour markers (both $P < 0.001$). The SII did not significantly differ between seminomas and non-seminomas, although it was significantly higher in patients with extragonadal primary in the DS but not in the VS ($P < 0.001$ vs 0.238).

A prognostic role of the SII. Median SII in DS was 1003. Patients with low SII ($SII < 1003$) had a significantly longer PFS (HR = 0.22, 95% CI 0.12–0.41, $P < 0.001$) (Figure 1A) and OS (HR = 0.16, 95% CI 0.08–0.32, $P < 0.001$) (Figure 1B) opposite to patients with high SII ($SII \geq 1003$).

A model with median obtained from the discovery data set was tested in an independent VS as defined above. This analysis confirmed prognostic value of SII in GCTs. Patients within the VS, who had a low SII, calculated with a median obtained from the DS had a significantly longer PFS (HR = 0.30, 95% CI 0.11–0.81, $P = 0.004$) and OS (HR = 0.15, 95% CI 0.05–0.47, $P < 0.001$) (Figure 2) as opposed to patients with the high SII.

Survival analysis of both study cohorts have reported significantly longer PFS (HR = 0.22, 95% CI 0.13–0.37, $P < 0.0001$) and OS (HR = 0.14, 95% CI 0.08–0.25, $P < 0.0001$) for patients with low SII compared to patients with high SII (Supplementary Figure 2). A multivariable Cox regression analysis has shown that SII was prognostic independently of IGCCCG for OS, but not for PFS when we compared IGCCCG poor vs good/intermediate-risk patients (Table 3). When we performed the multivariable Cox regression analysis with three IGCCCG categories, the analysis lost the statistical significance for OS (data not shown).

The association of the SII and PD-L1 expression. No statistically significant correlation between the SII and PD-L1 expression on tumour or TILs in the VS was observed. The mean HS for PD-L1 on TILs was 107.3 (95% CI 89.0–125.5) in patients with low SII, compared to 89.1 (95% CI 59.0–119.1) in patients with high SII ($P = 0.376$). Similarly, the mean HS for PD-L1 on tumour cells in patients with low vs high SII was 79.5 (95% CI 66.2–92.8) vs 61.2 (95% CI 41.6–80.8) ($P = 0.199$).

A Fisher's exact test of the SII and PD-L1 expression on tumour or TILs also reported no significant correlations ($P = 0.510$ and $P = 0.484$, respectively).

A combined prognostic role of the SII and PD-L1 expressing TILs. Previously, we have shown the prognostic value of PD-L1 expressing TILs in GCTs (16). In the subsequent analysis, we assessed a combined prognostic value of SII and PDL-1 on TILs within the VS. The analysis identified three prognostic groups of patients. The best prognosis was seen in patients who had a high expression of PD-L1 on TILs ($HS \geq 160$) and a low SII ($SII < 1003$)

Table 1. Patient characteristics

	Discovery set		Validation set	
	N = 171	%	N = 181	%
Age (years)				
Median (range)	30 (17–62)		30 (16–67)	
Histology				
Pure seminoma	31	18	33	18
Non-seminoma or mixed GCT	127	74	148	82
N/A	13	8.0		
Primary tumour				
Gonadal	143	84	179	99
Extragenadal	28	16	2	1
IGCCCG risk group				
Good risk	90	53	146	81
Intermediate risk	24	14	19	11
Poor risk	57	33	16	8
Sites of metastases				
Retroperitoneum	146	85	121	67
Mediastinum	45	26	18	10
Lungs	84	49	39	22
Liver	62	35	9	5
Other	29	17	16	9
Non-pulmonary visceral metastases	46	27	8	4
No. of metastatic sites				
0	8	5	55	30
1–2	103	60	101	57
>3	60	35	25	13
Mean (range) of pretreatment markers				
AFP mIU ml ⁻¹	4324 (0–164 946)	998 (0–60 570)		
HCG IU ml ⁻¹	164 474 (0–1 888 840)	10 633 (0–423 338)		
LDH (mkat l ⁻¹)	16 (2–130)	12 (1.97–81)		
Abbreviations: AFP = alpha-fetoprotein; GCT = germ-cell tumour; HCG = human chorionic gonadotropin; IGCCCG = International Germ Cell Cancer Collaborative Group; LDH = lactate dehydrogenase; N/A = not available.				

with a 5-year PFS and OS of both 100%, while the worst prognosis was seen in patients with a low expression of PD-L1 on TILs ($HS < 160$) and a high SII ($SII \geq 1003$) with a 5-year PFS and OS of 70% and 70%, respectively (HR = 0.29, 95% CI 0.10–0.78; $P < 0.001$ for PFS and HR = 0.13, 95% CI 0.04–0.40; $P < 0.001$ for OS) (Supplementary Figure 1). Patients with SII and PD-L1 on TILs both low or high had similar intermediate prognosis.

DISCUSSION

Immune mechanisms have been associated with the pathogenesis of cancer (Sharma and Allison, 2015). Recent advancements in anticancer treatment with immune therapy unleashing the immunity of the host and driving the anti-tumour response resulted in long-term remissions, and even cure in several malignancies (Brahmer *et al*, 2015; Larkin *et al*, 2015; Motzer *et al*, 2015). Testicular cancer has been traditionally referred to as chemotherapy sensitive and few facts are known about the underlying immune mechanisms in this disease. In this study, we analysed a prognostic value of a SII in two independent retrospective cohorts of patients and its association with PD-L1 expression on TILs in our VS. We found that SII calculated prior to chemotherapy is an indicator of prognosis among GCT patients. We did not observe correlation between SII and PD-L1 expression in tumour cells and/or PD-L1 expressing TILs, but the combination of SII and PD-L1 on TILs have created a robust prognostic tool for clinical outcome in GCTs. Our results suggest that immune processes have a role in the mechanisms of progression in GCTs, however, based on our data we cannot determine whether systemic inflammation as expressed by SII creates a permissive micro-environment that leads to the manifestation of the disease with poor prognostic features or if the SII reflects an aggressive disease.

Table 2. Patient characteristics in association with the SII

Variable	N	Discovery set				Validation set				
		SII				SII				
		Mean	s.e.m.	Median	P-value	N	Mean	s.e.m.	Median	P-value
All patients	171	1664	144	1003	NA	181	1034	98	611	NA
Histology										
Seminoma	31	1259	336	932	0.703	33	966	231	608	0.059
Non-seminoma	127	1686	166	971		148	1050	109	1437	
N/A	13									
Tumour primary										
Primary testicular	143	1487	153	909	<0.001	179	1032	99	600	0.238
Extragenadal	28	2571	347	1627		2	1302	939	1302	
IGCCCG risk group										
Good	90	900	178	644	<0.001	146	744	93	534	<0.001
Intermediate	24	2163	345	1319		19	1440	258	739	
Poor	57	2661	224	1991		16	3203	281	2961	
Number of metastatic sites										
0	8	530	647	467	<0.001	55	548	153	461	<0.001
1–2	103	1437	180	751		101	882	113	611	
≥3	60	2207	236	1610		25	2725	227	1785	
Retroperitoneal LN metastases										
Absent	25	1213	348	709	<0.001	60	718	159	467	<0.001
1–5 cm	53	760	239	585		63	767	155	496	
>5 cm	93	2302	181	2256		55	1605	166	1019	
N/A	0					3				
Mediastinal LN metastases										
Absent	126	1480	165	759	<0.001	163	861	95	569	<0.001
Present	45	2180	276	1571		18	2609	287	1461	
Lung metastases										
Absent	109	1447	178	753	<0.001	142	760	102	522	<0.001
Present	62	2046	236	1464		39	2035	195	1153	
Liver										
Absent	137	1412	154	794	<0.001	172	933	95	580	<0.001
Present	34	2683	310	2411		9	2980	417	3154	
Non-pulmonary visceral metastases										
Absent	125	1357	162	757	<0.001	173	950	96	583	<0.001
Present	46	2500	266	1921		8	2860	448	2952	
S-stage										
0	25	799	339	638	<0.001	82	614	126	495	<0.001
1	65	939	210	660		65	982	141	634	
2	24	2163	346	1320		21	1416	248	743	
3	57	2662	225	1991		13	3338	315	2768	

Abbreviations: IGCCCG = International Germ Cell Cancer Collaborative Group; LN = lymph node; NA = not applicable; SII = systemic immune-inflammation index.

However, the ability to predict the clinical outcome using the host's immune parameters could allow the pre-selection of patients with different prognostic profiles and the consequent planning of tailored treatment. Interestingly, Yuksel *et al* (2016) recently reported a neutrophil-to-lymphocyte ratio as a simple marker predictive of the presence of stage I testicular cancer. However, Bolat *et al*, (2017) evaluated a prognostic significance of pre-orchietomy neutrophil-to-lymphocyte ratio in GCT patients and observed no difference in PFS and cancer-specific survival. A revised version of the IGCCCG classification (28) has been launched in 2016, which is collecting data also on *P*, *N* and *L* at baseline in patients treated with first-line chemotherapy (Collette, 2017). This large series of thousands of cases could contribute to better understanding the impact of these parameters including the SII in GCTs.

A cytokine signalling suggesting pro-inflammatory and immunosuppressive pathways that predicts prognosis in GCTs has been

previously described in our works (Chovanec *et al*, 2015; Svetlovska *et al*, 2017). Pro-inflammatory TME has been reported to correspond with poor prognosis in other malignancies as well (Li *et al*, 2014; Tsai *et al*, 2014, 2017). However, the immune TME and its impact on outcome in patients with GCTs is not entirely clear. The SII in our DS have shown strong correlations with essentially all poor clinical characteristics in GCTs. All correlations with clinical features such as the extra-gonadal primary, bulky retroperitoneal disease, NPVM or elevated tumour markers, were strongly significant. We have demonstrated the prognostic value of SII in GCTs for the first time, evidenced by a significant difference in PFS and OS. These findings were replicated in the independent VS. While all the poor clinical characteristics were significantly associated high SII, similar to the data from the DS, one exception was seen. Only two patients were categorised as having primary extra-gonadal tumour within the VS, which contributed to the low statistical power in this tumour characteristic. In a study by Lolli

Figure 1. Kaplan-Meier estimates of probabilities of PFS and OS according to the SII. (A) Estimates of probabilities of PFS in the discovery set of patients with GCTs ($n=171$), HR=0.22, 95% CI 0.12–0.41, $P<0.001$; low SII <1003 ; high SII ≥ 1003 . (B) Estimates of probabilities of OS according to the SII; in the discovery set of patients with GCTs ($n=171$), HR=0.16, 95% CI 0.08–0.32, $P<0.001$; low SII <1003 ; high SII ≥ 1003 . CI = confidence interval; GCTs = germ-cell tumours; HR = hazard ratio; OS = overall survival; PFS = progression-free survival; SII = Systemic immune-inflammation index.

Figure 2. Kaplan-Meier estimates of probabilities of PFS and OS according to the SII. (A) Estimates of probabilities of PFS according to the SII; in the validation set of patients with GCTs ($n=171$), HR=0.30, 95% CI 0.11–0.81, $P=0.004$; low SII <1003 . (B) Estimates of probabilities of OS according to the SII; in the validation set of patients with GCTs ($n=171$), HR=0.15, 95% CI 0.05–0.47, $P<0.001$; low SII <1003 ; high SII ≥ 1003 . CI = confidence interval; GCTs = germ-cell tumours; HR = hazard ratio; OS = overall survival; PFS = progression-free survival; SII = Systemic immune-inflammation index.

et al (2016b), a higher SII was also associated with poor outcome and poor clinical features, such as Gleason score ≥ 8 , visceral metastases or ECOG ≥ 2 , in patients with prostate cancer. Similar data were reported by the same group and others in kidney cancer, oesophageal squamous, hepatocellular, gastric or colorectal carcinoma (Huang *et al*, 2016; Passardi *et al*, 2016; Wang *et al*, 2016; Lolli *et al*, 2016a; Feng *et al*, 2017). With the recent development of immune therapies, PD-1, PD-L1 and CTLA4 have become treatment targets in cancer. Germ-cell tumours express PD-L1 in abundance, as was shown by Fankhauser and our group previously (Fankhauser *et al*, 2015; Cierna *et al*, 2016). In addition, PD-L1 but not PD-1 expression was prognostic when expressed on tumour cells and TILs (Cierna *et al*, 2016; Chovanec *et al*, 2017). A phase II clinical study from Indiana University evaluating an anti-PD-1 agent pembrolizumab failed to prove an efficacy in the treatment of refractory GCTs (Adra *et al*, 2017). Initial results in seven cases with platinum-refractory GCTs treated with anti-PD1 agents (pembrolizumab or nivolumab) after high-dose chemotherapy have been recently reported with possible activity in three patients (Zschabitz *et al*, 2016, 2017) and a single case report provided evidence of ongoing partial remission with marker stabilisation with nivolumab (Chi and Schweizer, 2017). However, predictive markers associated with tumour response have not been reported

and larger prospective clinical trials are suggested. The associations of inflammatory pathways in relation with PD-1/PD-L1 signalling is unknown. Our further investigation of the association of SII and PD-L1 on tumour cells and TILs did not confirm significant correlations among these, suggesting that PD-L1 and SII mirror different aspects of immunity. An analysis of a combined prognostic value of the SII and PD-L1 on TILs in our VS have discovered three groups of patients; (i) with an excellent prognosis (no events reported) if PD-L1 on TILs was high and SII was low, (ii) a poor prognosis if PD-L1 on TILs was low and SII was high, (iii) and an intermediate prognosis if both were low or high. While the survival curves showed smaller differences in patients with better outcomes for OS compared to PFS in this survival analysis, no events were observed in the best prognostic group (100% PFS and 100% OS at 14 years (Supplementary Figure 1). This is an interesting observation, as well established IGCCCG prognostic criteria show 90% 5-year PFS and 97% 5-year OS for good risk patients (1997). Therefore, combined SII and PD-L1 seems to provide more prognostic power compared to the IGCCCG and may be a useful tool for prognosis prediction in the future. We speculate that the favourable prognosis that might be driven by PD-L1 on TILs can be reduced in the pro-inflammatory environment and vice-versa. The underlying mechanism is

Table 3. A multivariable Cox proportional hazards model for PFS and OS assessing differences in outcome on the basis of SII and prognosis according to IGCCCG (N = 352)

Variable	PFS		OS	
	HR (95% CI)	P-value	HR (95% CI)	P-value
SII in GCTs	1.6404	0.1337	2.6502	0.0200
High vs low	(0.8590–3.1324)		(1.1656–6.0256)	
IGCCCG risk group	7.5204	<0.0001	6.8666	<0.0001
Poor vs intermediate/good risk	(4.0501–13.9645)		(3.3483–14.0819)	

Abbreviations: CI = confidence interval; GCT = germ-cell tumour; HR = hazard ratio; IGCCCG = International Germ Cell Cancer Collaborative Group; OS = overall survival; PFS = progression-free survival; SII = systemic immune-inflammation index.

however unclear and more research is needed to provide a detailed explanation.

Our study have some strengths and limitations. The strength of the study is the existence of DS and VS as well as the size of the patient population. Limitations include the retrospective nature of the analysis and under-representation of extragonadal GCTs. Also, we noted some imbalances between the sets that could be responsible for differences in the median of SII. While the most of patients in both cohorts were in IGCCCG good risk, the VS included less poor risk and extragonadal GCTs than the DS, which could explain lower overall SII index in the VS compared to the DS.

In conclusion, this is the first translational study to show a prognostic potency of SII in GCTs. On the basis of the acquired data, we suggest SII and its' combined prognostic value with PD-L1 as an interesting novel finding that requires further larger and prospective study for validation and implementation into clinical practice. Additional research is needed to provide detailed insights into immunobiology of GCTs and uncover possible implications for treatment of this disease.

ACKNOWLEDGEMENTS

We would like to acknowledge our collaborators from departments of pathology in Slovakia: Antol M, Benko J, Danis D, Durcansky D, Fiala P, Galbavy S, Gogora M, Hudcovsky P, Macuch J, Martanovic P, Ondrias F, Plank L and Svajdlar M. We would like to acknowledge Theodore Logan for the critical input, Mrs Daniela Jantekova, from the Population Registry of Slovak Republic for updating patients' database, Mrs Zlatica Pekova for administration support, and Emilia Klincova and Ludovit Gaspar for excellent technical assistance. This work was supported by the Slovak Research and Development Agency under contract APVV-15-0086.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Adra N, Einhorn LH, Althouse SK, Ammakkanavar NR, Musapatika D, Albany C, Vaughn D, Hanna NH (2017) Phase II trial of pembrolizumab in patients with platinum refractory germ cell tumors: a Hoosier Cancer Research Network study GU14-206. *Ann Oncol* (pii): 4555286.

Albany C, Hever-Jardine MP, von Herrmann KM, Yim CY, Tam J, Warzecha JM, Shin L, Bock SE, Curran BS, Chaudhry AS, Kim F,

Sandusky GE, Taverna P, Freemantle SJ, Christensen BC, Einhorn LH, Spinella MJ (2017) Refractory testicular germ cell tumors are highly sensitive to the second generation DNA methylation inhibitor guadecitabine. *Oncotarget* 8(2): 2949–2959.

Ansell SM, Lesokhin AM, Borrello I, Halwani A, Scott EC, Gutierrez M, Schuster SJ, Millenson MM, Cattrly D, Freeman GJ, Rodig SJ, Chapuy B, Ligon AH, Zhu L, Grosso JF, Kim SY, Timmerman JM, Shipp MA, Armand P (2015) PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma. *N Engl J Med* 372(4): 311–319.

Balta S, Demirkol S, Kucuk U, Cakar M, Unlu M (2014) Neutrophil-to-lymphocyte ratio as a novel independent prognostic factor in urothelial carcinoma. *Clin Genitourin Cancer* 12(2): e69–e70.

Bolat D, Aydogdu O, Polat S, Yarimoglu S, Bozkurt IH, Yonguc T, Sen V (2017) Predictive value of preoperative neutrophil-to-lymphocyte ratio on the prognosis of germ cell testicular tumors. *Turk J Urol* 43(1): 55–61.

Brahmer J, Reckamp KL, Baas P, Crino L, Eberhardt WE, Poddubskaya E, Antonia S, Pluzanski A, Vokes EE, Holgado E, Waterhouse D, Ready N, Gainor J, Aren Frontera O, Havel L, Steins M, Garassino MC, Aerts JG, Domine M, Paz-Ares L, Reck M, Baudelet C, Harbison CT, Lestini B, Spigel DR (2015) Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med* 373(2): 123–135.

Chi EA, Schweizer MT (2017) Durable response to immune checkpoint blockade in a platinum-refractory patient with nonseminomatous germ cell tumor. *Clin Genitourin Cancer* 15(5): e855–e857.

Chovanec M, Cierna Z, Miskovska V, Machalekova K, Svetlovskaja D, Kalavska K, Rejlekova K, Spanik S, Kajo K, Babal P, Mardiak J, Mego M (2017) Prognostic role of programmed-death ligand 1 (PD-L1) expressing tumor infiltrating lymphocytes in testicular germ cell tumors. *Oncotarget* 8(13): 21794–21805.

Chovanec M, Mego M, Cholujovala D, Gronesova P, Miskovska V, Sycova-Mila Z, Usakova V, Svetlovskaja D, Bujdak P, Spanik S, Ondrus D, Mardiak J (2015) A cytokine and angiogenic factor (CAF) analysis in plasma in testicular germ cell tumor patients (TGCTs). *J Clin Oncol* 33(15): e15552.

Cierna Z, Mego M, Miskovska V, Machalekova K, Chovanec M, Svetlovskaja D, Hainova K, Rejlekova K, Macak D, Spanik S, Ondrus D, Kajo K, Mardiak J, Babal P (2016) Prognostic value of programmed-death-1 receptor (PD-1) and its ligand 1 (PD-L1) in testicular germ cell tumors. *Ann Oncol* 27(2): 300–305.

Collette L (2017) Update of the international prognostic classification for first line metastatic germ-cell cancers. An international initiative. *Eur J Cancer* 72(suppl 1): S196–S197.

Cools-Lartigue J, Spicer J, McDonald B, Gowing S, Chow S, Giannias B, Bourdeau F, Kubes P, Ferri L (2013) Neutrophil extracellular traps sequester circulating tumor cells and promote metastasis. *J Clin Invest* (pii): 67484.

Einhorn LH (1979) Combination chemotherapy with cis-dichlorodiammineplatinum(II) in disseminated testicular cancer. *Cancer Treat Rep* 63(9–10): 1659–1662.

Einhorn LH (1990) Treatment of testicular cancer: a new and improved model. *J Clin Oncol* 8(11): 1777–1781.

Einhorn LH, Williams SD, Chamness A, Brames MJ, Perkins SM, Abonour R (2007) High-dose chemotherapy and stem-cell rescue for metastatic germ-cell tumors. *N Engl J Med* 357(4): 340–348.

Fankhauser CD, Curioni-Fontecedro A, Allmann V, Beyer J, Tischler V, Sulser T, Moch H, Bode PK (2015) Frequent PD-L1 expression in testicular germ cell tumors. *Br J Cancer* 113(3): 411–413.

Feng JF, Chen S, Yang X (2017) Systemic immune-inflammation index (SII) is a useful prognostic indicator for patients with squamous cell carcinoma of the esophagus. *Medicine* 96(4): e5886.

Garon EB, Rizvi NA, Hui R, Leighl N, Balmanoukian AS, Eder JP, Patnaik A, Aggarwal C, Gubens M, Horn L, Carcereny E, Ahn MJ, Felip E, Lee JS, Hellmann MD, Hamid O, Goldman JW, Soria JC, Dolled-Filhart M, Rutledge RZ, Zhang J, Lunceford JK, Rangwala R, Lubiniecki GM, Roach C, Emancipator K, Gandhi L, Investigators K (2015) Pembrolizumab for the treatment of non-small-cell lung cancer. *N Engl J Med* 372(21): 2018–2028.

Herbst RS, Soria JC, Kowanetz M, Fine GD, Hamid O, Gordon MS, Sosman JA, McDermott DF, Powderly JD, Gettinger SN, Kohrt HE, Horn L, Lawrence DP, Rost S, Leabman M, Xiao Y, Mokatriin A, Koeppen H, Hegde PS, Mellman I, Chen DS, Hodi FS (2014) Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature* 515(7528): 563–567.

- Hong X, Cui B, Wang M, Yang Z, Wang L, Xu Q (2015) Systemic Immune-inflammation Index, Based on Platelet Counts and Neutrophil-Lymphocyte Ratio, Is Useful for Predicting Prognosis in Small Cell Lung Cancer. *Tohoku J Exp Med* **236**(4): 297–304.
- Hu B, Yang XR, Xu Y, Sun YF, Sun C, Guo W, Zhang X, Wang WM, Qiu SJ, Zhou J, Fan J (2014) Systemic immune-inflammation index predicts prognosis of patients after curative resection for hepatocellular carcinoma. *Clin Cancer Res* **20**(23): 6212–6222.
- Hu K, Lou L, Ye J, Zhang S (2015) Prognostic role of the neutrophil-lymphocyte ratio in renal cell carcinoma: a meta-analysis. *BMJ Open* **5**(4): e006040.
- Huang L, Liu S, Lei Y, Wang K, Xu M, Chen Y, Liu B, Chen Y, Fu Q, Zhang P, Qin K, Cai Y, Fu S, Ge S, Yuan X (2016) Systemic immune-inflammation index, thymidine phosphorylase and survival of localized gastric cancer patients after curative resection. *Oncotarget* **7**(28): 44185–44193.
- International Germ Cell Cancer Collaborative Group (1997) International Germ Cell Consensus Classification: a prognostic factor-based staging system for metastatic germ cell cancers. International Germ Cell Cancer Collaborative Group. *J Clin Oncol* **15**(2): 594–603.
- Kao SC, Pavlakos N, Harvie R, Vardy JL, Boyer MJ, van Zandwijk N, Clarke SJ (2010) High blood neutrophil-to-lymphocyte ratio is an indicator of poor prognosis in malignant mesothelioma patients undergoing systemic therapy. *Clin Cancer Res* **16**(23): 5805–5813.
- Kirkegaard T, Edwards J, Tovey S, McGlynn LM, Krishna SN, Mukherjee R, Tam L, Munro AF, Dunne B, Bartlett JM (2006) Observer variation in immunohistochemical analysis of protein expression, time for a change? *Histopathology* **48**(7): 787–794.
- Kishi Y, Kopetz S, Chun YS, Palavecino M, Abdalla EK, Vauthey JN (2009) Blood neutrophil-to-lymphocyte ratio predicts survival in patients with colorectal liver metastases treated with systemic chemotherapy. *Ann Surg Oncol* **16**(3): 614–622.
- Kondagunta GV, Bacik J, Donadio A, Bajorin D, Marion S, Sheinfeld J, Bosl GJ, Motzer RJ (2005) Combination of paclitaxel, ifosfamide, and cisplatin is an effective second-line therapy for patients with relapsed testicular germ cell tumors. *J Clin Oncol* **23**(27): 6549–6555.
- Labelle M, Begum S, Hynes RO (2011) Direct signaling between platelets and cancer cells induces an epithelial-mesenchymal-like transition and promotes metastasis. *Cancer Cell* **20**(5): 576–590.
- Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Cowey CL, Lao CD, Schadendorf D, Dummer R, Smylie M, Rutkowski P, Ferrucci PF, Hill A, Wagstaff J, Carlino MS, Haanen JB, Maio M, Marquez-Rodas I, McArthur GA, Ascierto PA, Long GV, Callahan MK, Postow MA, Grossmann K, Sznol M, Dreno B, Bastholt L, Yang A, Rollin LM, Horak C, Hodi FS, Wolchok JD (2015) Combined nivolumab and ipilimumab or monotherapy in untreated melanoma. *N Engl J Med* **373**(1): 23–34.
- Li G, Wang Z, Ye J, Zhang X, Wu H, Peng J, Song W, Chen C, Cai S, He Y, Xu J (2014) Uncontrolled inflammation induced by AEG-1 promotes gastric cancer and poor prognosis. *Cancer Res* **74**(19): 5541–5552.
- Li TJ, Jiang YM, Hu YF, Huang L, Yu J, Zhao LY, Deng HJ, Mou TY, Liu H, Yang Y, Zhang Q, Li GX (2017) Interleukin-17-producing neutrophils link inflammatory stimuli to disease progression by promoting angiogenesis in gastric cancer. *Clin Cancer Res* **23**(6): 1575–1585.
- Lippitz BE (2013) Cytokine patterns in patients with cancer: a systematic review. *Lancet Oncol* **14**(6): e218–e228.
- Lolli C, Basso U, Derosa L, Scarpi E, Sava T, Santoni M, Crabb SJ, Massari F, Aieta M, Conteduca V, Maruzzo M, La Russa F, Wheeler M, Berardi R, Galli L, De Giorgi U (2016a) Systemic immune-inflammation index predicts the clinical outcome in patients with metastatic renal cell cancer treated with sunitinib. *Oncotarget* **7**(34): 54564–54571.
- Lolli C, Caffo O, Scarpi E, Aieta M, Conteduca V, Maines F, Bianchi E, Massari F, Vecchia A, Chiuri VE, Facchini G, De Giorgi U (2016b) Systemic immune-inflammation index predicts the clinical outcome in patients with mCRPC treated with abiraterone. *Front Pharmacol* **7**: 376.
- Mantovani A, Allavena P, Sica A, Balkwill F (2008) Cancer-related inflammation. *Nature* **454**(7203): 436–444.
- Mardiak J, Salek T, Sycova-Mila Z, Obertova J, Hlavata Z, Mego M, Reckova M, Koza I (2005) Paclitaxel plus ifosfamide and cisplatin in second-line treatment of germ cell tumors: a phase II study. *Neoplasma* **52**(6): 497–501.
- Mead GM, Cullen MH, Huddart R, Harper P, Rustin GJ, Cook PA, Stenning SP, Mason M, Party MRCCTW (2005) A phase II trial of TIP (paclitaxel, ifosfamide and cisplatin) given as second-line (post-BEP) salvage chemotherapy for patients with metastatic germ cell cancer: a medical research council trial. *Br J Cancer* **93**(2): 178–184.
- Mego M, Cierna Z, Svetlovská D, Macak D, Machalekova K, Miskovska V, Chovanec M, Usakova V, Obertova J, Babal P, Mardiak J (2013) PARP expression in germ cell tumours. *J Clin Pathol* **66**(7): 607–612.
- Moch H, Cubilla AL, Humphrey PA, Reuter VE, Ulbright TM (2016) The 2016 WHO classification of tumours of the urinary system and male genital organs-part a: renal, penile, and testicular tumours. *Eur Urol* **70**(1): 93–105.
- Motzer RJ (2000) Paclitaxel (Taxol) combination therapy for resistant germ cell tumors. *Semin Oncol* **27**(1 Suppl 1): 33–35.
- Motzer RJ, Escudier B, McDermott DF, George S, Hammers HJ, Srinivas S, Tykodi SS, Sosman JA, Procopio G, Plimack ER, Castellano D, Choueiri TK, Gurney H, Donskov F, Bono P, Wagstaff J, Gauder TC, Ueda T, Tomita Y, Schutz FA, Kollmannsberger C, Larkin J, Ravaud A, Simon JS, Xu LA, Waxman IM, Sharma P, CheckMate I (2015) Nivolumab versus everolimus in advanced renal-cell carcinoma. *N Engl J Med* **373**(19): 1803–1813.
- Passardi A, Scarpi E, Cavanna L, Dall'Agata M, Tassinari D, Leo S, Bernardini I, Gelsomino F, Tamberi S, Brandes AA, Tenti E, Vespignani R, Frassinetti GL, Amadori D, De Giorgi U (2016) Inflammatory indexes as predictors of prognosis and bevacizumab efficacy in patients with metastatic colorectal cancer. *Oncotarget* **7**(22): 33210–33219.
- Placke T, Salih HR, Kopp HG (2012) G1TR ligand provided by thrombopoietic cells inhibits NK cell antitumor activity. *J Immunol* **189**(1): 154–160.
- Powles T, Eder JP, Fine GD, Braiteh FS, Loriot Y, Cruz C, Bellmunt J, Burris HA, Petrylak DP, Teng SL, Shen X, Boyd Z, Hegde PS, Chen DS, Vogelzang NJ (2014) MPDL3280A (anti-PD-L1) treatment leads to clinical activity in metastatic bladder cancer. *Nature* **515**(7528): 558–562.
- Romano FJ, Rossetti S, Conteduca V, Schepisi G, Cavaliere C, Di Franco R, La Mantia E, Castaldo L, Nocerino F, Ametrano G, Cappuccio F, Malzone G, Montanari M, Vanacore D, Quagliariello V, Piscitelli R, Pepe MF, Berretta M, D'Aniello C, Perdoni S, Muto P, Botti G, Ciliberto G, Veneziani BM, De Falco F, Maiolino P, Caraglia M, Montella M, De Giorgi U, Facchini G (2016) Role of DNA repair machinery and p53 in the testicular germ cell cancer: a review. *Oncotarget* **7**(51): 85641–85649.
- Rossi L, Santoni M, Crabb SJ, Scarpi E, Burattini L, Chau C, Bianchi E, Savini A, Burgio SL, Conti A, Conteduca V, Cascinu S, De Giorgi U (2015) High neutrophil-to-lymphocyte ratio persistent during first-line chemotherapy predicts poor clinical outcome in patients with advanced urothelial cancer. *Ann Surg Oncol* **22**(4): 1377–1384.
- Santoni M, De Giorgi U, Iacovelli R, Conti A, Burattini L, Rossi L, Luca Burgio S, Berardi R, Muzzonigro G, Cortesi E, Amadori D, Cascinu S (2013) Pre-treatment neutrophil-to-lymphocyte ratio may be associated with the outcome in patients treated with everolimus for metastatic renal cell carcinoma. *Br J Cancer* **109**(7): 1755–1759.
- Schumacher D, Strlic B, Sivaraj KK, Wettschurek N, Offermanns S (2013) Platelet-derived nucleotides promote tumor-cell transendothelial migration and metastasis via P2Y2 receptor. *Cancer Cell* **24**(1): 130–137.
- Seruga B, Zhang H, Bernstein LJ, Tannock IF (2008) Cytokines and their relationship to the symptoms and outcome of cancer. *Nat Rev Cancer* **8**(11): 887–899.
- Sestakova Z, Kalavská K, Hurbanova L, Jurkovicova D, Gursky J, Chovanec M, Svetlovská D, Miskovska V, Obertova J, Palacka P, Rejlekova K, Sycova-Mila Z, Cingelova S, Spanik S, Mardiak J, Chovanec M, Mego M (2016) The prognostic value of DNA damage level in peripheral blood lymphocytes of chemotherapy-naïve patients with germ cell cancer. *Oncotarget* **7**(46): 75996–76005.
- Sharma P, Allison JP (2015) The future of immune checkpoint therapy. *Science* **348**(6230): 56–61.
- Svetlovská D, Miskovska V, Cholujova D, Gronesova P, Cingelova S, Chovanec M, Sycova-Mila Z, Obertova J, Palacka P, Rajec J, Kalavská K, Usakova V, Luha J, Ondrus D, Spanik S, Mardiak J, Mego M (2017) Plasma cytokines correlated with disease characteristics, progression-free survival, and overall survival in testicular germ-cell tumor patients. *Clin Genitourin Cancer* **15**(3): 411–416.e2.
- Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF, Powderly JD, Carvajal RD, Sosman JA, Atkins MB, Leming PD, Spigel DR, Antonia SJ, Horn LM, Drake CG, Pardoll DM, Chen L, Sharfman WH, Anders RA, Taube JM, McMiller TL, Xu H, Korman AJ, Jure-Kunkel M, Agrawal S, McDonald D, Kollia GD, Gupta A, Wigginton JM, Sznol M (2012) Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* **366**(26): 2443–2454.

- Tsai CY, Wang CS, Tsai MM, Chi HC, Cheng WL, Tseng YH, Chen CY, Lin CD, Wu JI, Wang LH, Lin KH (2014) Interleukin-32 increases human gastric cancer cell invasion associated with tumor progression and metastasis. *Clin Cancer Res* **20**(9): 2276–2288.
- Wang BL, Tian L, Gao XH, Ma XL, Wu J, Zhang CY, Zhou Y, Guo W, Yang XR (2016) Dynamic change of the systemic immune inflammation index predicts the prognosis of patients with hepatocellular carcinoma after curative resection. *Clin Chem Lab Med* **54**(12): 1963–1969.
- Wang K, Diao F, Ye Z, Zhang X, Zhai E, Ren H, Li T, Wu H, He Y, Cai S, Chen J (2017) Prognostic value of systemic immune-inflammation index in patients with gastric cancer. *Chin J Cancer* **36**(1): 75.
- Xiao GQ, Liu C, Liu DL, Yang JY, Yan LN (2013) Neutrophil-lymphocyte ratio predicts the prognosis of patients with hepatocellular carcinoma after liver transplantation. *World J Gastroenterol* **19**(45): 8398–8407.
- Yu J, Wu X, Yu H, Li S, Mao L, Chi Z, Si L, Sheng X, Cui C, Dai J, Ma M, Tang H, Xu T, Yan J, Kong Y, Guo J (2017) Systemic immune-inflammation index and circulating T-cell immune index predict outcomes in high-risk acral melanoma patients treated with high-dose interferon. *Transl Oncol* **10**(5): 719–725.
- Yuksel OH, Verit A, Sahin A, Urkmez A, Uruc F (2016) White blood cell counts and neutrophil to lymphocyte ratio in the diagnosis of testicular cancer: a simple secondary serum tumor marker. *Int Braz J Urol* **42**(1): 53–59.
- Zheng L, Zou K, Yang C, Chen F, Guo T, Xiong B (2017) Inflammation-based indexes and clinicopathologic features are strong predictive values of preoperative circulating tumor cell detection in gastric cancer patients. *Clin Transl Oncol* **19**(9): 1125–1132.
- Zheng YB, Zhao W, Liu B, Lu LG, He X, Huang JW, Li Y, Hu BS (2013) The blood neutrophil-to-lymphocyte ratio predicts survival in patients with advanced hepatocellular carcinoma receiving sorafenib. *Asian Pac J Cancer Prev* **14**(9): 5527–5531.
- Zschabitz S, Lasitschka F, Hadaschik B, Hofheinz RD, Jentsch-Ullrich K, Gruner M, Jager D, Grulich C (2017) Response to anti-programmed cell death protein-1 antibodies in men treated for platinum refractory germ cell cancer relapsed after high-dose chemotherapy and stem cell transplantation. *Eur J Cancer* **76**: 1–7.
- Zschabitz S, Lasitschka F, Jager D, Grulich C (2016) Activity of immune checkpoint inhibition in platinum refractory germ-cell tumors. *Ann Oncol* **27**(7): 1356–1360.

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 4.0 Unported License.

Supplementary Information accompanies this paper on British Journal of Cancer website (<http://www.nature.com/bjc>)