

Keywords: colorectal; colon; proximal; distal; rectal cancer; physical activity; sedentary behaviours

Physical activity, sedentary behaviour and colorectal cancer risk in the UK Biobank

Jessica S Morris¹, Kathryn E Bradbury², Amanda J Cross¹, Marc J Gunter³ and Neil Murphy^{*,3}

¹Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK; ²Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK and ³Section of Nutrition and Metabolism, International Agency for Research on Cancer, Lyon, France

Background: Observational studies have shown that physical activity levels are inversely, and sedentary behaviours are positively, associated with colorectal cancer risk; however, whether these relationships are consistent across anatomical subsites is uncertain.

Methods: We investigated the associations between colorectal cancer and physical activity (metabolic equivalents (METs)-hours per week), and indicators of sedentary behaviour (television watching time and time spent using computers) among 430 584 men and women enrolled in the UK Biobank. Multivariable hazard ratios (HRs) and 95% confidence intervals (CI) were estimated using Cox proportional hazards models.

Results: After a median follow-up time of 5.6 years, 2391 incident colorectal cancer cases were recorded. High (≥ 60 -MET-hours per week) vs low (< 10 -MET-hours per week) total physical activity was associated with a lower colon cancer risk (HR = 0.84, 95% CI: 0.72–0.98; p -trend = 0.04), with comparable relationships observed for proximal and distal colon tumours, but no association for rectal cancer. Higher levels of television watching time were associated with greater colon cancer risk (HR for ≥ 5 h per day vs ≤ 1 h per day = 1.32, 95% CI: 1.04–1.68; p -trend = 0.007). Time spent using computers was not associated with colorectal cancer risk.

Conclusions: Higher levels of physical activity were associated with lower colon cancer risk, with no heterogeneity by colonic subsite. Sedentary behaviour (television watching) was associated with elevated colon cancer risk.

Colorectal cancer is the third most commonly diagnosed cancer in men and the second in women worldwide (IARC-WHO, 2015). A large body of evidence has shown that greater levels of physical activity are associated with a lower risk of colorectal cancer (WCRF-AICR, 2011; Moore *et al*, 2016). However, colorectal cancers are a heterogeneous collection of tumours with variable molecular characteristics, and it is hypothesised that tumours located at different anatomical subsites may have distinct aetiological risk factors (Wei *et al*, 2004; Limsui *et al*, 2010). For physical activity, generally consistent inverse relationships have been found for colon and rectal cancers, with a recent participant-level pooled analysis, which included 1.44 million participants, reporting that leisure-time physical activity was associated with lower risks of colon cancer and rectal cancer when analysed separately (Moore *et al*, 2016). It is unclear, however, how physical

activity levels are related to colon cancer risk across the proximal and distal anatomical regions as few studies have been of sufficient size to undertake analyses by subsite, and inconsistent results have emerged. For instance, in the European Prospective Investigation into Cancer and Nutrition (EPIC), an inverse relationship was reported only between total physical activity and proximal colon cancer risk, but null results were found for distal colon cancer risk (Friedenreich *et al*, 2006). In contrast, in other large prospective studies, inverse relationships of similar magnitude have been observed across the colonic subsites (Howard *et al*, 2008; Burón Pust *et al*, 2017; Wei *et al*, 2017).

Sedentary behaviours, defined as sitting or reclined posture activities which expend ≤ 1.5 metabolic equivalents (METs) (Owen *et al*, 2000), may also influence colorectal cancer risk. In the National Institutes of Health (NIH)-AARP Diet and Health

*Correspondence: Dr N Murphy; E-mail: murphyn@iarc.fr

Received 20 September 2017; revised 19 December 2017; accepted 20 December 2017; published online 8 March 2018

© 2018 Cancer Research UK. All rights reserved 0007–0920/18

Study, sedentary behaviour (time spent watching television and videos) was associated with a greater colorectal cancer risk (Howard *et al*, 2008). A meta-analysis of case-control and cohort studies reported that total sitting time, television viewing time, and occupational sitting time were all positively associated with colon cancer development (Schmid and Leitzmann, 2014). However, whether this relationship differs within the colon for proximal and distal tumours has been largely unstudied.

In the current analysis, we investigated how total physical activity levels and sedentary behaviours, such as television watching and time spent using computers, are related to colorectal cancer risk in the UK Biobank study. The UK Biobank is a large-scale prospective cohort study including over 500 000 participants. The large number of recorded colorectal cancer cases provides substantial statistical power to investigate relationships across colorectal cancer subsites (colon, proximal colon, distal colon, and rectum) and according to sex and body habitus.

MATERIALS AND METHODS

Study participants. The UK Biobank is a prospective cohort study which aims to investigate the genetic, lifestyle, and environmental causes of a range of diseases (UK-Biobank, 2010; Allen *et al*, 2012). Between 2006 and 2010, 502 656 adults aged between 40 and 69 years (229 182 men and 273 474 women) were recruited. All participants were registered with the UK National Health Service (NHS) and lived within ~25 miles (40 km) of 1 of the 22 study assessment centres. The UK Biobank invited ~9.2 million people to participate through postal invitation with a telephone follow-up, with a response rate of 5.7%. The UK Biobank has approval from the North West Multi-centre Research Ethics Committee, the National Information Governance Board for Health and Social Care in England and Wales, and the Community Health Index Advisory Group in Scotland. In addition, an independent Ethics and Governance Council was formed in 2004 to oversee UK Biobank's continuous adherence to the Ethics and Governance Framework, which were developed for the study (<http://www.uk-biobank.ac.uk/ethics/>). All participants provided written informed consent.

During the baseline recruitment visit, participants were asked to complete a self-administered touchscreen questionnaire, which included questions on socio-demographics (including age, sex, education and Townsend deprivation score), health and medical history, lifestyle exposures (including smoking habits, dietary intakes, and alcohol consumption), early life exposures, and medication use. At the baseline interview, participants also underwent physical measurements, including body weight, height, and waist circumference. Exclusions prior to the onset of analyses included: participants with prevalent cancer at recruitment ($n = 27\,058$) and those with missing self-reported physical activity information and body size measurements ($n = 45\,014$). Our analysis therefore included 430 584 participants (201 225 men and 229 359 women).

Assessment of exposure. The questions on physical activity that were included in the UK Biobank baseline questionnaire were adapted from the International Physical Activity Questionnaire (IPAQ) short form, a validated survey instrument (Craig *et al*, 2003). Questions are listed in Supplementary Table 1 and are available to view on the UK Biobank Web site (<http://biobank.ctsu.ox.ac.uk/crystal/label.cgi?id=100054>). The questions captured the frequency and duration of three different levels of activity (walking, moderate, and vigorous). Participants were asked on how many days in a typical week they engaged in each activity level for 10 min or more. For each level in which an answer of one or more days was given, the participant was then asked how many minutes they usually spent doing that activity on a typical day. Using this

information, METs were used to quantify the intensity of physical activity. The MET values represent the ratio of energy expended per kilogram of body weight per hour to the standard resting metabolic rate (Ainsworth *et al*, 2011). The number of days per week that participants engaged in each level of physical activity was multiplied by the number of minutes spent per day doing that activity. This gave the total number of minutes spent per week engaged in each activity category. Total MET values for each category from the International Physical Activity Questionnaire short form were: 3.3 for walking, 4.0 for moderate physical activity and 8.0 for vigorous physical activity (IPAQ Research Committee, 2005). We report excess METs, which are calculated by subtracting one MET from the value for each activity, and represent the energy expenditure above that of an inactive person (Howley, 2001). Excess MET values were therefore 2.3 for walking, 3.0 for moderate physical activity and 7.0 for vigorous physical activity. Excess MET-hours per week were calculated by multiplying the excess MET value for each activity by the duration of activity in hours per week (IPAQ Research Committee, 2005). In the baseline questionnaire, participants were also asked in a typical day how many hours they spent watching television and using a computer. This information was used as an indicator of sedentary behaviours as television watching and time spent using computers are two of the most wide-spread leisure-time sedentary behaviours in adults (Office for National Statistics, 2006)

Assessment of outcome. Prevalent and incident cancer cases within the UK Biobank cohort were identified through linkage to national cancer registries. Complete follow-up was available through 30th November 2014 for England and Wales and 31st December 2014 for Scotland. Cancer incidence data were coded using the 10th Revision of the International Classification of Diseases (ICD-10). Proximal colon cancers included those found within the caecum, appendix, ascending colon, hepatic flexure, transverse colon, and splenic flexure (C18.0-18.5). Distal colon cancers included those found within the descending (C18.6) and sigmoid (C18.7) colon. Overlapping (C18.8) and unspecified (C18.9) lesions of the colon were included in colon cancers only. Cancer of the rectum included cancers occurring at the recto sigmoid junction (C19) and rectum (C20).

Statistical analysis. Hazard ratios (HRs) and 95% confidence intervals (CIs) were estimated using Cox proportional hazards models. Age was the primary time variable in all models. Time at entry was age at recruitment. Exit time was age at whichever of the following came first: colorectal cancer diagnosis, death, or the last date at which follow-up was considered complete. Models were stratified by age at recruitment in 5-year categories, Townsend deprivation index fifths, and region of the recruitment assessment centre. Deviations from proportionality was assessed using an analysis of Schoenfeld residuals (Schoenfeld, 1982), with no evidence of non-proportionality being detected. Participants were grouped into predefined categories for total physical activity levels (<10, 10-<20, 20-<40, 40-<60, ≥ 60 MET hours per week), television watching time (≤ 1 , 2-3, 4-5, > 5 h per day for sexes combined analyses, and ≤ 1 , 2-3, ≥ 4 h per day for sex-specific analyses), and time spent using computers (none, $> 0-1$, 2-3, ≥ 4 h per day for sexes combined analyses, and none, $> 0-1$, ≥ 2 h per day for sex-specific analyses).

The multivariable models were adjusted for a set of a priori-determined colorectal cancer risk factors, namely height (continuous, cm), smoking status and intensity (never; former; current— < 15 cigarettes per day; current— ≥ 15 cigarettes per day; current - intensity unknown; unknown), alcohol consumption frequency (never; special occasions only; 1-3 times per month; 1-2 times per week; 3-4 times per week; daily or almost daily; unknown), family history of colorectal cancer (no; yes; unknown), prevalent diabetes (no; yes; unknown), regular aspirin/ibuprofen

Table 1. Characteristics of study participants by category of total physical activity (MET-hours per week)

Baseline characteristic	Total physical activity (MET hours per week)				
	< 10	10–<20	20–<40	40–<60	≥ 60
Men					
Colorectal cancer (N)	372	284	317	168	248
Age at recruitment (years) ^a	56.5 (8.0)	56.6 (8.2)	56.5 (8.3)	56.7 (8.3)	56.4 (8.3)
Body mass index (kg m ⁻²) ^a	28.5 (4.6)	27.7 (4.2)	27.5 (3.9)	27.4 (3.9)	27.4 (3.9)
Waist circumference (cm) ^a	99.3 (11.9)	97.0 (11.0)	95.7 (10.6)	95.1 (10.6)	95.0 (10.7)
Height (cm) ^a	175.7 (6.9)	176.0 (6.8)	176.0 (6.8)	175.7 (6.8)	175.1 (6.7)
Smoking status (%)					
Never	48.8	50.6	51.1	50	46.9
Current	13.6	11.3	10.3	10.5	14.2
Alcohol consumption (%)					
Never	7.3	5.7	5.1	5.3	6.1
Daily or almost daily	25.4	26.7	26.3	25.9	24.1
Qualifications (%)					
College/university degree	37.1	40.9	39.5	34.5	21.6
Family history of colorectal cancer (%)					
Yes	11.3	10.9	10.7	11.2	11.5
Aspirin/ibuprofen use (%)					
Yes	30.1	28.3	27.3	28.4	27.5
Red and processed meat (%)					
< 1 occasion per week	6.6	6.9	7.4	8	7.5
≥ 3 occasions per week	25.3	24.5	24	24.9	27.4
Women					
Colorectal cancer (N)	282	237	224	108	151
Age at recruitment (years) ^a	55.9 (7.9)	55.9 (8.0)	55.9 (8.1)	56.3 (8.2)	56.6 (8.1)
Body mass index (kg m ⁻²) ^a	27.9 (5.6)	26.9 (5.0)	26.4 (4.7)	26.2 (4.6)	26.2 (4.6)
Waist circumference (cm) ^a	86.9 (13.2)	84.4 (12.2)	83.0 (11.6)	82.4 (11.5)	82.4 (11.6)
Height (cm) ^a	162.4 (6.4)	162.7 (6.3)	162.8 (6.3)	162.6 (6.2)	162.3 (6.2)
Smoking status (%)					
Never	60.1	60.7	59.9	60	58.6
Current	9.6	8	7.6	7.8	9.4
Alcohol consumption (%)					
Never	10.2	8.4	7.9	8.3	9.8
Daily or almost daily	15.4	16.8	17	17.3	16.3
Qualifications (%)					
College/university degree	31.7	35.5	35.4	32.9	26.5
Family history of colorectal cancer (%)					
Yes	10.6	10.4	10.5	10.7	10.9
Aspirin/ibuprofen use (%)					
Yes	26.4	25.6	24.7	24.9	25.2
Red and processed meat (%)					
< 1 occasion per week	11.4	12.2	13	13.8	14.4
≥ 3 occasions per week	19.4	18.9	18.6	18.8	19.8
Ever menopausal hormone use (%)					
Yes	37	36.6	36.4	37.8	39.6

^aMean and standard deviation.

use (no; yes; unknown), qualifications (none; national exams at age 16 years (CSEs/O-levels/GCSEs or equivalent); vocational qualifications (NVQ/HND/HNC) or optional national exams at ages 17–18 years (A-levels/AS-levels or equivalent); other professional qualifications; college/university degree; unknown, frequency of red and processed meat consumption (<2 times per week; 2–3 times per week; 3–4 times per week; >4 times per week; unknown), and, among women, ever use of hormone replacement therapy (HRT; no, yes, unknown). Further adjustment for previous colorectal cancer screening resulted in virtually unchanged risk estimates, so this variable was not included in the final multi-variable models. As body size/adiposity is potentially on the causal pathway linking physical activity and sedentary activities with colorectal cancer, we also ran all models with and without adjustment for waist circumference. In addition, in sensitivity

analyses, the television watching time and time spent using computers analyses were mutually adjusted for one another.

Trend tests across physical activity categories were calculated by assigning the median value of each measurement and modelling as continuous terms in Cox regression models. For television watching time and time spent using computers, trend tests were calculated by entering the categorical exposure variables into the models as continuous variables. As well as for overall colorectal cancer, analyses were undertaken for colon cancer, proximal colon cancer, distal colon cancer, and rectal cancer for sexes combined and for men and women separately. Heterogeneity of associations by sex and across anatomical cancer subsites was assessed by calculating X^2 statistics.

The physical activity and colorectal cancer associations were further assessed across subgroups of body mass index (BMI; <25,

Table 2. Risk (hazard ratios) of colorectal cancer associated with total physical activity

	Total physical activity (MET hours per week)					p-trend
	<10	10–<20	20–<40	40–<60	≥60	
Colorectal cancer						
Both sexes						
N	654	521	541	276	399	
Multivariable, HR (95% CI)	1	0.99 (0.88–1.11)	0.90 (0.80–1.01)	0.98 (0.85–1.13)	0.92 (0.81–1.04)	0.25
Multivariable plus waist circumference, HR (95% CI)	1	1.01 (0.90–1.13)	0.93 (0.83–1.04)	1.02 (0.88–1.18)	0.96 (0.85–1.09)	0.63
Men						
N	372	284	317	168	248	
Multivariable, HR (95% CI)	1	0.95 (0.81–1.11)	0.91 (0.78–1.06)	1.00 (0.84–1.21)	0.89 (0.76–1.05)	0.26
Multivariable plus waist circumference, HR (95% CI)	1	0.98 (0.84–1.14)	0.95 (0.82–1.11)	1.06 (0.88–1.27)	0.94 (0.80–1.11)	0.65
Women						
N	282	237	224	108	151	
Multivariable, HR (95% CI)	1	1.03 (0.87–1.23)	0.88 (0.74–1.05)	0.94 (0.75–1.17)	0.96 (0.78–1.17)	0.52
Multivariable plus waist circumference, HR (95% CI)	1	1.05 (0.88–1.24)	0.90 (0.75–1.07)	0.96 (0.77–1.20)	0.98 (0.80–1.20)	0.71
Colon cancer						
Both sexes						
N	449	356	323	181	248	
Multivariable, HR (95% CI)	1	0.98 (0.85–1.12)	0.78 (0.67–0.90)	0.93 (0.78–1.11)	0.84 (0.72–0.98)	0.04
Multivariable plus waist circumference, HR (95% CI)	1	1.00 (0.87–1.15)	0.81 (0.70–0.94)	0.98 (0.82–1.16)	0.89 (0.76–1.04)	0.17
Men						
N	241	186	170	97	139	
Multivariable, HR (95% CI)	1	0.96 (0.79–1.16)	0.75 (0.62–0.91)	0.89 (0.70–1.13)	0.77 (0.62–0.95)	0.02
Multivariable plus waist circumference, HR (95% CI)	1	0.99 (0.82–1.20)	0.79 (0.65–0.97)	0.95 (0.75–1.21)	0.83 (0.67–1.02)	0.10
Women						
N	208	170	153	84	109	
Multivariable, HR (95% CI)	1	1.00 (0.81–1.22)	0.81 (0.66–0.99)	0.98 (0.76–1.26)	0.93 (0.74–1.17)	0.58
Multivariable plus waist circumference, HR (95% CI)	1	1.01 (0.83–1.24)	0.83 (0.67–1.02)	1.01 (0.78–1.30)	0.96 (0.76–1.21)	0.78
Proximal colon cancer						
Both sexes						
N	227	182	163	92	120	
Multivariable, HR (95% CI)	1	0.98 (0.81–1.20)	0.78 (0.63–0.95)	0.93 (0.73–1.18)	0.79 (0.63–0.99)	0.04
Multivariable plus waist circumference, HR (95% CI)	1	1.02 (0.84–1.24)	0.81 (0.66–0.99)	0.99 (0.77–1.26)	0.84 (0.67–1.06)	0.15
Men						
N	128	90	86	44	65	
Multivariable, HR (95% CI)	1	0.87 (0.66–1.14)	0.71 (0.54–0.94)	0.76 (0.54–1.07)	0.67 (0.49–0.90)	0.01
Multivariable plus waist circumference, HR (95% CI)	1	0.91 (0.69–1.19)	0.76 (0.58–1.00)	0.82 (0.58–1.16)	0.72 (0.53–0.98)	0.05
Women						
N	99	92	77	48	55	
Multivariable, HR (95% CI)	1	1.13 (0.85–1.50)	0.85 (0.63–1.15)	1.15 (0.81–1.62)	0.95 (0.69–1.33)	0.75
Multivariable plus waist circumference, HR (95% CI)	1	1.15 (0.87–1.54)	0.88 (0.65–1.19)	1.21 (0.85–1.71)	1.01 (0.72–1.41)	0.99
Distal colon cancer						
Both sexes						
N	202	154	143	81	114	
Multivariable, HR (95% CI)	1	0.95 (0.77–1.17)	0.77 (0.62–0.96)	0.94 (0.72–1.22)	0.88 (0.70–1.11)	0.38
Multivariable plus waist circumference, HR (95% CI)	1	0.97 (0.78–1.19)	0.80 (0.64–0.99)	0.97 (0.75–1.26)	0.91 (0.72–1.15)	0.57
Men						
N	104	86	75	48	68	
Multivariable, HR (95% CI)	1	1.04 (0.78–1.38)	0.77 (0.57–1.04)	1.03 (0.73–1.45)	0.89 (0.65–1.22)	0.52
Multivariable plus waist circumference, HR (95% CI)	1	1.06 (0.80–1.42)	0.81 (0.60–1.09)	1.08 (0.76–1.52)	0.94 (0.69–1.29)	0.75
Women						
N	98	68	68	33	46	
Multivariable, HR (95% CI)	1	0.86 (0.63–1.17)	0.77 (0.57–1.06)	0.83 (0.56–1.24)	0.86 (0.60–1.22)	0.48
Multivariable plus waist circumference, HR (95% CI)	1	0.86 (0.63–1.18)	0.78 (0.57–1.07)	0.84 (0.57–1.26)	0.87 (0.61–1.24)	0.55
Rectal cancer						
Both sexes						
N	200	162	214	93	151	
Multivariable, HR (95% CI)	1	1.01 (0.82–1.24)	1.16 (0.95–1.41)	1.09 (0.85–1.39)	1.11 (0.90–1.37)	0.34
Multivariable plus waist circumference, HR (95% CI)	1	1.02 (0.83–1.26)	1.19 (0.98–1.44)	1.12 (0.87–1.43)	1.15 (0.92–1.42)	0.23
Men						
N	133	96	144	69	107	
Multivariable, HR (95% CI)	1	0.90 (0.69–1.17)	1.16 (0.92–1.47)	1.17 (0.87–1.56)	1.07 (0.83–1.38)	0.41
Multivariable plus waist circumference, HR (95% CI)	1	0.92 (0.71–1.20)	1.20 (0.94–1.52)	1.21 (0.90–1.63)	1.11 (0.86–1.45)	0.28

Table 2. (Continued)

	Total physical activity (MET hours per week)					p-trend
	<10	10–<20	20–<40	40–<60	≥60	
Women						
N	67	66	70	24	44	
Multivariable, HR (95% CI)	1	1.21 (0.86–1.71)	1.15 (0.82–1.61)	0.89 (0.56–1.42)	1.18 (0.80–1.72)	0.78
Multivariable plus waist circumference, HR (95% CI)	1	1.22 (0.87–1.72)	1.16 (0.83–1.63)	0.90 (0.56–1.44)	1.19 (0.81–1.75)	0.74

Abbreviations: CI=confidence interval; HR=hazard ratio. Multivariable model: Cox regression using smoking status and intensity (never; former; current - <15 cigarettes per day; current - ≥15 cigarettes per day; current - intensity unknown; unknown), alcohol consumption frequency (never; special occasions only; 1–3 times/month; 1–2 times/week; 3–4 times per week; daily or almost daily; unknown), family history of colorectal cancer (no; yes; unknown), prevalent diabetes (no; yes; unknown), regular aspirin/ibuprofen use (no; yes; unknown), qualifications (none; CSEs/O-levels/GCSEs or equivalent; NVQ/HND/HNC/A-levels/AS-levels or equivalent; other professional qualifications; college/university degree; unknown), ever use of hormone replacement therapy (no, yes, unknown), frequency of red and processed meat consumption (<2 times per week; 2–3 times per week; 3–4 times/week; ≥4 times per week; unknown), and stratified by sex, age (5-year categories), Townsend deprivation index fifths, and region of the recruitment assessment centre.

25–<30, ≥30 kg m⁻²), waist circumference (sex-specific thirds), smoking status (never, former, current), and television watching time (≤1, 2–3, ≥4 h per day). Interaction terms (multiplicative scale) between these variables and physical activity were included in separate models. The statistical significance of the cross-product terms were evaluated using the likelihood ratio test. Similarly, the television watching time and colorectal cancer relationship was further assessed across subgroups of time spent using computers each day (none, anytime). To assess the possibility of reverse causality, cases diagnosed within the first two years of follow-up were excluded from all of the analyses. We repeated the analyses with multiple imputations to account for missing total physical activity information; however no differences were detected with the complete case analysis results presented (data not shown).

Statistical tests were all two-sided and a *P*-value of <0.05 was considered statistically significant. Analyses were conducted using Stata version 13.1.

RESULTS

After a median follow-up time of 5.6 years, 2391 cases of colorectal cancer (1389 in men and 1002 in women) were recorded. Of these cases, 1557 were colon tumours (784 proximal colon, 694 distal colon, and 79 overlapping or unspecified), 820 were rectal tumours, and 14 were overlapping between the colon and rectum. Compared with those in the lowest category (<10 MET hours per week), men and women in the highest total physical activity category (≥60 MET hours per week) had lower BMI and waist circumferences, were less likely to have attained a college education or university degree, to be never smokers, and to be regular aspirin/ibuprofen users (Table 1).

Physical activity

Colorectal cancer. There was no association between physical activity and colorectal cancer risk in the sexes-combined multivariable model (HR for ≥60 vs <10 MET hours per week = 0.92, 95% CI: 0.81–1.04; *p*-trend = 0.25), with similar relationships found for men and women (*p*-interaction = 0.57) (Table 2).

Colon cancer. For colon cancer, in the sexes-combined model, the highest physical activity (METs) category had a 16% lower risk compared to those in the lowest category (HR for ≥60 vs <10 MET hours per week = 0.84, 95% CI: 0.72–0.98; *p*-trend = 0.04). A statistically significant inverse association between physical activity and colon cancer risk was found only among men (HR for ≥60 vs <10 MET hours per week = 0.77, 95% CI: 0.62–0.95; *p*-trend = 0.02), but not women (HR for ≥60 vs <10 MET hours per week = 0.93, 95% CI: 0.74–1.17; *p*-trend = 0.58); however, this

difference between sexes did not reach statistical significance (*p*-interaction = 0.25).

For the sexes-combined models, no heterogeneity by colonic subsite was found (*p*-heterogeneity = 0.52), with a statistically significant inverse association observed between total physical activity and proximal colon cancer risk (HR for ≥60 vs <10 MET hours per week = 0.79, 95% CI: 0.63–0.99; *p*-trend = 0.04), and a non-significant inverse association for distal colon cancer risk (HR for ≥60 vs <10 MET hours per week = 0.88, 95% CI: 0.70–1.11; *p*-trend = 0.38). Among men, a statistically significant inverse association was found between total physical activity and proximal colon cancer risk (HR for ≥60 vs ≤10 MET hours per week = 0.67, 95% CI: 0.49–0.90; *p*-trend = 0.01), but no relationship was observed for distal colon cancer risk. Among women, no significant associations were observed between total physical activity and risks proximal colon and distal colon. The differences between the sexes did not reach statistical significance for proximal colon cancer (*p*-interaction = 0.15) and distal colon cancer (*p*-interaction = 0.89).

Additional statistical adjustment for waist circumference slightly attenuated the physical activity associations for colon cancer and its subsites (attenuated the HR for the ≥60 MET hours per week group by between 2 and 8%), and rendered the HRs in this physical activity group no longer statistically significant for all models, except for proximal colon cancer among men (HR for ≥60 vs <10 MET hours per week = 0.72, 95% CI: 0.53–0.98) (Table 2).

Rectal cancer. There was no relationship observed between physical activity and risk of rectal cancer for sexes-combined models, and for when men and women were analysed separately (*p*-interaction = 0.69).

Television watching time and time spent using computers and colorectal cancer risk

Colorectal cancer. In the sexes-combined multivariable model, higher levels of television watching time was associated with a greater colorectal cancer risk (HR for >5 h per day vs ≤1 h per day = 1.26, 95% CI: 1.04–1.53; *p*-trend = 0.01) (Table 3). A statistically significant positive association between television watching time and colorectal cancer was only found for men (HR for ≥4 h per day vs ≤1 h per day = 1.35, 95% CI: 1.13–1.61; *p*-trend = 0.02), but not women (HR for ≥4 h/day vs ≤1 h/day = 1.11, 95% CI: 0.91–1.35; *p*-trend = 0.25); however, this difference between sexes did not reach statistical significance (*p*-interaction = 0.21). Additional statistical adjustment for waist circumference slightly attenuated the association between television watching time colorectal cancer (5.5% reduction in HR of the ≥5 h per day group) and rendered the HRs in this television watching time group to be no longer statistically significant. Time

Table 3. Risk (hazard ratios) of colorectal cancer associated with television watching time

	Television watching time				
Colorectal cancer					
Both sexes	≤1 h per day	2–3 h per day	4–5 h per day	> 5 h per day	<i>p</i> -trend
<i>N</i>	373	1186	663	159	
Multivariable, HR (95% CI)	1	1.17 (1.04–1.32)	1.24 (1.08–1.41)	1.26 (1.04–1.53)	0.01
Multivariable plus waist circumference, HR (95% CI)	1	1.15 (1.02–1.29)	1.19 (1.04–1.36)	1.19 (0.98–1.45)	0.07
Men	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	196	685	501		
Multivariable, HR (95% CI)	1	1.24 (1.05–1.46)	1.35 (1.13–1.61)		0.02
Multivariable plus waist circumference, HR (95% CI)	1	1.21 (1.03–1.42)	1.28 (1.07–1.53)		0.09
Women	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	177	501	321		
Multivariable, HR (95% CI)	1	1.09 (0.92–1.30)	1.11 (0.91–1.35)		0.25
Multivariable plus waist circumference, HR (95% CI)	1	1.08 (0.90–1.29)	1.08 (0.88–1.32)		0.40
Colon cancer					
Both sexes	≤1 h per day	2–3 h per day	4–5 h per day	> 5 h per day	<i>p</i> -trend
<i>N</i>	246	753	441	111	
Multivariable, HR (95% CI)	1	1.14 (0.98–1.32)	1.25 (1.06–1.47)	1.32 (1.04–1.68)	0.007
Multivariable plus waist circumference, HR (95% CI)	1	1.11 (0.96–1.29)	1.19 (1.01–1.41)	1.23 (0.97–1.57)	0.045
Men	≤1 h/d	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	111	408	311		
Multivariable, HR (95% CI)	1	1.31 (1.05–1.62)	1.45 (1.15–1.83)		0.02
Multivariable plus waist circumference, HR (95% CI)	1	1.27 (1.02–1.57)	1.37 (1.08–1.72)		0.09
Women	≤1 h/d	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	135	345	241		
Multivariable, HR (95% CI)	1	1.00 (0.82–1.23)	1.09 (0.87–1.37)		0.16
Multivariable plus waist circumference, HR (95% CI)	1	0.98 (0.80–1.21)	1.05 (0.84–1.33)		0.28
Proximal colon cancer					
Both sexes	≤1 h per day	2–3 h per day	4–5 h per day	> 5 h per day	<i>p</i> -trend
<i>N</i>	127	356	239	58	
Multivariable, HR (95% CI)	1	1.04 (0.84–1.27)	1.28 (1.02–1.61)	1.29 (0.93–1.80)	0.009
Multivariable plus waist circumference, HR (95% CI)	1	1.00 (0.81–1.23)	1.20 (0.96–1.52)	1.18 (0.85–1.65)	0.047
Men	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	58	191	162		
Multivariable, HR (95% CI)	1	1.18 (0.88–1.60)	1.45 (1.05–1.99)		0.03
Multivariable plus waist circumference, HR (95% CI)	1	1.14 (0.84–1.54)	1.35 (0.98–1.86)		0.08
Women	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	69	165	135		
Multivariable, HR (95% CI)	1	0.91 (0.68–1.22)	1.14 (0.83–1.57)		0.16
Multivariable plus waist circumference, HR (95% CI)	1	0.88 (0.66–1.18)	1.08 (0.78–1.48)		0.30
Distal colon cancer					
Both sexes	≤1 h per day	2–3 h per day	4–5 h per day	> 5 h per day	<i>p</i> -trend
<i>N</i>	103	357	186	47	
Multivariable, HR (95% CI)	1	1.29 (1.03–1.62)	1.26 (0.97–1.62)	1.33 (0.92–1.92)	0.28
Multivariable plus waist circumference, HR (95% CI)	1	1.27 (1.02–1.59)	1.22 (0.94–1.58)	1.27 (0.88–1.83)	0.44
Men	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	48	196	137		
Multivariable, HR (95% CI)	1	1.43 (1.04–1.98)	1.45 (1.02–2.06)		0.36
Multivariable plus waist circumference, HR (95% CI)	1	1.40 (1.02–1.94)	1.39 (0.98–1.97)		0.54
Women	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	55	161	96		
Multivariable, HR (95% CI)	1	1.16 (0.85–1.59)	1.09 (0.76–1.56)		0.59
Multivariable plus waist circumference, HR (95% CI)	1	1.15 (0.84–1.58)	1.07 (0.74–1.53)		0.66
Rectal cancer					
Both sexes	≤1 h per day	2–3 h per day	4–5 h per day	> 5 h per day	<i>p</i> -trend
<i>N</i>	126	423	221	46	
Multivariable, HR (95% CI)	1	1.22 (0.99–1.50)	1.24 (0.99–1.56)	1.13 (0.79–1.61)	0.43
Multivariable plus waist circumference, HR (95% CI)	1	1.21 (0.99–1.48)	1.21 (0.96–1.53)	1.09 (0.76–1.55)	0.60
Men	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	84	269	192		
Multivariable, HR (95% CI)	1	1.13 (0.88–1.45)	1.24 (0.95–1.63)		0.29
Multivariable plus waist circumference, HR (95% CI)	1	1.11 (0.87–1.43)	1.20 (0.91–1.58)		0.43
Women	≤1 h per day	2–3 h per day	≥4 h per day		<i>p</i> -trend
<i>N</i>	42	154	75		
Multivariable, HR (95% CI)	1	1.42 (0.99–2.01)	1.15 (0.77–1.73)		0.87
Multivariable plus waist circumference, HR (95% CI)	1	1.41 (0.99–1.99)	1.14 (0.76–1.71)		0.82

Abbreviations: CI=confidence interval; HR=hazard ratio. Multivariable model: Cox regression using physical activity (MET hours per week; fifths), smoking status and intensity (never; former; current <15 cigarettes/day; current ≥15 cigarettes per day; current – intensity unknown; unknown), alcohol consumption frequency (never; special occasions only; 1–3 times/month; 1–2 times per week; 3–4 times per week; daily or almost daily; unknown), family history of colorectal cancer (no; yes; unknown), prevalent diabetes (no; yes; unknown), regular aspirin/ibuprofen use (no; yes; unknown), qualifications (none; CSEs/O-levels/GCSEs or equivalent; NVQ/HND/HNC/A-levels/AS-levels or equivalent; other professional qualifications; college/university degree; unknown), ever use of hormone replacement therapy (no, yes, unknown), frequency of red and processed meat consumption (<2 times per week; 2–3 times/week; 3–4 times/week; ≥4 times per week; unknown), and stratified by sex, age (5-year categories), Townsend deprivation index fifths, and region of the recruitment assessment centre.

Table 4. Risk (hazard ratios) of colorectal cancer associated with time spent on computers

		Time spent on computers				
		None	0.1–1 h per day	2–3 h/d	≥4 h per day	p-trend
Colorectal cancer						
Both sexes	N	1837	408	92	40	
	Multivariable, HR (95% CI)	1	0.95 (0.85–1.06)	1.07 (0.87–1.32)	0.94 (0.68–1.29)	0.75
	Multivariable plus waist circumference, HR (95% CI)	1	0.93 (0.83–1.04)	1.04 (0.84–1.29)	0.91 (0.67–1.25)	0.50
Men	N	995	292	96		p-trend
	Multivariable, HR (95% CI)	1	0.98 (0.86–1.12)	1.04 (0.84–1.29)		0.89
	Multivariable plus waist circumference, HR (95% CI)	1	0.96 (0.84–1.10)	1.01 (0.82–1.25)		0.82
Women	N	842	116	36		p-trend
	Multivariable, HR (95% CI)	1	0.90 (0.74–1.10)	1.03 (0.73–1.44)		0.58
	Multivariable plus waist circumference, HR (95% CI)	1	0.89 (0.73–1.09)	1.01 (0.72–1.41)		0.49
Colon cancer						
Both sexes	N	1199	263	61	23	p-trend
	Multivariable, HR (95% CI)	1	0.96 (0.84–1.10)	1.12 (0.86–1.45)	0.86 (0.56–1.30)	0.81
	Multivariable plus waist circumference, HR (95% CI)	1	0.94 (0.82–1.08)	1.08 (0.84–1.41)	0.83 (0.55–1.26)	0.55
Men	N	599	171	59		p-trend
	Multivariable, HR (95% CI)	1	0.94 (0.79–1.12)	1.05 (0.80–1.37)		0.83
	Multivariable plus waist circumference, HR (95% CI)	1	0.92 (0.77–1.09)	1.00 (0.76–1.32)		0.58
Women	N	600	92	25		p-trend
	Multivariable, HR (95% CI)	1	1.01 (0.81–1.27)	1.01 (0.68–1.52)		0.98
	Multivariable plus waist circumference, HR (95% CI)	1	1.00 (0.80–1.25)	0.99 (0.66–1.49)		0.86
Proximal colon cancer						
Both sexes	N	601	140	26	11	p-trend
	Multivariable, HR (95% CI)	1	1.04 (0.86–1.25)	0.97 (0.65–1.44)	0.84 (0.46–1.53)	0.79
	Multivariable plus waist circumference, HR (95% CI)	1	1.01 (0.84–1.22)	0.93 (0.63–1.38)	0.81 (0.44–1.47)	0.55
Men	N	292	92	27		p-trend
	Multivariable, HR (95% CI)	1	1.03 (0.81–1.31)	0.95 (0.64–1.42)		0.86
	Multivariable plus waist circumference, HR (95% CI)	1	1.00 (0.79–1.27)	0.91 (0.61–1.35)		0.63
Women	N	309	48	10		p-trend
	Multivariable, HR (95% CI)	1	1.06 (0.78–1.44)	0.84 (0.45–1.59)		0.77
	Multivariable plus waist circumference, HR (95% CI)	1	1.03 (0.75–1.40)	0.81 (0.43–1.54)		0.64
Distal colon cancer						
Both sexes	N	532	114	32	12	p-trend
	Multivariable, HR (95% CI)	1	0.93 (0.75–1.14)	1.31 (0.91–1.88)	1.00 (0.56–1.79)	0.62
	Multivariable plus waist circumference, HR (95% CI)	1	0.91 (0.74–1.12)	1.28 (0.89–1.84)	0.98 (0.55–1.75)	0.74
Men	N	278	72	30		p-trend
	Multivariable, HR (95% CI)	1	0.85 (0.66–1.11)	1.18 (0.81–1.74)		0.91
	Multivariable plus waist circumference, HR (95% CI)	1	0.84 (0.65–1.10)	1.15 (0.78–1.69)		0.96
Women	N	254	42	14		p-trend
	Multivariable, HR (95% CI)	1	1.09 (0.78–1.52)	1.31 (0.76–2.26)		0.36
	Multivariable plus waist circumference, HR (95% CI)	1	1.08 (0.78–1.51)	1.30 (0.75–2.24)		0.39
Rectal cancer						
Both sexes	N	624	145	31	17	p-trend
	Multivariable, HR (95% CI)	1	0.94 (0.78–1.13)	1.00 (0.70–1.45)	1.10 (0.68–1.79)	0.98
	Multivariable plus waist circumference, HR (95% CI)	1	0.93 (0.77–1.12)	0.99 (0.68–1.42)	1.08 (0.66–1.76)	0.86
Men	N	389	122	36		p-trend
	Multivariable, HR (95% CI)	1	1.06 (0.86–1.30)	1.03 (0.72–1.45)		0.67
	Multivariable plus waist circumference, HR (95% CI)	1	1.04 (0.85–1.29)	1.00 (0.71–1.42)		0.79
Women	N	235	23	12		p-trend
	Multivariable, HR (95% CI)	1	0.62 (0.40–0.96)	1.19 (0.66–2.13)		0.56
	Multivariable plus waist circumference, HR (95% CI)	1	0.62 (0.40–0.95)	1.17 (0.65–2.11)		0.54

Abbreviations: CI=confidence interval; HR=hazard ratio. Multivariable model: Cox regression using physical activity (MET hours per week; fifths), smoking status and intensity (never; former; current < 15 cigarettes per day; current ≥ 15 cigarettes per day; current – intensity unknown; unknown), alcohol consumption frequency (never; special occasions only; 1–3 times per month; 1–2 times per week; 3–4 times per week; daily or almost daily; unknown), family history of colorectal cancer (no; yes; unknown), prevalent diabetes (no; yes; unknown), regular aspirin/ibuprofen use (no; yes; unknown), qualifications (none; CSEs/O-levels/GCSEs or equivalent; NVQ/HND/HNC/A-levels/AS-levels or equivalent; other professional qualifications; college/university degree; unknown), ever use of hormone replacement therapy (no, yes, unknown), frequency of red and processed meat consumption (<2 times per week; 2–3 times/week; 3–4 times/week; ≥4 times/week; unknown), and stratified by sex, age (5-year categories), Townsend deprivation index fifths, and region of the recruitment assessment centre.

spent using computers was not associated with risk of colorectal cancer (Table 4), and this relationship did not differ after additional adjustment for television watching time (Supplementary Table 3).

Colon cancer. In the sexes-combined multivariable model, higher levels of television watching time was associated with a greater colon cancer risk (HR for >5 h per day vs ≤ 1 h per day = 1.32, 95% CI: 1.04–1.68; p -trend = 0.007) (Table 3). This positive relationship was only present among men, and not women, although this difference did not reach statistical significance (p -interaction = 0.10). The sexes-combined association between television watching time and colon cancer risk did not differ by subsite with HRs of similar strength observed for proximal and distal colon cancers (p -heterogeneity = 0.93) (Table 3). The positive relationships between television watching time and proximal and distal colon cancer were stronger and statistically significant for men than for women; however, these differences did not reach statistical significance (proximal colon p -interaction = 0.31; distal colon p -interaction = 0.28). Similar relationships between television watching time and colon cancer were observed after additional statistical adjustment for time spent on computers (Supplementary Table 2). Time spent using computers was not associated with risk of colon cancer, proximal colon cancer, and distal colon cancer (Table 4), and these relationships did not differ after additional adjustment for television watching time (Supplementary Table 3). Additional statistical adjustment for waist circumference slightly attenuated the association between television watching time and cancers of the colon and proximal colon (the HRs were attenuated for the ≥ 5 h per day group by between 6 and 8%), and rendered the HRs in this television watching time group to be no longer statistically significant.

Rectal cancer. There were no relationships observed between television watching time and time spent using computers and rectal cancer risk in the sexes-combined models, and for when men and women were analysed separately (television watching time p -interaction = 0.77; time spent using computers p -interaction = 0.70).

Sensitivity and subgroup analyses. The associations between total physical activity and colorectal cancer were similar across subgroups of BMI (p -interaction = 0.61), waist circumference (p -interaction = 0.93), smoking status (p -interaction = 0.14), and television watching time (p -interaction = 0.24) (data not shown). The relationship between television watching time and colorectal cancer did not differ across subgroups of time spent using computers (p -interaction = 0.23). Similar relationships were also observed when colorectal cancer cases which occurred during the first two years of follow-up were excluded from the analyses (data not shown).

DISCUSSION

In this prospective analysis of UK Biobank participants, total physical activity levels were inversely associated with colon cancer risk. This relationship was most apparent among men and for proximal colon cancer. Conversely, sedentary behaviour, as indicated by television watching time, was associated with elevated colon cancer risk. Total physical activity and the sedentary behaviours were not associated with rectal cancer risk in this population.

Our finding that total physical activity was associated with lower colon cancer risk is in accordance with a large body of epidemiological evidence (Friedenreich *et al*, 2006; Wolin *et al*, 2009; Moore *et al*, 2016). This relationship was consistent across body habitus and smoking status subgroups, and was most

apparent for men compared to women, although this difference was not statistically significant. Consistent with other individual prospective analyses (Lee *et al*, 2007; Burón Pust *et al*, 2017; Wei *et al*, 2017), we found no relationship between total physical activity and rectal cancer risk. However, a recent participant-level pooled analysis which included 5531 rectal cancer cases reported an inverse relationship between physical activity and rectal cancer incidence (highest vs lowest leisure-time physical activity group, relative risk (RR) = 0.87, 95% CI: 0.80–0.95) (Moore *et al*, 2016), which suggests that the null results reported by individual studies such as the current analysis ($n = 820$ rectal cancer cases) may be a consequence of insufficient sample size to detect weak-to-moderate associations.

For colon cancer, although we observed statistically significant inverse relationships for tumours in the proximal region and not the distal region with physical activity, this heterogeneity did not differ statistically. These results are generally consistent with those reported by other large prospective studies which did not observe heterogeneous physical activity relationships across the proximal and distal colon regions (Howard *et al*, 2008; Burón Pust *et al*, 2017; Wei *et al*, 2017), with the notable exception of the EPIC study (Friedenreich *et al*, 2006). Recently, in the Million's Women Study, which included 12 761 colon cancer cases, similar magnitude inverse relationships were observed between strenuous exercise and proximal and distal colon cancers (Burón Pust *et al*, 2017). Similarly, in the Nurses' Health Study and the NIH-AARP Diet and Health Study, lower risks were generally observed between various indicators of physical activity and tumours in both colonic regions (Howard *et al*, 2008; Wei *et al*, 2017).

The results of our study contribute to a growing body of evidence which has reported sedentary behaviour as a risk factor for the development of colon cancer, independent of physical activity levels (Howard *et al*, 2008; Schmid and Leitzmann, 2014). A recent meta-analysis reported that when comparing the highest with the lowest total sitting time group, a 24% greater risk (RR = 1.24, 95% CI: 1.03–1.50) of developing colon cancer was observed (Schmid and Leitzmann, 2014). Consistent with the findings from our analysis, for television watching time the same meta-analysis reported a 54% elevated risk (RR = 1.54, 95% CI: 1.19–1.98) of colon cancer when the highest and lowest viewing time groups were compared (Schmid and Leitzmann, 2014). Television viewing time has been shown to be accompanied with other lifestyle behaviours, such as smoking and eating energy dense foods (Wiecha *et al*, 2006; Schmid and Leitzmann, 2014), which are also related to cancer risk (Schmid and Leitzmann, 2014). We did not, however, observe any relationship between time spent on computers and colorectal cancer risk. It has been previously shown that for sedentary behaviour questionnaires, television watching assessment has higher test-retest reliability than computer time assessment (Lynch *et al*, 2014; Wijndaele *et al*, 2014). Such differences in the assessment of domain-specific sedentary behaviours may have impacted upon the contrasting associations we observed for television watching and time spent on computers with colorectal cancer risk.

The biological mechanisms through which physical activity potentially lowers colon cancer risk are uncertain. Being physically active is associated with less weight gain and body fatness (Lee *et al*, 2010; May *et al*, 2012), and lower adiposity is associated with a reduced risk of colorectal cancer (Larsson and Wolk, 2007; Moghaddam *et al*, 2007). Since body size/adiposity is possibly on the causal pathway linking physical activity and sedentary activities with colorectal cancer, we ran all multivariable models with and without additional adjustment for waist circumference. Adjusting for waist circumference resulted in a modest attenuation of the risk estimates (1–8%). These results support the hypothesis that the inverse physical activity and colorectal cancer relationship is at least partially mediated through changes in body size and

composition. As well as influencing weight gain and body fatness, greater physical activity and lower sedentary behaviours have also been associated with lower insulin levels and inflammation (Woods *et al*, 2006; Dunstan *et al*, 2007; Helmerhorst *et al*, 2009), both of which have been linked to the development of colorectal cancer (Kaaks *et al*, 2000; Terzic *et al*, 2010; Murphy *et al*, 2016). Physical activity may also reduce colorectal cancer risk by stimulating digestion and reducing transit time through the intestine (Slattery, 2004; Friedenreich *et al*, 2010).

The current analysis represents one of the largest single studies investigating the relationships between physical activity levels and sedentary behaviour with colorectal cancer risk to date. The large number of incident colorectal cancer cases allowed analyses by sex and across colorectal subsites, and the detailed phenotypic information collected from UK Biobank participants enabled us to carefully adjust for known colorectal cancer risk factors. A limitation of our analysis is that due to the relatively short follow-up time (median of 5.6 years) preclinical disease may be influencing the observed associations. However, we observed similar relationships when we excluded cases which occurred during the first 2 years of follow-up. In addition, a recent large participant level pooled analysis reported similar magnitude inverse relationships between leisure-time physical activity and colorectal cancer when stratified by follow-up period (<5 years and ≥ 5 years) (Moore *et al*, 2016), which suggests our results observed in a relatively short follow-up time may not be substantially influenced by preclinical disease. Another limitation of the study is that physical activity and sedentary behaviour information was self-reported by participants at the baseline questionnaire. Recently, the UK Biobank has objectively measured activity levels using 7-day accelerometers, in over 100 000 participants (Doherty *et al*, 2017). Currently, the follow-up time since these accelerometer measurements were collected is too short to study incident colorectal cancer, but these data will offer unique future insights into the physical activity and cancer relationship.

In conclusion, in this prospective analysis of UK Biobank participants, total physical activity levels were inversely associated with colon cancer risk, with no heterogeneity found between tumours located in the proximal colon and distal colon. Sedentary behaviour, as indicated by television watching time, was associated with an elevated colon cancer risk. These findings add to the large body of evidence which supports the promotion of physical activity in population-wide cancer prevention programmes.

ACKNOWLEDGEMENTS

This work has been conducted using the UK Biobank Resource under Application Number 25897 and we express our gratitude to the participants and those involved in building the resource. UK Biobank is an open access resource. Bona fide researchers can apply to use the UK Biobank data set by registering and applying at <http://www.ukbiobank.ac.uk/register-apply/>. KEB was supported by a Girdlers' New Zealand Health Research Council Fellowship.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DRJ, Tudor-Locke C, Greer JL, Vezina J, Whitt-Glover MC, Leon AS (2011) Compendium of physical activities: a second update of codes and MET values. *Med Sci Sports Exerc* **43**: 1575–1581.

- Allen N, Sudlow C, Downey P, Peakman T, Danesh J, Elliott P, Gallacher J, Green J, Matthews P, Pell J, Sprosen T, Collins R (2012) UK Biobank: current status and what it means for epidemiology. *Health Policy Technol* **1**: 123–126.
- Burón Pust A, Alison R, Blanks R, Pirie K, Gaitskell K, Barnes I, Gathani T, Reeves G, Beral V, Green J. Million women study, C (2017) Heterogeneity of colorectal cancer risk by tumour characteristics: large prospective study of UK women. *Int J Cancer* **140**: 1082–1090.
- Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, Pratt M, Ekelund U, Yngve A, Sallis JF, Oja P (2003) International physical activity questionnaire: 12-Country reliability and validity. *Med Sci Sports Exerc* **35**: 1381–1395.
- Doherty A, Jackson D, Hammerla N, Plötz T, Olivier P, Granat MH, White T, VAN Hees VT, Trenell MI, Owen CG, Preece SJ, Gillions R, Sheard S, Peakman T, Brage S, Wareham NJ (2017) Large scale population assessment of physical activity using wrist worn accelerometers: the UK biobank study. *PLOS ONE* **12**: e0169649.
- Dunstan DW, Salmon J, Healy GN, Shaw JE, Jolley D, Zimmet PZ, Owen N (2007) Association of television viewing with fasting and 2-h postchallenge plasma glucose levels in adults without diagnosed diabetes. *Diabetes Care* **30**: 516–522.
- Friedenreich C, Norat T, Steindorf K, Boutron-Ruault MC, Pischon T, Mazuir M, Clavel-Chapelon FO, Linseisen J, Boeing H, Bergman M, Johnsen NF, Tjønneland A, Overvad K, Mendez M, Quiros JR, Martinez C, Dorronsoro M, Navarro C, Gurrea AB, Bingham S, Khaw KT, Allen N, Key T, Trichopoulos A, Trichopoulos D, Orfanou N, Krogh V, Palli D, Tumino R, Panico S, Vineis P, Bueno-De-Mesquita HB, Peeters PHM, Monninkhof E, Berglund G, Manjer J, Ferrari P, Slimani N, Kaaks R, Riboli E (2006) Physical activity and risk of colon and rectal cancers: the European prospective investigation into cancer and nutrition. *Cancer Epidemiol Biomark Prev* **15**: 2398–2407.
- Friedenreich CM, Neilson HK, Lynch BM (2010) State of the epidemiological evidence on physical activity and cancer prevention. *Eur J Cancer* **46**: 2593–2604.
- Helmerhorst HJF, Wijndaele K, Brage S, Wareham NJ, Ekelund U (2009) Objectively measured sedentary time may predict insulin resistance independent of moderate- and vigorous-intensity physical activity. *Diabetes* **58**: 1776–1779.
- Howard RA, Freedman DM, Park Y, Hollenbeck A, Schatzkin A, Leitzmann MF (2008) Physical activity, sedentary behavior, and the risk of colon and rectal cancer in the NIH-AARP Diet and Health Study. *Cancer Causes Control* **19**: 939–953.
- Howley ET (2001) Type of activity: resistance, aerobic and leisure versus occupational physical activity. *Med Sci Sports Exerc* **33**: S364–S369.
- IARC-WHO (2015) GLOBOCAN 2012: Estimated Cancer Incidence, Mortality and Prevalence Worldwide in 2012.
- IPAQ research committee (2005) Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ)—Short and Long Forms.
- Kaaks R, Toniolo P, Akhmedkhanov A, Lukanova A, Biessy C, Dechaud H, Rinaldi S, Zeleniuch-Jacquotte A, Shore RE, Riboli E (2000) Serum C-peptide, insulin-like growth factor (IGF)-I, IGF-binding proteins, and colorectal cancer risk in women. *J Natl Cancer Inst* **92**: 1592–1600.
- Larsson SC, Wolk A (2007) Obesity and colon and rectal cancer risk: a meta-analysis of prospective studies. *Am J Clin Nutr* **86**: 556–565.
- Lee I, Djoussé L, Sesso HD, Wang L, Buring JE (2010) Physical activity and weight gain prevention. *J Am Med Assoc* **303**: 1173–1179.
- Lee K-J, Inoue M, Otani T, Iwasaki M, Sasazuki S, Tsugane S (2007) Physical activity and risk of colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective Study. *Cancer Causes Control* **18**: 199–209.
- Limsui D, Vierkant RA, Tillmans LS, Wang AH, Weisenberger DJ, Laird PW, Lynch CF, Anderson KE, French AJ, Haile RW, Harnack LJ, Potter JD, Slager SL, Smyrk TC, Thibodeau SN, Cerhan JR, Limburg PJ (2010) Cigarette smoking and colorectal cancer risk by molecularly defined subtypes. *J Natl Cancer Inst* **102**: 1012–1022.
- Lynch BM, Friedenreich CM, Khandwala F, Liu A, Nicholas J, Csizmadia I (2014) Development and testing of a past year measure of sedentary behavior: the SIT-Q. *BMC Public Health* **14**: 899.
- May AM, Romaguera D, Travier N, Ekelund U, Bergmann MM, Kaaks R, Teucher B, Steffen A, Boeing H, Halkjaer J, Tjønneland A, Jakobsen MU, Overvad K, Dartois L, Fagherazzi G, Boutron-Ruault M-C, Quiros JR, Agudo

- A, Gonzalez C, Sánchez M-J, Amiano P, Huerta J-M, Ardanaz E, Wareham NJ, Crowe FL, Naska A, Orfanos P, Trichopoulou A, Palli D, Agnoli C, Tumino R, Vineis P, Panico S, Bueno-De-Mesquita HB, Verschuren M, Drake I, Sonestedt E, Braaten T, Rinaldi S, Romieu I, Slimani N, Norat T, Riboli E, Peeters PHM (2012) Combined impact of lifestyle factors on prospective change in body weight and waist circumference in participants of the EPIC-PANACEA study. *PLoS ONE* 7: e50712.
- Moghaddam AA, Woodward M, Huxley R (2007) Obesity and risk of colorectal cancer: a meta-analysis of 31 studies with 70,000 events. *Cancer Epidemiol Biomark Prev* 16: 2533–2547.
- Moore SC, Lee IM, Weiderpass E, Campbell PT, Sampson JN, Kitahara CM, Keadle SK, Arem H, Berrington De Gonzalez A, Hartge P, Adami HO, Blair CK, Borch KB, Boyd E, Check DP, Fournier A, Freedman ND, Gunter M, Johansson M, Khaw KT, Linet MS, Orsini N, Park Y, Riboli E, Robien K, Schairer C, Sesso H, Spriggs M, VAN Dusen R, Wolk A, Matthews CE, Patel AV (2016) Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Int Med* 176: 816–825.
- Murphy N, Cross AJ, Abubakar M, Jenab M, Aleksandrova K, Boutron-Ruault MC, Dossus L, Racine A, Kuhn T, Katzke VA, Tjonneland A, Petersen KEN, Overvad K, Quiros JR, Jakszyn P, Molina-Montes E, Dorronsoro M, Huerta JM, Barricarte A, Khaw KT, Wareham N, Travis RC, Trichopoulou A, Lagiou P, Trichopoulos D, Masala G, Krogh V, Tumino R, Vineis P, Panico S, Bueno-De-Mesquita HB, Siersema PD, Peeters PH, Ohlsson B, Ericson U, Palmqvist R, Nystrom H, Weiderpass E, Skeie G, Freisling H, Kong SY, Tsilidis K, Muller DC, Riboli E, Gunter MJ (2016) A nested case–control study of metabolically defined body size phenotypes and risk of colorectal cancer in the European prospective investigation into cancer and nutrition (EPIC). *PLoS Med* 13: e1001988.
- Office for National Statistics (2006) *The Time Use Survey, 2005. How We Spend Our Time* London.
- Owen N, Leslie E, Salmon J, Fotheringham MJ (2000) Environmental determinants of physical activity and sedentary behavior. *Exerc Sport Sci Rev* 28: 153–158.
- Schmid D, Leitzmann MF (2014) Television viewing and time spent sedentary in relation to cancer risk: a meta-analysis. *J Natl Cancer Inst* 106: dju098–dju098.
- Schoenfeld D (1982) Partial residuals for the proportional hazards regression model. *Biometrika* 69: 239–241.
- Slattery ML (2004) Physical activity and colorectal cancer. *Sports Med* 34: 239–252.
- Terzic J, Grivnennikov S, Karin E, Karin M (2010) Inflammation and colon cancer. *Gastroenterology* 138: 2101–2114.e5.
- UK-Biobank (2010) UK Biobank. Protocol for a large-scale prospective epidemiological resources [Online] <http://www.ukbiobank.ac.uk/wp-content/uploads/2011/11/UK-Biobank-Protocol.pdf> Accessed 10 July 2016.
- WCRF-AICR (2011) World Cancer Research Fund–American Institute for Cancer Research. Continuous Update Project Report Summary. Food, Nutrition, Physical Activity, and the Prevention of Colorectal Cancer.
- Wei EK, Colditz GA, Giovannucci EL, WU K, Glynn RJ, Fuchs CS, Stampfer M, Willett W, Ogino S, Rosner B (2017) A comprehensive model of colorectal cancer by risk factor status and subsite using data From the nurses' health study. *Am J Epidemiol* 185: 224–237.
- Wei EK, Giovannucci E, WU K, Rosner B, Fuchs CS, Willett WC, Colditz GA (2004) Comparison of risk factors for colon and rectal cancer. *Int J Cancer* 108: 433–442.
- Wiecha JL, Peterson KE, Ludwig DS, Kim J, Sobol A, Gortmaker SL (2006) When children eat what they watch: impact of television viewing on dietary intake in youth. *Arch Pediatr Adolesc Med* 160: 436–442.
- Wijndaele K, DE Bourdeaudhuij I, Godino JG, Lynch BM, Griffin SJ, Westgate K, Brage S (2014) Reliability and validity of a domain-specific last 7-d sedentary time questionnaire. *Med Science in Sports Exerc* 46: 1248–1260.
- Wolin KY, Yan Y, Colditz GA, Lee IM (2009) Physical activity and colon cancer prevention: a meta-analysis. *Br J Cancer* 100: 611–616.
- Woods JA, Vieira VJ, Keylock KT (2006) Exercise, inflammation, and innate immunity. *Neurol Clin* 24: 585–599.

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 4.0 Unported License.

Supplementary Information accompanies this paper on British Journal of Cancer website (<http://www.nature.com/bjc>)