CORRIGENDUM

Achieving stringent CR is essential before reduced-intensity conditioning allogeneic hematopoietic cell transplantation in AML

C Ustun^{1,2}, AC Wiseman³, TE DeFor², S Yohe⁴, MA Linden⁴, B Oran¹, M Burke⁵, E Warlick^{1,2}, JS Miller^{1,2} and D Weisdorf^{1,2}¹Division of Hematology, Oncology and Transplantation, Department of Medicine, University of Minnesota, Minneapolis, MN, USA;²Blood and Marrow Transplant Program, University of Minnesota, Minneapolis, MN, USA; ³Medical School, University of Minnesota, Minneapolis, MN, USA; ⁴Division of Hematopathology, Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN, USA and ⁵Division of Hematology-Oncology, Department of Pediatrics, University of Minnesota, Minneapolis, MN, USA*Bone Marrow Transplantation* (2013) 48, 1488; doi:10.1038/bmt.2013.161**Correction to:** *Bone Marrow Transplantation* (2013) 48, 1415–1420; doi: 10.1038/bmt.2013.124; published online 12 August 2013

Since the publication of this article, the authors have noticed that the affiliations were incorrect. The correct affiliations are shown

above, and the corrected article appears in this issue. The html and online pdf versions of this paper have also been rectified, and now carry the correct paper.

The authors apologize for any inconvenience this has caused.