

Histone demethylase Kdm4b functions as a co-factor of C/EBP β to promote mitotic clonal expansion during differentiation of 3T3-L1 preadipocytes

L Guo¹, X Li^{1,2}, J-X Huang¹, H-Y Huang^{1,2}, Y-Y Zhang², S-W Qian², H Zhu², Y-D Zhang¹, Y Liu¹, Y Liu¹, K-K Wang³ and Q-Q Tang^{*1,2}

CCAAT/enhancer-binding protein (C/EBP) β is required for both mitotic clonal expansion (MCE) and terminal adipocyte differentiation of 3T3-L1 preadipocytes. Although the role of C/EBP β in terminal adipocyte differentiation is well defined, its mechanism of action during MCE is not. In this report, histone demethylase Kdm4b, as well as cell cycle genes *Cdc45l* (cell division cycle 45 homolog), *Mcm3* (mini-chromosome maintenance complex component 3), *Gins1* (GINS complex subunit 1) and *Cdc25c* (cell division cycle 25 homolog c), were identified as potential C/EBP β target genes during MCE by utilizing promoter-wide chromatin immunoprecipitation (ChIP)-on-chip analysis combined with gene expression microarrays. The expression of Kdm4b is induced during MCE and its induction is dependent on C/EBP β . ChIP, Electrophoretic Mobility Shift Assay (EMSA) and luciferase assay confirmed that the promoter of Kdm4b is bound and activated by C/EBP β . Knockdown of Kdm4b impaired MCE. Furthermore, Kdm4b interacted with C/EBP β and was recruited to the promoters of C/EBP β -regulated cell cycle genes, including *Cdc45l*, *Mcm3*, *Gins1*, and *Cdc25c*, demethylated H3K9me3 and activated their transcription. These findings suggest a novel feed forward mechanism involving a DNA binding transcription factor (C/EBP β) and a chromatin regulator (Kdm4b) in the regulation of MCE by controlling cell cycle gene expression.

Cell Death and Differentiation (2012) 19, 1917–1927; doi:10.1038/cdd.2012.75; published online 22 June 2012

Obesity is a major risk factor for type 2 diabetes, hypertension, hyperlipidemia, and arteriosclerosis.¹ Both the increase of adipocyte size (hypertrophy) and number (hyperplasia)² are major contributors to the development of obesity. The hyperplasia of adipocytes is mimicked during the adipogenic differentiation program of 3T3-L1 preadipocytes.³ Immediately after the treatment with differentiation inducers, growth-arrested 3T3-L1 preadipocytes re-enter the cell cycle, a process referred to as mitotic clonal expansion (MCE) in which cell number increases ~4-fold during differentiation. The gene expression program leading to terminal adipocyte differentiation is initiated during and after the MCE period. Although there is some controversy,⁴ numerous experimental observations support the view that MCE is a necessary step for the terminal adipocyte differentiation of 3T3-L1 preadipocytes. The extracellular signal-regulated kinase (MEK) inhibitor U0126 and cyclin-dependent kinase inhibitor roscovitine, which inhibit the cell cycle at different points, block MCE, as well as adipogenesis.⁵ Moreover, the DNA synthesis inhibitor aphidicolin and the anti-proliferation reagent rapamycin also block MCE and 3T3-L1 preadipocyte differentiation.^{6,7}

CCAAT/enhancer-binding protein β (C/EBP β) is an important transcription factor for the initiation of 3T3-L1 preadipocyte differentiation. C/EBP β is induced very early in adipocyte differentiation. Then it activates the expression of CCAAT enhancer-binding protein α (C/EBP α) and peroxisome proliferator-activated receptor γ (PPAR γ), two critical pro-adipogenic transcription factors, by binding to their promoters.⁸ Previous studies have shown that the expression and the activation of C/EBP β are required for MCE. When subjected to the same differentiation protocol as 3T3-L1 preadipocytes, a subset of mouse embryo fibroblasts (MEFs) undergo MCE and terminal differentiation into adipocytes. MEFs from C/EBP β (–/–) mice, however, neither undergo MCE nor differentiate into adipocytes.⁹ Furthermore, knockdown of C/EBP β by RNA interference (RNAi) in 3T3-L1 preadipocytes prevents MCE, as well as adipocyte differentiation.¹⁰ Additionally, overexpression of a dominant-negative C/EBP β (A-C/EBP) that blocks C/EBP β DNA binding by dimerizing through its leucine zipper, also disrupts MCE and adipogenesis of 3T3-L1 cells.¹¹ C/EBP β also has important roles in the proliferation of certain other cell types such as lobuloalveolar

¹Key Laboratory of Molecular Medicine, the Ministry of Education, Department of Biochemistry and Molecular Biology, Fudan University Shanghai Medical College, Shanghai 200032, People's Republic of China; ²Department of Stem Cell Research and Regenerative Medicine, Institutes of Biomedical Sciences, Fudan University, Shanghai 200032, People's Republic of China and ³Department of Medical Genomics, Sino-French Research Center for Life Sciences and Genomics, Rui-Jin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, People's Republic of China

*Corresponding author: Q-Q Tang, Key Laboratory of Molecular Medicine, the Ministry of Education, Department of Biochemistry and Molecular Biology, Fudan University Shanghai Medical College, Shanghai 200032, People's Republic of China. Tel: +86 21 54237198; Fax: +86 21 54237290; E-mail: qqtang@shmu.edu.cn

Keywords: 3T3-L1 preadipocytes; adipocyte differentiation; mitotic clonal expansion; C/EBP β ; Kdm4b; ChIP-on-chip

Abbreviations: Cdc25c, cell division cycle 25 homolog c; Cdc45l, cell division cycle 45 homolog; C/EBP β , CCAAT/enhancer-binding protein β ; C/EBP α , CCAAT enhancer-binding protein α ; ChIP, chromatin immunoprecipitation; EMSA, Electrophoretic Mobility Shift Assay; ER, estrogen receptor; Gins1, GINS complex subunit 1; LAP, liver-enriched activator protein; LIP, liver-enriched inhibitory protein; MCE, mitotic clonal expansion; Mcm3, mini-chromosome maintenance complex component 3; MECs, mammary epithelial cells; PPAR γ , peroxisome proliferator-activated receptor γ ; RNAi, RNA interference; TSS, transcription start site

Received 09.1.12; revised 20.4.12; accepted 08.5.12; Edited by RA Knight; published online 22.6.12

cells, osteoblasts, keratinocytes and augments Ha-*ras*-induced transformation of NIH 3T3 cells.^{12–15} Despite the important role of C/EBP β in growth stimulation, the mechanism underlying MCE is not fully understood.

The methylation of lysine residues in histones is a central epigenetic modification in the regulation of eukaryotic gene expression. While methylation at H3K4 and H3K36 primarily transduces activating signals, methylation at H3K9, H3K27, and H4K20 is associated primarily with repressed chromatin. Kdm4b is a JmjC-domain containing histone demethylase for H3K9me3. Recent studies have shown that Kdm4b is required for estrogen receptor α (ER α)-regulated breast cancer progression and mammary epithelial cells (MECs) proliferation.^{16–18} In the present study, Kdm4b was identified as a target gene of C/EBP β that functions as a co-factor of C/EBP β to demethylate H3K9me3 in the regulatory regions of C/EBP β -regulated cell cycle genes *Cdc45l* (cell division cycle 45 homolog), *Mcm3* (mini-chromosome maintenance complex component 3), *Gins1* (GINS complex subunit 1) and *Cdc25c* (cell division cycle 25 homolog c), thereby promoting their expression and MCE. This explains why C/EBP β is required for MCE during adipocyte differentiation.

Results

Identification of C/EBP β binding sites during the MCE of 3T3-L1 preadipocyte differentiation. A promoter-wide chromatin immunoprecipitation (ChIP)-on-chip analysis was employed on 3T3-L1 cells harvested at 20 h after hormonal induction when the cells are in the middle S phase during the first round of MCE.^{5,19} Nine-hundred and forty-one significant

C/EBP β binding regions were identified (Supplementary Table S1). To validate the ChIP-on-chip data, C/EBP β enrichment was measured by ChIP quantitative PCR (ChIP-qPCR) at 30 novel binding regions and 29 of these sites were positive (Supplementary Figure S1a). Fifteen of these sites were also tested by ChIP-qPCR with another C/EBP β antibody and a similar result was obtained as with the original antibody used for the ChIP-on-chip assays (Supplementary Figure S1b).

Distribution analysis of the binding sites relative to the TSS showed that the majority of C/EBP β binding regions were located at distances within 2 kb from the nearest annotated transcription start sites (TSSs) (Figure 1a). Further analysis revealed that ~30% of C/EBP β binding sites were at proximal promoters (Figure 1b, TSS to -1 kb), and ~30% of the sites fell within distal promoters, defined as >1 kb upstream from TSS (Figure 1b, >1 kb), while many other sites (26.84%) were located in introns (Figure 1b). Evolutionary conservation of the C/EBP β binding regions was examined to show that there is a high degree of conservation of C/EBP β binding sites among higher eukaryotes (Figure 1c), suggesting that these binding sites are likely to be functional transcriptional regulatory regions across species.

The sequences of C/EBP β -bound regions were mined to further identify potential binding motifs for transcription factors. A *de novo* motif search identified a sequence that strongly resembles the C/EBP recognition element as the top-scoring motif (Figure 1d). Consistently, the highest scoring motifs from the TRANSFAC database represented different matrices for C/EBP proteins (Figure 1e and Supplementary Table S2). Gene expression microarrays were further




Figure 1 Bioinformatic analyses of the ChIP-on-chip data. Post-confluent growth-arrested 3T3-L1 preadipocytes were induced to differentiation as described. At 20 h ChIP-on-chip were performed and analyzed. (a) Distribution in 200-bp intervals of C/EBP β binding regions relative to TSSs of the nearest RefSeq genes. (b) Mapping of C/EBP β binding sites relative to the nearest RefSeq genes. >1 kb: more than 1 kb upstream from the TSS; Downstream: downstream from the 3' end of the gene; UTR: untranslated region. (c) Average plot for conservation of the C/EBP β binding regions among higher eukaryotes. (d) Sequence logo illustrating the top-scoring *de novo* motif for C/EBP β -bound regions. (e) TRANSFAC matrices enriched around C/EBP β -bound regions. (f) Overlap between the genes targeted by C/EBP β (1039 genes) and the genes whose induction was dependent on C/EBP β (481 genes)

conducted for both control RNAi-treated cells and C/EBP β RNAi-treated cells to confirm the functional relationship between C/EBP β binding and gene transcription. Four hundred and eighty-one genes were identified to be induced (20h *versus* 0h) in a C/EBP β -dependent manner (Supplementary Tables S3 and S4). When overlapping ChIP-on-chip data and microarray data, 68 potential C/EBP β target genes were identified (Figure 1f and Supplementary Table S5).

Histone demethylase Kdm4b was identified as one of the C/EBP β target genes required for MCE during 3T3-L1 adipocyte differentiation. Lysine methylation on histones has a pivotal role in transcriptional regulation.²⁰ The histone demethylase Kdm4b (for H3K9me3) was among the 68 potential C/EBP β target genes identified above. This suggests that C/EBP β might regulate gene expression through indirect regulation of histone lysine methylation.

A DNA fragment containing the 5'-flanking region of *Kdm4b* gene was subcloned into the luciferase reporter construct pGL3-basic and transfected into 3T3-L1 preadipocytes with/without the C/EBP β expression plasmid. Luciferase activity was increased ~10-fold when co-transfected with C/EBP β expression plasmid (Figure 2a). Consistently, deletion of the C/EBP binding site (-577 to -564 bp) identified in our ChIP-on-chip analysis almost totally abolished this transactivation. ChIP-qPCR confirmed that the Kdm4b promoter fragment containing the region of -577 bp ~ -564 bp was significantly enriched by the C/EBP β antibody at 16h and 20h after adipogenic induction (Figure 2b). The specific binding of C/EBP β to this region was also confirmed by Electrophoretic Mobility Shift Assay (EMSA) (Supplementary Figure S2). Upon induction of differentiation, Kdm4b messenger RNA (mRNA) level was significantly induced and this upregulation was prevented when C/EBP β was knocked down by RNAi (Figure 2c). Consistently, Kdm4b protein level also increased post-induction and was impaired in C/EBP β RNAi-treated cells (Figure 2d). The knockdown efficiency of C/EBP β RNAi was also confirmed (Figures 2c and d) and its effect on MCE and terminal adipocyte differentiation is consistent with our previous report.¹⁰ While the expression of Kdm4b is significantly induced and regulated by C/EBP β , this is not the case for other Kdm4 family genes (Figure 2e).

To further confirm the role of C/EBP β in Kdm4b induction, C/EBP β was overexpressed in 3T3-L1 cells (Figure 2f). As expected, ectopic expression of C/EBP β promoted MCE as illustrated by increased EdU incorporation (an indicator of DNA synthesis, like BrdU incorporation) (Figure 2g), as well as adipocyte differentiation, as indicated by increased accumulation of cytoplasmic triglyceride (Figure 2h) and enhanced the expression of Kdm4b at both 0h and 20h after adipogenic induction (Figure 2i). These results demonstrate that Kdm4b is a *bona fide* target of C/EBP β during MCE.

Kdm4b has been shown to regulate cell proliferation, including estrogen-induced breast cancer growth and MECs proliferation.^{16,17} Therefore, we hypothesized that Kdm4b might also be involved in regulating MCE during 3T3-L1 preadipocyte differentiation. RNAi oligonucleotide of Kdm4b was transfected into 3T3-L1 preadipocytes. The efficiency




Figure 2 Transactivation of Kdm4b by C/EBP β during MCE of 3T3-L1 preadipocyte differentiation. (a) 3T3-L1 preadipocytes were transfected with control vector or C/EBP β -expressing vector, together with reporter constructs containing the Kdm4b promoter (-1000 to -1 bp) or mutants. Luciferase activities were measured 36 h later and plotted. (b) 3T3-L1 cells were induced to differentiation and binding of C/EBP β on the Kdm4b promoter was analyzed by ChIP-qPCR at times indicated. (c) 3T3-L1 cells were treated with control or C/EBP β RNAi. After induction, the mRNA levels of C/EBP β and Kdm4b at times indicated were analyzed by RT-qPCR. (d) Cell lysates were harvested at times indicated and subjected to western blotting with antibodies indicated. The C/EBP β antibody detects two bands: liver-enriched activator protein (LAP, 34 kDa) and liver-enriched inhibitory protein (LIP, 21 kDa, an N-terminally truncated isoform of C/EBP β). Hsp90 served as a loading control. (e) After induction, mRNA levels of the Kdm4 family members at times indicated were analyzed by RT-qPCR. Data normalized to 0h time point of each gene. * $P < 0.05$. (f-i) Overexpression of C/EBP β promotes Kdm4b expression. 3T3-L1 preadipocytes were infected with retrovirus expressing flag-tagged C/EBP β or with empty vector and induced to differentiation. (f) The overexpression of C/EBP β (LAP) was confirmed by western blotting at 20h. (g) At 16h after induction, cells were labeled with EdU for 2h and then subjected to FACS analysis. The percentages of cells in S phase were determined. (h) At 8d after induction, cells were stained with Oil red O. (i) The effect of C/EBP β overexpression on the expression of Kdm4b was detected by RT-qPCR

and specificity of Kdm4b knockdown were confirmed by western blotting (Figure 3a) and RT-qPCR (Supplementary Figure S3) respectively. The RNAi effect on DNA synthesis was evaluated by using EdU labeling. As shown in Figure 3b, Kdm4b knockdown caused less EdU incorporation. Furthermore, flow cytometry analysis revealed that knockdown of Kdm4b resulted in a significant reduction in the cell population in S phase (from 34 to 19%) with a concomitant increase in the




Figure 4 Four cell cycle genes *Cdc45l*, *Mcm3*, *Gins1* and *Cdc25c* were identified as C/EBP β target genes. (a, b) 3T3-L1 cells were induced to differentiation. The expression profiles of C/EBP β and the four cell cycle genes *Cdc45l*, *Mcm3*, *Gins1* and *Cdc25c* were analyzed by RT-qPCR (a) and western blotting (b) respectively. (c, d) 3T3-L1 preadipocytes were transfected with control RNAi or C/EBP β RNAi. After post-confluence, cells were induced to differentiation. The effect of knocking down C/EBP β on the expression of the four cell cycle genes was detected by RT-qPCR (c) and western blotting (d) respectively

of these cell cycle genes was significantly reduced as indicated by the reduction of their mRNA and protein levels (Figures 4c and d). These results confirm that C/EBP β is required for the induction of the four cell cycle genes during MCE in the differentiation program of 3T3-L1 preadipocytes.

The binding of C/EBP β to the promoters of the four cell cycle genes was then investigated. The promoter regions of the four genes identified in the C/EBP β ChIP-on-chip data and the C/EBP β consensus sequences are illustrated in Supplementary Figure S4. ChIP-qPCR assays were performed using primers covering these C/EBP β consensus sequences of the four genes.

As Figure 5a shows, C/EBP β associates with the promoters of the four genes at 16 h and 20 h after induction. In contrast, C/EBP β occupancy was hardly detectable in quiescent cells before induction (i.e., at 0 h). EMSA was also performed using probes containing the C/EBP β consensus sequences and nuclear extracts from 3T3-L1 cells at 20 h after induction. As shown in Figure 5b, DNA–protein complexes were formed with these probes (lanes 1, 6, 11, 16). The C/EBP β containing DNA–protein complexes were verified by the completely supershift of the complexes using C/EBP β antibody (lanes 3, 8, 13, and 18) compared with that with control IgG (lanes 2, 7, 12, and 17). Moreover, the inhibition of complex formation by unlabeled competitor oligonucleotide, but not by self probes mutated in the C/EBP consensus sequence (Supplementary Figure S4), verified specific binding (compare lanes 4, 9, 14, 19 with lanes 5, 10, 15, 20). Furthermore, promoter-luciferase reporter analysis indicates that C/EBP β transactivates the promoters of the four cell cycle genes and that mutation of the C/EBP β binding sites significantly reduces expression (Figure 5c).

The role of the four cell cycle genes in MCE was further examined. Knockdown of the four genes inhibits MCE (Supplementary Figure S5). Additionally, ectopic expression of the four genes together significantly rescues the inhibitory effect of C/EBP β RNAi on MCE (Supplementary Figure S6). Taken together, the RNAi and overexpression experiments demonstrate that *Cdc45l*, *Mcm3*, *Gins1* and *Cdc25c*, which are transactivated by C/EBP β , have an important role in MCE.

Histone demethylase Kdm4b serves as a co-factor of C/EBP β during MCE.

Recent studies have revealed that Kdm4b is a direct target gene of ER α , forms complex with ER α and facilitates ER α -regulated transcription.^{16,17} As indicated above, *Kdm4b* has been shown to be a target gene of C/EBP β (Figure 2) and like C/EBP β , Kdm4b is also required for MCE (Figure 3). Thus, we hypothesized that Kdm4 might act as a co-factor of C/EBP β to promote MCE.

ChIP assays were performed to determine whether Kdm4b interacts with the C/EBP β -targeted cell cycle gene loci. The results revealed that, similar to C/EBP β , Kdm4b is recruited to the promoter regions of the four cell cycle genes 20 h after, but not before, induction (Figure 6a). Furthermore, re-ChIP analysis indicated that the promoters of the four genes, which are immunoprecipitated by C/EBP β antibody, are re-immunoprecipitated by Kdm4b antibody (Figure 6b). The interaction between C/EBP β and Kdm4b was further confirmed in co-immunoprecipitation experiments. Immunoprecipitation of the nuclear extracts at 20 h post-induction with C/EBP β antibody pulls down Kdm4b (Figure 6c), and Kdm4b antibody also immunoprecipitates C/EBP β (Figure 6d). Collectively, these results verify that C/EBP β and Kdm4b form a complex that targets the promoters of the four cell cycle genes.




Figure 5 C/EBP β binds to and transactivates the promoters of the four cell cycle genes. (a) Growth-arrested 3T3-L1 preadipocytes were induced to differentiation. At times indicated ChIP was performed to show the binding of C/EBP β to the promoters of the four cell cycle gene. Data normalized to the IgG controls at each time point. One region of the insulin gene serves as a negative control. (b) EMSA was conducted with nuclear extracts from 3T3-L1 cells at 20 h after induction. Supershift experiments were performed with control IgG or C/EBP β antibody. For competition analysis, unlabeled competitor probe of the C/EBP element in the C/EBP α gene promoter (Consensus) or unlabeled self probes mutated in the C/EBP element (Mut, shown in Supplementary Figure S4) were used. The positions of the C/EBP β containing DNA–protein complex and the supershift band (SS) are indicated. (c) 3T3-L1 preadipocytes were transfected with reporter constructs with/without the C/EBP binding sites mutated and with/without the C/EBP β expression vector. After 48 h, luciferase activity was analyzed and plotted. * $P < 0.05$

Furthermore, luciferase assays demonstrate that Kdm4b enhances the ability of C/EBP β to activate the promoters of the four cell cycle genes (Figure 6e), which indicates that Kdm4b functions as a positive transcriptional co-factor for C/EBP β to regulate the four cell cycle genes.

Kdm4b is a histone demethylase that reverses tri-methylation at H3K9. H3K9me3 is generally associated with gene silencing and is enriched in heterochromatin of inactive genes. It was found that H3K9me3 modification of the promoter regions of Cdc45l, Mcm3, Gins1 and Cdc25c is high at 0 h and significantly decreases by 20 h after induction (Figures 7a–d). Consistently, the decrease in H3K9me3 associated with these promoters following adipogenic induction was only observed in control, but not in Kdm4b-depleted cells (Figure 7e). As expected, knockdown of the expression of Kdm4b impaired the induction of the four cell cycle genes (Figure 7f).

C/EBP β and Kdm4b specifically regulate MCE, but not normal 3T3-L1 preadipocyte proliferation. To determine whether the function of C/EBP β and Kdm4b is specific to MCE, the effects of knocking down C/EBP β and Kdm4b on the proliferation of normal 3T3-L1 preadipocyte were examined. Knockdown of C/EBP β or Kdm4b neither significantly altered the growth rate of pre-confluent 3T3-L1 cell, as assessed by cell number analysis, nor cell cycle phase (Figures 8a and b, respectively). Furthermore, only weak binding of C/EBP β or Kdm4b on the promoters of the four cell cycle genes was detected in pre-confluent 3T3-L1 cells

(Figures 8c and d, respectively). Consistently, the expression of the four cell cycle genes was not affected when C/EBP β or Kdm4b was knocked down (Figure 8e). These observations indicate that C/EBP β and Kdm4b has a specific role in MCE, but not in normal 3T3-L1 preadipocyte proliferation.

Discussion

C/EBP β is a basic leucine zipper transcription factor expressed in various cell types and has been characterized as a regulator of differentiation, such as the differentiation of adipocytes, the female reproductive system, mammary gland and bone.^{21–23} While these studies have shed light on its role in differentiation, some other studies also show that C/EBP β functions in cell proliferation of some experimental and neoplastic models. The mechanistic insight into C/EBP β action in those types of cell proliferation, however, is limited.

Kdm4b was originally shown to antagonize H3K9me3 at pericentric heterochromatin in mammalian cells.²⁴ Recent reports reveal that Kdm4b also functions in regulating gene expression. In ER signaling, Kdm4b demethylates H3K9me3 in the regulatory regions of ER responsive genes, thereby releasing a transcriptional block and promoting breast cancer cell proliferation and mammary gland development.^{16–18} Our findings have identified a specific role for Kdm4b in MCE during the proliferative phase of adipocyte differentiation. Kdm4b was found to be a *bona fide* target gene of C/EBP β during MCE. C/EBP β upregulates Kdm4b




Figure 6 Kdm4b acted as a co-factor of C/EBP β to activate the promoters of the four cell cycle genes. (a) ChIP analysis indicated that Kdm4b was also associated with the promoters of the four cell cycle genes. (b) At 20 h post-induction, soluble chromatin was prepared for ChIPs and Re-ChIPs with the indicated antibodies. Recovery of the four cell cycle gene promoters was analyzed by PCR. (c, d) Interaction between C/EBP β and Kdm4b. At 20 h post-induction, cell lysates were harvested and immunoprecipitated with antibody against C/EBP β (c) or Kdm4b (d). Immunoprecipitate complexes were detected by western blotting with the indicated antibodies. The band detected by the C/EBP β antibody is liver-enriched activator protein (LAP) (34 kDa). (e) 3T3-L1 preadipocytes were transfected with reporter constructs with/without the indicated expression vectors. After 48 h, luciferase activity was analyzed and plotted. * $P < 0.05$




Figure 7 Kdm4b regulates the expression of C/EBP β -targeted cell cycle genes through demethylation of H3K9me3. (a–d) Decrease of H3K9me3 modification level on the promoters of the four cell cycle genes during MCE. Upper panel: Schematic representation of the genomic locus. The regions from 'a' to 'd' were analyzed by ChIP-qPCR. The TSS is indicated by the arrow. Lower panel: ChIP-qPCR analysis of H3K9me3 occupancy on the sites in the gene loci indicated in the upper panel was performed at 0 h and 20 h after induction. Primers for ChIP-qPCR were available upon requested. (e) 3T3-L1 preadipocytes were treated with control RNAi or Kdm4b RNAi and then induced to differentiation. Cells were collected at times indicated and subjected to ChIP-qPCR analysis of H3K9me3 modification at position 'c' of each gene promoter loci. Data were normalized to 0 h time point of control RNAi-treated cells. (f) 3T3-L1 preadipocytes were treated as in (e). The relative mRNA level of C/EBP β -targeted cell cycle genes at times indicated was analyzed by RT-qPCR

expression (Figure 2), which in turn demethylates H3K9me3 associated with C/EBP β -regulated cell cycle genes, thereby activating the expression of these genes (Figure 7). Specifically, C/EBP β drives the expression of the four cell cycle genes via two pathways, one through transcriptional activation and the other through an epigenetic pathway

(Supplementary Figure S7). C/EBP β requires Kdm4b to reverse H3K9me3 modification thereby enhancing its transcriptional activity. Moreover, Kdm4b requires C/EBP β to be transcriptionally induced. This finding provides new insights into potential roles of C/EBP β in epigenetic and transcriptional regulation during MCE of preadipocyte differentiation.




Figure 8 Knockdown the expression of C/EBP β and Kdm4b has no effect on pre-confluent 3T3-L1 cell proliferation. (a) 3T3-L1 preadipocytes were plated at 20% confluence, next day the cells were transfected with C/EBP β or Kdm4b RNAi. Forty-eight hours later cell proliferation was determined by cell number counting at times indicated. (b) Cells were treated as in (a). At 48 h after RNAi transfection, cell cycle phase profiles were analyzed by PI staining followed by flow cytometry. (c, d) ChIP-qPCR analysis of C/EBP β (c) and Kdm4b (d) binding on the promoters of mouse Cdc45l, Mcm3, Gins1 and Cdc25c was performed in pre-confluent 3T3-L1 cells. (e) Forty-eight hours after transfection of C/EBP β or Kdm4b RNAi, cell lysates were harvested and the expression of the indicated genes was assessed by western blotting

Of interest, C/EBP β is also required for mammary gland development. Deletion of C/EBP β results in retarded proliferation of MECs and severe inhibition of lobuloalveolar development,²⁵ which is similar to the phenotype of conditional deletion of Kdm4b in mammary epithelium.¹⁷ Our findings suggest a collaboration between C/EBP β and Kdm4b during mammary gland development.

In the current study, four cell cycle genes were identified as direct targets of C/EBP β during MCE. Mcm3 is a component of MCM2-7 (mini-chromosome maintenance proteins 2–7) complex while Gins1 is a subunit of GINS (go-ichi-ni-san) complex.^{26,27} Cdc45l, MCM2-7 and GINS form a large complex referred to as CMG, which is found at the sites of DNA unwinding during replication and contains an intrinsic helicase activity.²⁸ The CMG complex is involved in the regulation of eukaryotic chromosomal DNA replication and is required for the initiation and elongation steps, as well as the co-ordination of DNA replication.²⁹ Cdc25c is a phosphotyrosine phosphatase belonging to the Cdc25 phosphatase family. It directs dephosphorylation of cyclin B-bound Cdc2 and triggers entry into mitosis.³⁰ A study indicates that Cdc25c not only has a role in mitosis, but also contributes to S-phase entry.³¹ Our results verify that the four cell cycle genes are downstream target genes of C/EBP β during MCE (Figures 4, 5; Supplementary Figure S5–S6). Knockdown of the four genes impairs MCE, followed by the inhibition of terminal adipocyte differentiation (Supplementary Figure S5). This observation lends additional credence to the idea that MCE is an early essential step in the adipocyte differentiation program of 3T3-L1 cells.

Propidium iodide (PI) staining and EdU labeling experiments indicate that the inhibition of MCE by the knockdown of Kdm4b and the four cell cycle genes is followed by a blockade of the G1/S transition (Figures 3b and c; Supplementary Figure S5b–S5c). These findings are also consistent with the role of C/EBP β in MCE, as functional disruption of C/EBP β blocks G1/S transition followed by the inhibition of MCE.^{5,11} Histone acetyltransferase binding to ORC1 (HBO1) is known

to positively regulate the initiation of DNA replication.³² Knockdown of HBO1 impairs the ability of 3T3-L1 preadipocytes to differentiate into mature adipocytes by inhibiting DNA replication and MCE.³³ In addition, aphidicolin and roscovitine, which are S-phase inhibitors, block DNA synthesis, MCE and adipogenesis in the process of 3T3-L1 adipocyte differentiation.^{5,6} Therefore, our present results together with previous reports, suggest that the regulation of G1/S transition is a key step in MCE. This hypothesis supports the notion that DNA replication during MCE increases the accessibility of promoter or enhancer elements to factors required for transcription of genes involved in the initiation of differentiation.³⁴

As proposed in this study, Kdm4b serves as a co-factor for C/EBP β to regulate cell cycle genes. To get a global sense of the potential collaboration between C/EBP β and Kdm4b during MCE, promoter-wide search of Kdm4b binding sites was performed. This analysis identified 379 Kdm4b binding regions at 20 h after 3T3-L1 preadipocytes were induced to differentiation (Supplementary Table S6). By integrating the C/EBP β and Kdm4b ChIP-on-chip data, 101 regions, which correspond to 119 genes, were found to be co-occupied by C/EBP β and Kdm4b (Supplementary Figure S8 and Table S7). Furthermore, 30 genes of the 68 potential C/EBP β target genes were also targeted by Kdm4b, including the four cell cycle genes described in this study (Supplementary Table S8). We have previously reported that histone H4, which is involved in chromatin assembly and is required for normal progression of S phase, is transactivated by C/EBP β during MCE.¹⁰ Intriguingly, histone H4 is also targeted by Kdm4b, as illustrated in the ChIP-on-chip data (Supplementary Table S6 and Table S8). Consistently, our study verifies that Kdm4b positively regulates the expression of histone H4 through demethylation of H3K9me3 on its promoter (Supplementary Figure S9). Thus, C/EBP β and Kdm4b co-regulate the expression of cell cycle genes, as well as chromatin assembly genes, such as histone H4, to efficiently promote MCE.

Knockdown of C/EBP β or Kdm4b does not appear to impair pre-confluent 3T3-L1 preadipocytes proliferation (Figure 8). This suggests that C/EBP β and Kdm4b are not general regulators of cell proliferation, but have specific functions in certain types of cell proliferation including MCE. Unlike the strong promoter bindings during MCE, C/EBP β and Kdm4b bind weakly to the promoters of the four cell cycle genes in pre-confluent 3T3-L1 cells (Figures 8c and d). It is possible that additional co-factors or specific post-translational modifications of C/EBP β , which might be absent in pre-confluent 3T3-L1 cells, are required for efficient binding of C/EBP β and Kdm4b to these promoters during MCE.

In conclusion, our data indicate that C/EBP β transactivates Kdm4b, which regulates the epigenetic signature in C/EBP β binding regions leading to the induction of cell cycle genes and chromatin assembly genes and thereby promoting MCE (Supplementary Figure S7). It is conceivable that through upregulation of Kdm4b, C/EBP β and Kdm4b form a feed forward regulatory loop to facilitate MCE. This suggests a profound role for C/EBP β in the epigenetic control of MCE and offers new insights into the critical action of C/EBP β in the proliferation of certain specific cell types.

Materials and Methods

Cell culture, induction of differentiation. 3T3-L1 preadipocytes were propagated and maintained in DMEM containing 10% calf serum and HEK293T cells were cultured in DMEM supplemented with 10% fetal bovine serum (FBS). To induce differentiation, 2-day post-confluent 3T3-L1 preadipocytes (designated 0 day or 0 h) were fed DMEM containing 10% FBS, 1 μ g/ml insulin, 1 μ M dexamethasone, and 0.5 mM 3-isobutyl-1-methylxanthine (MIX) until day 2. Cells were then fed DMEM supplemented with 10% FBS and 1 μ g/ml insulin for 2 days, after which they were fed every other day with DMEM containing 10% FBS. Adipocyte gene expression and acquisition of the adipocyte phenotype begins on day 3 and is maximal by day 8. Cell number was determined by hemocytometer.

Oil red O staining. Oil red O staining was conducted as described.¹⁹ Cells were washed three times with PBS and then fixed for 10 min with 3.7% formaldehyde. Oil red O (0.5% in isopropanol) was diluted with water (3:2), filtered through a 0.45 μ m filter, and incubated with the fixed cells for 1 h at room temperature. Cells were washed with water, and the stained fat droplets in the cells were visualized by light microscopy and photographed.

ChIP and Re-ChIP. Cells were fixed with 1% formaldehyde for 10 min at room temperature. Glycine was added to a final concentration of 0.125 M, and the incubation was continued for additional 5 min. Cells were then washed twice with ice-cold PBS, harvested by scraping, pelleted, and resuspended in 1 ml SDS lysis buffer containing 50 mM Tris-HCl pH 8.0, 1% SDS, 10 mM EDTA, and protease inhibitors. Samples were sonicated eight times for 30 s each with an interval of 30 s with a Bioruptor sonicator (Diagenode, Liège, Belgium). Samples were centrifuged at 14 000 \times g at 4 $^{\circ}$ C for 10 min. After removal of an input aliquot (whole-cell extract), supernates were diluted 10-fold in ChIP dilution buffer containing 20 mM Tris-HCl pH 8.0, 150 mM NaCl, 2 mM EDTA, 1% TritonX-100, and complete protease inhibitor tablets. Samples were immunoprecipitated using antibodies against C/EBP β (sc-150 and sc-7962, Santa Cruz Biotechnologies, Santa Cruz, CA, USA), Kdm4b (A301-477A, Bethyl, Montgomery, AL, USA), H3K9me3 (ab8898, Abcam, Cambridge, MA, USA) or the non-specific IgG control (Abcam). Immunoprecipitated samples were eluted and reverse cross-linked by incubation overnight at 65 $^{\circ}$ C. Genomic DNA was then extracted using a PCR purification kit (Qiagen, Valencia, CA, USA). Primers for ChIP-qPCR were in the Supplementary Table S9. ChIP data were normalized to control IgG or expressed as percentage of input. Re-ChIPs were done essentially the same as primary ChIPs. Beads from the first ChIP were incubated with 10 mM DTT at 37 $^{\circ}$ C for 30 min and the eluted immunocomplex were diluted 1:50 in dilution buffer followed by re-immunoprecipitation with the second antibodies.

ChIP-on-chip arrays and data analysis. ChIP-on-chip analysis was performed as described.³⁵ C/EBP β antibody (sc-150, Santa Cruz Biotechnologies) and Kdm4b antibody (A301-477A, Bethyl) were used for the ChIPs. ChIP samples were then amplified by ligation-mediated PCR, labeled and hybridized to Affymetrix mouse promoter 1.0 oligonucleotide tiling arrays (Affymetrix, Santa Clara, CA, USA, for C/EBP β) or Nimblegen mouse 385 K RefSeq promoter arrays (Roche NimbleGen, Madison, WI, USA, for Kdm4b). ChIP-on-chip assays were analyzed with MAT, with probes remapped to the February 2006 Mouse Genome Assembly (mm8). The threshold cutoffs for binding regions were $FDR \leq 1\%$. For each binding region, the nearest gene was determined, and the distance from the center of the binding region to the TSSs of the gene was calculated using CEAS^{35,36} according to gene coordinates in the February 2006 Mouse Genome Assembly (mm8). For conservation analysis, the ChIP regions were aligned at the center and uniformly expanded to 1500 bp at each end, and at each position the average phastCons score retrieved from UCSC Genome Browser (<http://genome.ucsc.edu>) was generated for the average conservation plot.

Analysis of the enriched motifs. An enumeration-based *de novo* motif discovery method was performed on the ChIP-on-chip data.³⁷ Z-score was used to assess the significance of motif enrichment. $Z - score = (C - E) / \sqrt{E}$, quantifying the significance of enrichment. Here, *C* and *E* represent the observed and expected numbers of instances in ChIP regions, respectively. Background regions were randomly selected from design regions of Affymetrix mouse Promoter array. Enrichment of known transcription factor motifs in the TRANSFAC database was evaluated by Z-scores. Using a matrix similarity threshold of 0.8, the numbers of instances in ChIP and background regions were counted for 554 annotated vertebrate PWMs.

GO analysis. GO analysis was performed using DAVID (<http://david.abcc.ncifcrf.gov>), by examining the biological processes in which these genes are involved, and using RefSeq mRNA IDs with the entire mouse genome as a background model.^{35,38} P-values were calculated using Fisher Exact statistics.

RNA interference. Synthetic RNAi oligonucleotides specific for regions in the C/EBP β , Kdm4b, Cdc45l, Mcm3, Gins1, Cdc25c mRNAs were designed and synthesized by Invitrogen (Carlsbad, CA, USA) Stealth RNAi. The silencing effects of several RNAi oligonucleotides were screened and tested initially for their ability to knock down the expression of these genes by RT-qPCR. The sequences for successful RNAi knockdown were listed in the Supplementary Table S10. Stealth RNAi Negative Control Duplexes with a similar GC content were used as a negative control. 3T3-L1 cells were transfected at $\sim 50\%$ confluence with RNAi oligonucleotides using Lipofectamine RNAiMAX (Invitrogen) according to the manufacturer's instructions.

RNA isolation, qPCR analysis, and gene expression profiling. Total RNAs were extracted using Trizol (Invitrogen). First-strand cDNAs were synthesized using the PrimeScript reverse transcriptase and Random primers (Takara Bio, Otsu, Japan). qPCR was performed using primers as described. Power SYBR Green PCR Mastermix (Applied Biosystems, Carlsbad, CA, USA) and the PRISM 7300 instrument (Applied Biosystems) were used. Analysis was performed using the standard curve method and normalization of all genes of interest to the control gene 18S rRNA. Primers for RT-qPCR were available in the Supplementary Table S9. RT-qPCR data were normalized to 0 h time point of control RNAi-treated cells. Gene expression profiling was carried out by KangChen Bio-tech (Shanghai, China). At 0 h or 20 h after induction, total RNA from 3T3-L1 cells was extracted, amplified, labeled and hybridized using the NimbleGen system (Roche NimbleGen).

Western blotting. Cells were lysed with lysis buffer containing 2% SDS, 10 mM dithiothreitol, 50 mM Tris-HCl, pH 6.8, 10% glycerol, 0.002% bromophenol blue, 1 \times protease inhibitor mixture. Equal amounts of protein were separated by SDS-PAGE, and transferred to PVDF membrane (Millipore, Billerica, MA, USA), immunoblotted with antibodies. Antibodies recognizing the following proteins were used: PPAR γ (Santa Cruz Biotechnologies), C/EBP α (Santa Cruz Biotechnologies), 422/ap2 (prepared in Dr. Lane's lab), Cdc45l (GeneTex, Irvine, CA, USA), Mcm3 (GeneTex), Gins1 (Abgent, San Diego, CA, USA), Cdc25c (Santa Cruz Biotechnologies), C/EBP β (sc-150, Santa Cruz Biotechnologies), Kdm4b (A301-477A, Bethyl) and Hsp90 (Abcam).

EdU labeling, microscopic fluorescence, PI staining, and flow cytometry analysis. EdU labeling was performed as previously described.³⁹

Cells were induced to differentiation with standard protocol. Sixteen hours after induction, cells were labeled for 2 h with 10 μ M EdU before fixation with 4% paraformaldehyde. EdU was stained by incubating for 20 min with 100 mM Tris (from 2 M stock, pH 8.5), 0.5 mM CuSO₄, 10 μ M fluorescent azide (from 100 mM stocks in DMSO), and 50 mM ascorbic acid (added last to the mix from a 0.5 M stock in water). The staining mix was prepared fresh each time and used for staining cells immediately after addition of ascorbate. Cells were then washed and counterstained with Hoechst. The incorporation of EdU was visualized by fluorescence microscopy. At each step described above, the cells were washed with PBS three times after treatment. For FACS analysis, EdU-labeled cells were trypsinized, treated with EdU-staining chemicals, washed and subjected to flow cytometry with FACScan (Becton Dickinson, Franklin Lakes, NJ, USA). For PI staining, cells were trypsinized, fixed with 2% (wt/vol) paraformaldehyde in 1 \times PBS. Then they were treated with 0.5 mg/ml RNase A for 1 h at room temperature and incubated with 0.1 mg/ml PI (Sigma, St. Louis, MO, USA) for 45 min at 37 $^{\circ}$ C. DNA content was determined by flow cytometry analysis.

Electrophoretic Mobility Shift Assay. Nuclei were isolated and nuclear extracts prepared using the buffer containing 0.3 M NaCl, 1 M urea, 1% Nonidet P-40, 25 mM HEPES (pH 7.9), and 1 mM DTT. Protein concentration was determined by the Bradford method (Bio-Rad, Hercules, CA, USA). EMSA was performed as follows. Reaction mixtures containing \sim 0.25 ng of ³²P-labeled oligonucleotide probe, 2 μ g of poly [d(I-C)], and 10 μ g of nuclear extract protein in 30 μ l of buffer (10 mM HEPES, 0.1 mM EDTA, 5% glycerol, 100 mM NaCl, 0.3 M urea, 0.3% NP-40) were incubated on ice for 15 min, at room temperature for 15 min, and then were separated electrophoretically on 5% polyacrylamide gels 0.5 \times TBE [44.5 mM Tris, 44.5 mM boric acid, 1 mM EDTA (pH 8.3)]. For the competition assays, 100-fold excess of unlabeled probes were added to the binding mixtures. In all, 2 μ g of antibodies against C/EBP β (sc-7962, Santa Cruz Biotechnology) were used in the supershift analyses. Oligonucleotides used are listed in the Supplementary Table S11.

Luciferase reporter assays. Promoter regions of mouse Kdm4b (–1000 to –1 bp), Cdc45l (–1000 to +208 bp), Mcm3 (–500 to +23 bp), Gins1 (–258 to +105 bp), Cdc25c (–500 to +145 bp) were amplified via PCR from genomic DNA of 3T3-L1 (primer sequences are available upon request) and cloned into PGL3-basic (Promega, Madison, WI, USA). Cells were transfected by Lipofectamine 2000 (Invitrogen). Luciferase activity was measured using Dual Luciferase Reporter Assay (Promega), normalizing firefly luciferase to renilla activity.

Immunoprecipitation assay. Cells were washed with PBS, scraped off and collected by centrifugation. Then cells were suspended in RIPA buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS) in the presence of protease inhibitors (Roche) for 3 h at 4 $^{\circ}$ C. After centrifugation, the supernates were incubated with the indicated antibodies at 4 $^{\circ}$ C overnight. The next day, protein A-agarose beads (Invitrogen) was added. After 3 h incubation, the beads were washed with TBS-T (TBS + 0.05% Tween-20). The immunoprecipitates were separated by SDS-PAGE and subjected to western blotting.

Statistical analysis. Results are expressed as mean \pm S.E. Comparisons between groups were made by unpaired two-tailed Student's *t*-test, where $P < 0.05$ was considered as statistically significant. All experiments were repeated at least three times and representative data are shown.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. This research is supported partially by National Key Basic Research Project Grant 2011CB910201 and 2009CB825604, The State Key Program of National Natural Science Foundation 31030048C120114, Shanghai Key Science and Technology Research Project 10JC1401000 (For QQT); National Natural Science Foundation Grant 30870510 (For XL), National Natural Science Foundation Grant 31000603 (For LG); The Department is supported by Shanghai Leading Academic Discipline Project B110 and 985 Project 985III-YFX0302. We thank KangChen Bio-tech and Dr. Kan-kan Wang for ChIP-on-chip assays.

- Haslam DW, James WP. Obesity. *Lancet* 2005; **366**: 1197–1209.
- Shepherd PR, Gnudi L, Tozzo E, Yang H, Leach F, Kahn BB. Adipose cell hyperplasia and enhanced glucose disposal in transgenic mice overexpressing GLUT4 selectively in adipose tissue. *J Biol Chem* 1993; **268**: 22243–22246.
- Green H, Kehinde O. An established preadipose cell line and its differentiation in culture. II. Factors affecting the adipose conversion. *Cell* 1975; **5**: 19–27.
- Qiu Z, Wei Y, Chen N, Jiang M, Wu J, Liao K. DNA synthesis and mitotic clonal expansion is not a required step for 3T3-L1 preadipocyte differentiation into adipocytes. *J Biol Chem* 2001; **276**: 11988–11995.
- Tang QQ, Otto TC, Lane MD. Mitotic clonal expansion: a synchronous process required for adipogenesis. *Proc Natl Acad Sci USA* 2003; **100**: 44–49.
- Reichert M, Eick D. Analysis of cell cycle arrest in adipocyte differentiation. *Oncogene* 1999; **18**: 459–466.
- Yeh WC, Bierer BE, McKnight SL. Rapamycin inhibits clonal expansion and adipogenic differentiation of 3T3-L1 cells. *Proc Natl Acad Sci USA* 1995; **92**: 11086–11090.
- Farmer SR. Transcriptional control of adipocyte formation. *Cell Metab* 2006; **4**: 263–273.
- Tang QQ, Otto TC, Lane MD. CCAAT/enhancer-binding protein beta is required for mitotic clonal expansion during adipogenesis. *Proc Natl Acad Sci USA* 2003; **100**: 850–855.
- Zhang YY, Li X, Qian SW, Guo L, Huang HY, He Q *et al*. Transcriptional activation of histone H4 by C/EBPbeta during the mitotic clonal expansion of 3T3-L1 adipocyte differentiation. *Mol Biol Cell* 2011; **22**: 2165–2174.
- Zhang JW, Tang QQ, Vinson C, Lane MD. Dominant-negative C/EBP disrupts mitotic clonal expansion and differentiation of 3T3-L1 preadipocytes. *Proc Natl Acad Sci USA* 2004; **101**: 43–47.
- Zhu S, Yoon K, Sterneck E, Johnson PF, Smart RC. CCAAT/enhancer binding protein-beta is a mediator of keratinocyte survival and skin tumorigenesis involving oncogenic Ras signaling. *Proc Natl Acad Sci USA* 2002; **99**: 207–212.
- Iyer VV, Kadakia TB, McCabe LR, Schwartz RC. CCAAT/enhancer-binding protein-beta has a role in osteoblast proliferation and differentiation. *Exp Cell Res* 2004; **295**: 128–137.
- Seagroves TN, Krnacik S, Raught B, Gay J, Burgess-Beusse B, Darlington GJ *et al*. C/EBPbeta, but not C/EBPalpha, is essential for ductal morphogenesis, lobuloalveolar proliferation, and functional differentiation in the mouse mammary gland. *Genes Dev* 1998; **12**: 1917–1928.
- Shuman JD, Sebastian T, Kaldis P, Copeland TD, Zhu S, Smart RC *et al*. Cell cycle-dependent phosphorylation of C/EBPbeta mediates oncogenic cooperativity between C/EBPbeta and H-RasV12. *Mol Cell Biol* 2004; **24**: 7380–7391.
- Shi L, Sun L, Li Q, Liang J, Yu W, Yi X *et al*. Histone demethylase JMJD2B coordinates H3K4/H3K9 methylation and promotes hormonally responsive breast carcinogenesis. *Proc Natl Acad Sci USA* 2011; **108**: 7541–7546.
- Kawazu M, Saso K, Tong KI, McQuire T, Goto K, Son DO *et al*. Histone demethylase JMJD2B functions as a co-factor of estrogen receptor in breast cancer proliferation and mammary gland development. *PLoS One* 2011; **6**: e17830.
- Yang J, Jubb AM, Pike L, Buffa FM, Turley H, Baban D *et al*. The histone demethylase JMJD2B is regulated by estrogen receptor alpha and hypoxia, and is a key mediator of estrogen induced growth. *Cancer Res* 2010; **70**: 6456–6466.
- Tang QQ, Lane MD. Activation and centromeric localization of CCAAT/enhancer-binding proteins during the mitotic clonal expansion of adipocyte differentiation. *Genes Dev* 1999; **13**: 2231–2241.
- Peterson CL, Laniel MA. Histones and histone modifications. *Curr Biol* 2004; **14**: R546–R551.
- Sterneck E, Tessarollo L, Johnson PF. An essential role for C/EBPbeta in female reproduction. *Genes Dev* 1997; **11**: 2153–2162.
- Robinson GW, Johnson PF, Hennighausen L, Sterneck E. The C/EBPbeta transcription factor regulates epithelial cell proliferation and differentiation in the mammary gland. *Genes Dev* 1998; **12**: 1907–1916.
- Gutierrez S, Javed A, Tennant DK, van Rees M, Montecino M, Stein GS *et al*. CCAAT/enhancer-binding proteins (C/EBP) beta and delta activate osteocalcin gene transcription and synergize with Runx2 at the C/EBP element to regulate bone-specific expression. *J Biol Chem* 2002; **277**: 1316–1323.
- Fodor BD, Kubicek S, Yonezawa M, O'Sullivan RJ, Sengupta R, Perez-Burgos L *et al*. Jmjd2b antagonizes H3K9 trimethylation at pericentric heterochromatin in mammalian cells. *Genes Dev* 2006; **20**: 1557–1562.
- Grimm SL, Rosen JM. The role of C/EBPbeta in mammary gland development and breast cancer. *J Mammary Gland Biol Neoplasia* 2003; **8**: 191–204.
- Lin DI, Aggarwal P, Diehl JA. Phosphorylation of MCM3 on Ser-112 regulates its incorporation into the MCM2-7 complex. *Proc Natl Acad Sci USA* 2008; **105**: 8079–8084.
- Nagahama Y, Ueno M, Miyamoto S, Morii E, Minami T, Mochizuki N *et al*. PSF1, a DNA replication factor expressed widely in stem and progenitor cells, drives tumorigenic and metastatic properties. *Cancer Res* 2010; **70**: 1215–1224.
- Aparicio T, Ibarra A, Mendez J. Cdc45-MCM-GINS a new power player for DNA replication. *Cell Div* 2006; **1**: 18.
- Tercero JA, Labib K, Diffley JF. DNA synthesis at individual replication forks requires the essential initiation factor Cdc45p. *EMBO J* 2000; **19**: 2082–2093.

30. Aressy B, Ducommun B. Cell cycle control by the CDC25 phosphatases. *Anticancer Agents Med Chem* 2008; **8**: 818–824.
31. Turowski P, Franckhauser C, Morris MC, Vaglio P, Fernandez A, Lamb NJ. Functional cdc25C dual-specificity phosphatase is required for S-phase entry in human cells. *Mol Biol Cell* 2003; **14**: 2984–2998.
32. Doyon Y, Cayrou C, Ullah M, Landry AJ, Cote V, Selleck W *et al*. ING tumor suppressor proteins are critical regulators of chromatin acetylation required for genome expression and perpetuation. *Mol Cell* 2006; **21**: 51–64.
33. Johmura Y, Osada S, Nishizuka M, Imagawa M. FAD24 acts in concert with histone acetyltransferase HBO1 to promote adipogenesis by controlling DNA replication. *J Biol Chem* 2008; **283**: 2265–2274.
34. Cornelius P, MacDougald OA, Lane MD. Regulation of adipocyte development. *Annu Rev Nutr* 1994; **14**: 99–129.
35. Lefterova MI, Zhang Y, Steger DJ, Schupp M, Schug J, Cristancho A *et al*. PPAR γ and C/EBP factors orchestrate adipocyte biology via adjacent binding on a genome-wide scale. *Genes Dev* 2008; **22**: 2941–2952.
36. Ji X, Li W, Song J, Wei L, Liu XS. CEAS: cis-regulatory element annotation system. *Nucleic Acids Res*, 2006; **34**: W551–W554.
37. Wang K, Wang P, Shi J, Zhu X, He M, Jia X *et al*. PML/RAR α targets promoter regions containing PU.1 consensus and RARE half sites in acute promyelocytic leukemia. *Cancer Cell* 2010; **17**: 186–197.
38. Steger DJ, Grant GR, Schupp M, Tomaru T, Lefterova MI, Schug J *et al*. Propagation of adipogenic signals through an epigenomic transition state. *Genes Dev* 2010; **24**: 1035–1044.
39. Aparicio T, Guillou E, Coloma J, Montoya G, Mendez J. The human GINS complex associates with Cdc45 and MCM and is essential for DNA replication. *Nucleic Acids Res* 2009; **37**: 2087–2095.

Supplementary Information accompanies the paper on Cell Death and Differentiation website (<http://www.nature.com/cdd>)