Review

Novel roles for hERG K⁺ channels in cell proliferation and apoptosis

J Jehle¹, PA Schweizer¹, HA Katus¹ and D Thomas^{*,1}

The human ether-a-go-go-related gene potassium channel (hERG, Kv11.1, KCNH2) has an essential role in cardiac action potential repolarization. Electrical dysfunction of the voltage-sensitive ion channel is associated with potentially lethal ventricular arrhythmias in humans. hERG K⁺ channels are also expressed in a variety of cancer cells where they control cell proliferation and apoptosis. In this review, we discuss molecular mechanisms of hERG-associated cell cycle regulation and cell death. In addition, the significance of hERG K⁺ channels as future drug target in anticancer therapy is highlighted.

Cell Death and Disease (2011) 2, e193; doi:10.1038/cddis.2011.77; published online 18 August 2011

Subject Category: Cancer

Ion Channels Involved in Cell Proliferation and Death

Ion channels have been implicated in signaling pathways leading to cell proliferation or apoptosis (programmed cell death). Their identification and functional characterization in tumor cells suggest potential significance in anticancer therapy. Transient receptor potential channels form a superfamily of ubiquitously expressed channels influencing the balance between cell survival and death.^{1,2} In addition, hyperpolarization-activated cyclic nucleotide-gated channels were detected in embryonic stem cells where they exert proliferative effects. Potassium channels represent the largest group of channels involved in cell death and proliferation.^{3,4} Calcium-activated K_{Ca}3.1 channels contribute to proliferation and atherosclerosis, and inhibition of the current attenuates fibrosis and lymphocyte proliferation.^{5–8} Furthermore, voltage-gated K⁺ channels (e.g. Kv1.3) or two-pore-domain channels (e.g. K_{2P}5.1) determine growth of adenocarcinomas.^{9,10} Voltage-sensitive human ether-a-go-go-related gene (hERG) potassium channels have recently emerged as novel regulators of growth and death in cancer cells. This review focuses on hERG channels in proliferation and apoptosis. Current knowledge on expression, function and regulation is reviewed, and clinical implications are discussed.

Differential Expression of hERG Potassium Channels

Cardiac expression and function of hERG K⁺ channels.

Repolarization of cardiac ventricular myocytes is mainly regulated by outward potassium currents. One of the most important currents is the delayed rectifier potassium current,

I_K, which has rapidly and slowly activating components (*I_{Kr}* and *I_{Ks}*).¹¹ Activation of the rapid component of the delayed rectifier potassium current, *I_{Kr}*, terminates the plateau phase and initiates repolarization of the cardiac action potential. The hERG encodes the voltage-gated potassium channel α -subunit underlying *I_{Kr}*.^{12–14} hERG potassium channels form homo-tetramers of identical six transmembrane spanning domains, with a cluster of positive charges localized in the S4 domain serving as voltage sensor. hERG channels are a primary target for the pharmacological management of arrhythmias with class III antiarrhythmic agents.^{15,16} Blockade of hERG currents causes lengthening of the cardiac action potential, which may produce a beneficial class III antiarrhythmic effect. Excessive reduction of hERG currents due to mutations in hERG or *via* blockade produces chromosome-7-linked congenital long QT syndrome (LQTS-2) and acquired long QT syndrome, respectively. Both forms of LQTS are associated with delayed cardiac repolarization, prolonged electrocardiographic QT intervals, and a risk for the development of ventricular ‘torsade de pointes’ arrhythmias and sudden cardiac death. hERG channels are inhibited by a variety of non-antiarrhythmic compounds. This undesirable side effect is now considered a significant hurdle in the development of new and safer drugs, and has forced removal of several drugs from the market. In addition to LQTS, cardiomyocyte apoptosis has been reported following pharmacological hERG K⁺ channel blockade.¹⁷

hERG K⁺ channels in cancer. Various cancer cell lines of epithelial, neuronal, leukemic, and connective tissue origin express hERG K⁺ channels (Table 1), whereas corresponding non-cancerous cells and cell lines do not

¹Department of Cardiology, Medical University Hospital, Heidelberg, Germany

*Corresponding author: D Thomas, Department of Cardiology, Medical University Hospital, Heidelberg, Im Neuenheimer Feld 410, D-69120 Heidelberg, Germany. Tel: +49 6221 568855; Fax: +49 6221 565514; E-mail: dierk.thomas@med.uni-heidelberg.de

Keywords: hERG; K⁺ channel; cell proliferation; apoptosis; anticancer therapy

Abbreviations: ALLHAT, Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial; AML, acute myeloid leukemia; BPH, benign prostatic hyperplasia; CHOP, c/EBP homologous protein; DOC-1, downstream of CHOP-1; FAK, focal adhesion kinase; hERG, human ether-a-go-go-related gene; Kv, voltage-gated potassium channel; LQTS, long QT syndrome; TNF α , tumor necrosis factor α

Received 20.5.11; revised 14.7.11; accepted 18.7.11; Edited by V De Laurenzi

Table 1 Cells and cell lines expressing hERG[illegible]

Table 1 (Continued)

Derivation	Cell type/cell line	Comment
Neuronal	<i>Blood</i> PBCD34 peripheral blood, hemopoietic progenitor cells ²⁴	Rapid induction of hERG expression by cytokines/growth factors
	<i>Uterus</i> Myometrium ²³	
	<i>Brain</i> Rat brain ⁴⁴	
	Mouse brain ⁴⁴	
	Human prolactin-secreting adenoma cells ⁴⁵	
	Native rat lactotroph cells ⁴⁵	
Non-cancerous cell lines		
Connective and soft tissue	<i>Heart</i> HL-1 (murine atrial tumor cell line) ⁴⁶	

exhibit significant hERG protein levels. In corresponding human cancers, hERG protein may serve as biomarkers of malignant transition. Furthermore, hERG expression is implicated in enhanced cell proliferation, invasiveness, lymph node dissemination, and reduced cell differentiation and prognosis.^{21,22} In addition, increased neoangiogenesis, another hallmark of malignant tissue growth, has been reported for glioblastoma where the generation of blood vessels was stimulated by hERG-dependent secretion of vascular endothelial growth factor.²⁷

Differential hERG expression patterns during ontogenesis.

While hERG expression in normal adult human tissue is limited to heart, brain, myometrium, pancreas, and hematopoietic progenitors, other species have been described to undergo changes in their ERG expression profile during ontogenesis: quail embryos express ERG K⁺ channels in peripheral ganglia and skeletal muscle in addition to heart and central nervous system.⁴⁷ This observation illustrates that hERG expression in tumor cells might either represent ectopic re-expression of a gene that remains silent in differentiated cells, or reflect re-activation of embryonic genes, which is well recognized in cancers.³⁵

Cell Proliferation

Functional role of hERG K⁺ channels in cell proliferation. In differentiated adult cells, resting membrane potential varies from −40 mV to about −90 mV.⁴⁸ These distinct differences are closely correlated to the proliferative potential of respective cell types, ranging from slowly proliferating or non-proliferative neurons or muscle cells (−70 mV to −90 mV) to highly proliferative glandular epithelia of liver, thyroid, pancreas, or salivary glands (−40 mV to −55 mV).⁴⁸ hERG K⁺ channels are closed at membrane potentials below a threshold of ~−60 mV¹ whereas classical inwardly rectifying channels remain open at more negative membrane potentials.⁴⁹ The predominance of hERG in cycling cells may thus account for the depolarized resting membrane potential in these cells.³¹ The membrane potential of cycling cells is particularly depolarized during the G1 phase. However, K⁺ channel-dependent hyperpolarization appears to be critical for progression to the S phase. Hyperpolarization evokes

Ca²⁺ influx, which is further augmented by calcium-dependent K⁺ (K_{Ca}) channels and permits synthesis of mitogenic factors. In addition, hyperpolarization provides the electrical gradient necessary for Na⁺-dependent transport of metabolic substrates and ions across the plasma membrane, which is required for DNA synthesis.⁵⁰ Considering that K⁺ channels are involved in cell cycle progression, abundant expression of K⁺ channels is expected to cause loss of proliferative control if endogenous pathways fail to block excessively expressed K⁺ channels.⁵⁰ Interestingly, the promoter region of the hERG gene harbors multiple binding sites for oncoproteins, such as specificity protein 1 and nuclear factor kappa light chain enhancer of activated B-cells, and for the tumor suppressor protein Nkx3.1 (Nk3 homeobox 1).³⁰ We may hypothesize that mutations in oncoproteins constitutively activate hERG gene expression, shifting resting membrane potentials of cancerous cells toward more depolarized values and repolarizing them at the end of G1 phase, thereby facilitating cell cycle progression and thus leading to cell proliferation. Here, pharmacological intervention using hERG antagonists will serve to arrest the cell cycle in the G1 phase. Furthermore, human gastric cancer cells exhibit reduced levels of the regulatory β -subunit KCNE2, leading to hERG current increase.^{51,52} In addition, genetic deletion of KCNE2 is associated with gastric neoplasia and increased nuclear cyclin D1 levels in mice, revealing genetic manipulation of cell proliferation mediated by a hERG β -subunit.⁵²

Various cancer cell lines and cardiomyocytes have been reported to express an N terminally truncated splice variant of hERG, hERG1b, that confers specific electrophysiological properties.⁵³ Pharmacological approaches targeting the hERG1/hERG1b ratio may modulate the resting membrane potential of cycling cells. Increased hERG1b levels are expected to depolarize cells, while high hERG1 levels will shift membrane potential toward more hyperpolarized values³⁵ and suppress cell proliferation.

hERG potassium channel blockers modulate proliferation.

Leukemic cell lines express hERG K⁺ channels whereas non-cancerous lymphocytes do not exhibit hERG protein. Selective hERG channel blockade by E-4031 reduced proliferation in cancerous cell lines.²⁵ Unspecific deceleration of the cell cycle and reduction of cell proliferation⁵⁰ were ruled out in

Table 2 Cell cycle arrest induced by hERG K⁺ channel inhibitors

Cell type	hERG blocker	Comment
Human osteoclast/preosteoclast cells FLG 29.1 ²⁴	E-4031; WAY 123398; CsCl	Arrest in G1 phase
Human leukemia cell lines K562 and HL60 ⁵⁴	E-4031	Arrest in G1 phase
Human neuroblastoma SH-SY5Y ³⁶	HERG1/1b shRNA	Arrest in G1 phase
Human gastric cancer cell line SGC7901 ²¹	HERG-specific siRNA	Arrest in G1 phase
Murine corticotroph AtT20 cells ⁵⁵	Doxazosin	Arrest in G1 phase
Rat somatotroph GH3 cells ⁵⁵	Doxazosin	Arrest in G1 phase
MCF-7 breast cancer cell line ⁵⁶	Astemizole	Arrest in G1 phase
Human colon carcinoma cell line HT-29 ²⁹	Erythromycin (+vincristine)	Potential of the effect of vincristine (arrest in G2/M phase)
Prostate cancer cell line LNCaP ⁵⁷	Doxazosin (25 μ M); terazosin (25 μ M)	No antiproliferative effect, no change in cell cycle distribution

mechanistic analyses, confirming specific cell cycle arrest as underlying mechanism. Cell cycle analysis of FLG29.1 leukemia cells revealed accumulation of cells in the G1 phase following treatment with hERG channel blockers.²⁴ Furthermore, additional structurally different hERG blockers have been shown to achieve cell cycle arrest in G1 phase of hERG-positive cells (Table 2). It is noteworthy that the hERG blocker erythromycin blocks cell cycle in G2 phase if administered together with vincristine.²⁹ In addition, hERG blockers doxazosin and terazosin did not cause cell cycle arrest despite hERG expression in distinct cell lines, for example, LNCaP prostate carcinoma cells.^{30,57}

Significance of hERG Ion Channels in Apoptosis

Proapoptotic effects of hERG K⁺ channel inhibitors. hERG channel blockers have been shown to induce apoptosis in different cell types. This mechanism is independent of their capacity to inhibit cell proliferation via cell cycle arrest. The significance of hERG K⁺ channels in apoptotic pathways has been demonstrated in hERG-transfected HEK293 cells, which underwent apoptosis upon administration of doxazosin, compared with control HEK293 cells lacking endogenous hERG.⁵⁸ Doxazosin is an α_1 -adrenoceptor antagonist with hERG-blocking properties that is clinically used as antihypertensive drug.⁵⁹ In the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT), which compared novel antihypertensive drugs to diuretic treatment in 33 000 patients, the doxazosin arm had to be discontinued due to an increase in congestive heart failure that may be attributed to cardiomyocyte apoptosis.^{60,61} The proapoptotic effect of doxazosin has been confirmed *in vitro* in the murine atrial tumor cell line HL-1 and in isolated adult human cardiomyocytes,¹⁷ providing a possible explanation for the increased incidence of congestive heart failure in the doxazosin arm of the ALLHAT trial. In addition to hypertension, doxazosin is used for treatment of lower urinary tract symptoms caused by benign prostatic hyperplasia (BPH). Smooth muscle relaxation due to α_1 -adrenergic blockade was initially thought to underlie the relief of symptoms in BPH patients. However, subsequent studies revealed an apoptotic effect of doxazosin in hyperplastic prostatic tissue that may contribute to its clinical efficacy.⁶² Furthermore, doxazosin induced apoptosis in

prostatic cancer cells.⁶³ Limitations arise from the lack of studies directly comparing hERG expression in normal, hyperplastic, and cancerous prostatic tissue, respectively. Finally, hERG channel expression is well documented in pituitary adenoma cells.⁴⁵ When treated with doxazosin *in vitro*, antiproliferative and proapoptotic effects were observed in pituitary adenoma cells independent of antiadrenergic properties of the drug.⁵⁵

Molecular mechanisms of hERG-associated apoptosis.

hERG K⁺ channel blockers such as doxazosin activate multiple apoptotic pathways. However, evidence for a direct mechanistic link between hERG K⁺ channels and apoptotic proteins remains sparse to date. In HL-1 cardiomyocytes, doxazosin induces apoptosis via the endoplasmic reticulum pathway, involving enhanced phosphorylation of p38 mitogen-activated protein kinase, which activates GADD153/CHOP (growth arrest and DNA damage-induced gene 153/c/EBP homologous protein). GADD153/CHOP subsequently forms heterodimers with DNA-binding protein c/EBP β (CCAAT enhancer-binding protein beta) and translocates into the nucleus, where it augments transcription of the carbonic anhydrase DOC-1 (downstream of CHOP-1). DOC-1 then acidifies intracellular pH and facilitates apoptosis.⁶⁴ Finally, the CHOP pathway results in activation of a key apoptotic enzyme, caspase 3.⁶⁵ Caspase activation by doxazosin induces cleavage of the protein-tyrosine kinase FAK (focal adhesion kinase) in HL-1 cells, which compromises cell adhesion and leads to apoptosis.⁶⁴ FAK is an essential component of integrin signaling and is phosphorylated when cells are adhered to the extracellular matrix. Thus, it provides a survival signal and prevents apoptosis.⁶⁶ In prostate cancer cells, FAK is cleaved by caspase 3 upon treatment with doxazosin, which leads to apoptosis or anoikis (i.e. apoptosis due to loss of cell adhesion).⁶⁷ Furthermore, hERG1, integrin β 1, and FAK form a macromolecular complex in hERG1-transfected HEK293 cells and SH-SY5Y neuroblastoma cells. Cell adhesion via integrin β 1 causes activation of hERG1, which is essential for direct FAK phosphorylation (Figure 1).³⁷

FAK and hERG overexpression have independently been related to enhanced dissemination and invasiveness of tumors.^{20,66} FAK phosphorylation due to hERG activation may explain the ability of malignant cells to circumvent apoptosis once they have lost contact to the extracellular

Figure 1 Pathways of hERG-associated apoptosis. Doxazosin induces apoptosis via two independent mechanisms, inhibition of FAK phosphorylation via blockade of hERG K⁺ channels³⁷ and caspase 3-mediated cleavage of FAK⁶⁷ via induction of ER stress,⁶⁴ respectively. In addition, DOC-1 causes a decrease in intracellular pH, which facilitates apoptosis⁶⁴

matrix. The abundant expression of hERG and FAK might provide crucial survival signals in the absence of cell adhesion, and thus account for increased invasiveness and dissemination of hERG-positive tumors. In addition, colocalization with hERG potassium channels activates the GTPase Rac1 and may contribute to adhesion-dependent modulation of tumor cell motility.³⁷

Cell type- and environment-specific effects on apoptosis are suggested by reports of hERG activity promoting apoptosis. In hERG-positive SKBr3, SH-SY5Y, and HL-1 cells, apoptosis occurs via a caspase 3-dependent pathway in response to extracellular administration of H₂O₂ or TNF α (tumor necrosis factor α), whereas selective inhibition of hERG conductance by dofetilide attenuates the proapoptotic effect of H₂O₂ and TNF α .³³ The methodology in the latter study is different from investigations mentioned above. Cells were first incubated with H₂O₂ or TNF α to induce apoptosis, followed by application of hERG blockers. In the same study, hERG is revealed to recruit TNF α receptor 1 to the plasma membrane, which might explain increased responsiveness to TNF α in these cells.³³ The authors describe a proliferative effect in hERG-expressing cells at low doses of TNF α and an antiapoptotic effect of the hERG inhibitor dofetilide upon pretreatment with H₂O₂ and TNF α . These observations appear to be at odds with proapoptotic effects of hERG K⁺ channel blockers. The hERG blocker doxazosin has been proven as a proapoptotic agent in a wide range of *in vitro* and *in vivo* studies. Doxazosin increases the intracellular H₂O₂ content in BPH stromal cells. This is considered to facilitate TNF α -related pathways.⁶⁸ Administration of H₂O₂ before hERG inhibition appears to interfere with hERG-induced

signaling pathways, which augment intracellular H₂O₂ levels. The antiapoptotic effect of hERG channel blockade may be due to this interference. However, pro- and antiapoptotic effects of hERG blockers might coexist, and proapoptotic effects, including the increase in intracellular H₂O₂, could outweigh a possible antiapoptotic effect through suppression of the apoptotic H₂O₂ – TNF α pathway. However, an unambiguous differentiation between effects of hERG conductance and hERG expression is lacking, and the mechanism by which hERG conductance facilitates H₂O₂- and TNF α -mediated apoptosis remains unclear at the molecular level.

Clinical and Therapeutic Implications

Diagnostic value of hERG K⁺ channel expression in tumors. hERG may be utilized as a potential tumor marker, given their expression in a variety of tumor cells and their absence from most non-cancerous human tissues. Specifically, hERG was detected in endometrial cancer at mRNA (sensitivity = 67%; *n* = 18) and protein levels (sensitivity = 82%; *n* = 18), whereas only 18% (*n* = 11) of non-cancerous endometrial samples exhibited hERG mRNA or protein.²³ In colon carcinomas, hERG mRNA was a more sensitive and more specific indicator for malignancy (100% sensitivity and specificity; *n* = 23) than mRNA of the established tumor markers CEA (sensitivity = 94.4%; *n* = 18), CK19 (sensitivity = 77.8%; *n* = 18), or CK20 (sensitivity = 94.4%; *n* = 18).¹⁸ Immunohistochemical staining for hERG protein

reached similar sensitivity and specificity as hERG mRNA.¹⁸ Further validation is required in larger patient populations.

Prognostic significance of hERG K⁺ channel expression in tumors. The prognostic value of hERG expression in tumors has been evaluated in several tissues. In acute myeloid leukemia (AML) blasts, hERG K⁺ channel expression is associated with a 50% reduction of relapse-free and overall survival time compared with patients with hERG-negative AML (12 *versus* 23 months).⁶⁹ Patients with esophageal squamous cell carcinomas similarly exhibit reduced survival (30 *versus* 56 months) when hERG is detected.²² However, hERG K⁺ channel expression was not significantly associated with invasiveness, dissemination, or tumor grade in this study. In gastric cancer cells, levels of hERG expression are positively correlated to tumor differentiation and TNM stage.²¹ Moreover, tumor growth was observed in BALB/c nu/nu mice following injection of gastric cancer cells. Injection of cancer cells that were pretreated with hERG siRNA significantly attenuated tumorigenesis,²¹ confirming the pathological significance of hERG in tumor growth and suggesting a potential novel target in anticancer therapy (see below). In colonic adenocarcinomas, there is a significant correlation between hERG K⁺ channel expression and invasiveness or dissemination. hERG is not detected in normal colonic mucosa (0%; *n* = 60) and rarely observed in adenoma (9%; *n* = 11). In contrast, substantial hERG was found in patients with non-metastatic adenocarcinoma (75%; *n* = 52) and metastatic adenocarcinoma (100%; *n* = 8), with the most pronounced staining found in hepatic and peritoneal metastasis.²⁰

Anticancer therapy. The antihypertensive α_1 -adrenoceptor blocker doxazosin is an established treatment option in BPH. Its therapeutic efficacy has been attributed to induction of apoptosis in hyperplastic and cancerous prostate cells.⁵⁷ Furthermore, hERG-positive cancer cells have been reported to be particularly susceptible to chemotherapeutics vincristine, paclitaxel, and hydroxycamptothecin.²⁹ Direct effects of vincristine, paclitaxel, and hydroxycamptothecin on hERG channels remain to be investigated. Erythromycin, a macrolide antibiotic with hERG-blocking properties, further enhances the antiproliferative effect of these chemotherapeutics.²⁹ The most intriguing perspective of anticancer therapy targeting hERG channels is direct blockade of the potassium channel, which is expected to produce antiproliferative and proapoptotic effects that diminish tumor growth and invasiveness. The first proof of concept study confirmed prevention of gastric cancer cell proliferation by the hERG K⁺ channel blocker cisapride.⁷⁰ A systematic *in vivo* investigation of chemotherapeutic properties and potential cardiac side effects of hERG inhibitors is required.

Potential side effects and limitations of anticancer therapy based on hERG current inhibition. Proarrhythmic¹⁴ and cardiotoxic risks of hERG inhibitors require careful evaluation⁷ when applying these compounds in clinical oncology. Systemic treatment of cancers with hERG antagonists may affect cardiac myocytes, resulting in

apoptosis and heart failure. In addition, application of hERG antagonists may induce QT prolongation and ventricular tachycardia. Although cancer treatment usually occurs in life-threatening situations, and in some cases potential cardiac damage is accepted (e.g. during use of anthracyclines), optimal suppression of these events will be required. To prevent proarrhythmic side effects, short-term drug application may be sufficient to induce apoptosis in tumor cells with minimal effects on cardiac electrophysiology. ECG monitoring should be performed during application of the drug. Additional pharmacological inhibition of cardiac L-type calcium channels or β -adrenoceptors may offset the limiting proarrhythmic effects of hERG channel inhibitors.^{71–73} Cardiomyocyte apoptosis may be circumvented through targeted delivery techniques such as direct injection or trans-arterial drug application. Gene therapy represents an additional therapeutic approach to targeted suppression of hERG channel expression in cancers. Different proliferative states of cardiac and tumor cells may render cancerous tissue more susceptible to proapoptotic and antiproliferative stimuli, reducing the overall risk of heart failure during systemic application of hERG antagonists. Feasibility of tumor-selective hERG-based anticancer therapy will further depend on differential drug effects on cancerous and non-cancerous tissue expressing hERG K⁺ channels.

Conclusion

hERG potassium channels, previously recognized to promote cardiac action potential repolarization, are now revealed to serve as regulators of proliferation and apoptosis in cancer cells. Their significance in anticancer therapy is supported by mechanistic data and preliminary *in vivo* studies. Limitations arise from potential cardiac side effects that require attention. Further studies are warranted to provide a more complete understanding of hERG effects on apoptotic pathways. Downstream signaling proteins may serve as more specific therapeutic drug targets in future anticancer therapy.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. This study was supported in part by research grants from the ADUMED foundation (to DT), the German Heart Foundation/German Foundation of Heart Research (to DT), and the Max-Planck-Society (TANDEM project to PAS).

- Shapovalov G, Lehen'kyi V, Skryma R, Prevarskaya N. TRP channels in cell survival and cell death in normal and transformed cells. *Cell Calcium* 2011; e-pub ahead of print 29 May 2011; doi:10.1016/j.ceca.2011.05.006.
- Lau YT, Wong CK, Luo J, Leung LH, Tsang PF, Bian ZX *et al*. Effects of hyperpolarization-activated cyclic nucleotide-gated (HCN) channel blockers on the proliferation and cell cycle progression of embryonic stem cells. *Pflügers Arch* 2011; **461**: 191–202.
- Shieh CC, Coghlan M, Sullivan JP, Gopalakrishnan M. Potassium channels: molecular defects, diseases, and therapeutic opportunities. *Pharmacol Rev* 2000; **52**: 557–594.
- Lang F, Föllmer M, Lang KS, Lang PA, Ritter M, Gulbins E *et al*. Ion channels in cell proliferation and apoptotic cell death. *J Membr Biol* 2005; **205**: 147–157.
- Chou CC, Lunn CA, Murgolo NJ. KCa3.1: target and marker for cancer, autoimmune disorder and vascular inflammation? *Expert Rev Mol Diagn* 2008; **8**: 179–187.

6. Srivastava S, Zhdanova O, Di L, Li Z, Albaquimi M, Wulff H *et al*. Protein histidine phosphatase 1 negatively regulates CD4 T cells by inhibiting the K⁺ channel KCa3.1. *Proc Natl Acad Sci USA* 2008; **105**: 14442–14446.
7. Toyama K, Wulff H, Chandy KG, Azam P, Raman G, Saito T *et al*. The intermediate-conductance calcium-activated potassium channel KCa3.1 contributes to atherogenesis in mice and humans. *J Clin Invest* 2008; **118**: 3025–3037.
8. Grgic I, Kiss E, Kaistha BP, Busch C, Kloss M, Sautter J *et al*. Renal fibrosis is attenuated by targeted disruption of KCa3.1 potassium channels. *Proc Natl Acad Sci USA* 2009; **106**: 14518–14523.
9. Jang SH, Choi SY, Ryu PD, Lee SY. Anti-proliferative effect of Kv1.3 blockers in A549 human lung adenocarcinoma *in vitro* and *in vivo*. *Eur J Pharmacol* 2011; **651**: 26–32.
10. Alvarez-Baron CP, Jonsson P, Thomas C, Dryer SE, Williams C. The two-pore domain potassium channel KCNK5: induction by estrogen receptor [alpha] and role in proliferation of breast cancer cells. *Mol Endocrinol* 2011; **25**: 1326–1336.
11. Sanguinetti MC, Jurkiewicz NK. Two components of delayed rectifier K⁺ current. *J Gen Physiol* 1990; **96**: 195–215.
12. Warmke JW, Ganetzky B. A family of potassium channel genes related to eag in *Drosophila* and mammals. *Proc Natl Acad Sci USA* 1994; **91**: 3438–3442.
13. Sanguinetti MC, Jiang C, Curran ME, Keating MT. A mechanistic link between an inherited and an acquired cardiac arrhythmia: HERG encodes the I_{Kr} potassium channel. *Cell* 1995; **81**: 299–307.
14. Sanguinetti MC, Tristani-Firouzi M. hERG potassium channels and cardiac arrhythmia. *Nature* 2006; **440**: 463–469.
15. Thomas D, Karle CA, Kiehn J. The cardiac hERG/I_{Kr} potassium channel as pharmacological target: structure, function, regulation, and clinical applications. *Curr Pharm Des* 2006; **12**: 2271–2283.
16. Staudacher I, Schweizer PA, Katus HA, Thomas D. hERG: protein trafficking and potential for therapy and drug side effects. *Curr Opin Drug Discov Devel* 2010; **13**: 23–30.
17. Gonzalez-Juanatey JR, Iglesias MJ, Alcáide C, Pineiro R, Lago F. Doxazosin induces apoptosis in cardiomyocytes cultured *in vitro* by a mechanism that is independent of alpha1-adrenergic blockade. *Circulation* 2003; **107**: 127–131.
18. Dolderer JH, Schuldes H, Bockhorn H, Altmannberger M, Lambers C, von Zabern D *et al*. HERG1 gene expression as a specific tumor marker in colorectal tissues. *Eur J Surg Oncol* 2010; **36**: 72–77.
19. Gong JH, Liu XJ, Shang BY, Chen SZ, Zhen YS. HERG K⁺ channel related chemosensitivity to sparfloxacin in colon cancer cells. *Oncol Rep* 2010; **23**: 1747–1756.
20. Lastraioli E, Guasti L, Crociani O, Polvani S, Hofmann G, Witchel H *et al*. herg1 gene and HERG1 protein are overexpressed in colorectal cancers and regulate cell invasion of tumor cells. *Cancer Res* 2004; **64**: 606–611.
21. Shao XD, Wu KC, Guo XZ, Xie MJ, Zhang J, Fan DM. Expression and significance of HERG protein in gastric cancer. *Cancer Biol Ther* 2008; **7**: 45–50.
22. Ding XW, Luo HS, Luo B, Xu DQ, Gao S. Overexpression of HERG1 in resected esophageal squamous cell carcinomas: a marker for poor prognosis. *J Surg Oncol* 2008; **97**: 57–62.
23. Cherubini A, Taddei GL, Crociani O, Paglierani M, Buccoliero AM, Fontana L *et al*. HERG potassium channels are more frequently expressed in human endometrial cancer as compared to non-cancerous endometrium. *Br J Cancer* 2000; **83**: 1722–1729.
24. Pillozzi S, Brizzi MF, Balzi M, Crociani O, Cherubini A, Guasti L *et al*. HERG potassium channels are constitutively expressed in primary human acute myeloid leukemias and regulate cell proliferation of normal and leukemic hemopoietic progenitors. *Leukemia* 2002; **16**: 1791–1798.
25. Smith GA, Tsui HW, Newell EW, Jiang X, Zhu XP, Tsui FW *et al*. Functional up-regulation of HERG K⁺ channels in neoplastic hematopoietic cells. *J Biol Chem* 2002; **277**: 18528–18534.
26. Li H, Liu L, Guo L, Zhang J, Du W, Li X *et al*. HERG K⁺ channel expression in CD34+/CD38–/CD123(high) cells and primary leukemia cells and analysis of its regulation in leukemia cells. *Int J Hematol* 2008; **87**: 387–392.
27. Masi A, Becchetti A, Restano-Cassulini R, Polvani S, Hofmann G, Buccoliero AM *et al*. hERG1 channels are overexpressed in glioblastoma multiforme and modulate VEGF secretion in glioblastoma cell lines. *Br J Cancer* 2005; **93**: 781–792.
28. Patt S, Preussat K, Beetz C, Kraft R, Schrey M, Kalff R *et al*. Expression of ether a go-go potassium channels in human gliomas. *Neurosci Lett* 2004; **368**: 249–253.
29. Chen SZ, Jiang M, Zhen YS. HERG K⁺ channel expression-related chemosensitivity in cancer cells and its modulation by erythromycin. *Cancer Chemother Pharmacol* 2005; **56**: 212–220.
30. Lin H, Xiao J, Luo X, Wang H, Gao H, Yang B *et al*. Overexpression HERG K(+) channel gene mediates cell-growth signals on activation of oncoproteins SP1 and NF-kappaB and inhibition of tumor suppressor Nkx3.1. *J Cell Physiol* 2007; **212**: 137–147.
31. Bianchi L, Wible B, Arcangeli A, Tagliatela M, Morra F, Castaldo P *et al*. herg encodes a K⁺ current highly conserved in tumors of different histogenesis: a selective advantage for cancer cells? *Cancer Res* 1998; **58**: 815–822.
32. Roy J, Vantol B, Cowley EA, Blay J, Linsdell P. Pharmacological separation of hEAG and hERG K⁺ channel function in the human mammary carcinoma cell line MCF-7. *Oncol Rep* 2008; **19**: 1511–1516.
33. Wang H, Zhang Y, Cao L, Han H, Wang J, Yang B *et al*. HERG K⁺ channel, a regulator of tumor cell apoptosis and proliferation. *Cancer Res* 2002; **62**: 4843–4848.
34. Afrasiabi E, Hietamaki M, Viitanen T, Sukumaran P, Bergelin N, Tornquist K. Expression and significance of HERG (KCNH2) potassium channels in the regulation of MDA-MB-435S melanoma cell proliferation and migration. *Cell Signal* 2010; **22**: 57–64.
35. Crociani O, Guasti L, Balzi M, Becchetti A, Wanke E, Olivetto M *et al*. Cell cycle-dependent expression of HERG1 and HERG1B isoforms in tumor cells. *J Biol Chem* 2003; **278**: 2947–2955.
36. Zhao J, Wei XL, Jia YS, Zheng JQ. Silencing of herg gene by shRNA inhibits SH-SY5Y cell growth *in vitro* and *in vivo*. *Eur J Pharmacol* 2008; **579**: 50–57.
37. Cherubini A, Hofmann G, Pillozzi S, Guasti L, Crociani O, Cilia E *et al*. Human ether-a-go-go-related gene 1 channels are physically linked to beta1 integrins and modulate adhesion-dependent signaling. *Mol Biol Cell* 2005; **16**: 2972–2983.
38. D'Amico M, Biagiotti T, Fontana L, Restano-Cassulini R, Lasagna N, Arcangeli A *et al*. HERG current sustains a cardiac-type action potential in neuroblastoma S cells. *Biochem Biophys Res Commun* 2003; **302**: 101–108.
39. Meves H. Slowing of ERG current deactivation in NG108-15 cells by the histidine-specific reagent diethylpyrocarbonate. *Neuropharmacology* 2001; **41**: 220–228.
40. Pancrazio JJ, Ma W, Grant GM, Shaffer KM, Kao WY, Liu QY *et al*. A role for inwardly rectifying K⁺ channels in differentiation of NG108-15 neuroblastoma x glioma cells. *J Neurobiol* 1999; **38**: 466–474.
41. Hofmann G, Bernabei PA, Crociani O, Cherubini A, Guasti L, Pillozzi S *et al*. HERG K⁺ channels activation during beta(1) integrin-mediated adhesion to fibronectin induces an up-regulation of alpha(v)beta(3) integrin in the preosteoclastic leukemia cell line FLG 29.1. *J Biol Chem* 2001; **276**: 4923–4931.
42. Agarwal JR, Griesinger F, Stuhmer W, Pardo LA. The potassium channel Ether a go-go is a novel prognostic factor with functional relevance in acute myeloid leukemia. *Mol Cancer* 2010; **9**: 18.
43. Rosati B, Marchetti P, Crociani O, Lecchi M, Lupi R, Arcangeli A *et al*. Glucose- and arginine-induced insulin secretion by human pancreatic beta-cells: the role of HERG K(+) channels in firing and release. *FASEB J* 2000; **14**: 2601–2610.
44. Pond AL, Scheve BK, Benedict AT, Petrecca K, Wagoner RD, Shrier A *et al*. Expression of distinct ERG proteins in rat, mouse, and human heart. Relation to functional I(Kr) channels. *J Biol Chem* 2000; **275**: 5997–6006.
45. Bauer CK, Wulfsen I, Schäfer R, Glassmeier G, Wimmers S, Flitsch J *et al*. HERG K(+) currents in human prolactin-secreting adenoma cells. *Pflugers Arch* 2003; **445**: 589–600.
46. Staudacher I, Wang L, Wan X, Obers S, Wenzel W, Tristram F *et al*. hERG K⁺ channel expression during development of the nervous and muscular system of quail embryos. *Schmiedeberg's Arch Pharmacol* 2011; **383**: 119–139.
47. Crociani O, Cherubini A, Piccini E, Polvani S, Costa L, Fontana L *et al*. Erg gene(s) expression during development of the nervous and muscular system of quail embryos. *Mech Dev* 2000; **95**: 239–243.
48. Binggeli R, Weinstein RC. Membrane potentials and sodium channels: hypotheses for growth regulation and cancer formation based on changes in sodium channels and gap junctions. *J Theor Biol* 1986; **123**: 377–401.
49. Ishihara K, Hiraoka M. Gating mechanism of the cloned inward rectifier potassium channel from mouse heart. *J Membr Biol* 1994; **142**: 55–64.
50. Wonderlin WF, Strobl JS. Potassium channels, proliferation and G1 progression. *J Membr Biol* 1996; **154**: 91–107.
51. Abbott GW, Sesti F, Splawski I, Buck ME, Lehmann MH, Timothy KW *et al*. MiRP1 forms I_{Kr} potassium channels with HERG and is associated with cardiac arrhythmia. *Cell* 1999; **97**: 175–187.
52. Roeppke TK, Purtell K, King EC, La Perle KM, Lerner DJ, Abbott GW. Targeted deletion of Kcne2 causes gastritis cystica profunda and gastric neoplasia. *PLoS One* 2010; **5**: e11451.
53. Jones EMC, Roti Roti EC, Wang J, Delfosse SA, Robertson GA. Cardiac I_{Kr} channels minimally comprise HERG 1a and 1b subunits. *J Biol Chem* 2004; **279**: 44690–44694.
54. Li H, Liu L, Guo T, Zhang J, Li X, Du W *et al*. Expression and functional role of HERG1, K⁺ channels in leukemic cells and leukemic stem cells. *J Huazhong Univ Sci Technolog Med Sci* 2007; **27**: 257–260.
55. Fernando MA, Heaney AP. Alpha1-adrenergic receptor antagonists: novel therapy for pituitary adenomas. *Mol Endocrinol* 2005; **19**: 3085–3096.
56. Borowiec AS, Hague F, Harir N, Guenin S, Guerin F, Gouilleux F *et al*. IGF-1 activates hEAG K(+) channels through an Akt-dependent signaling pathway in breast cancer cells: role in cell proliferation. *J Cell Physiol* 2007; **212**: 690–701.
57. Benning CM, Kyprianou N. Quinazoline-derived alpha1-adrenoceptor antagonists induce prostate cancer cell apoptosis via an alpha1-adrenoceptor-independent action. *Cancer Res* 2002; **62**: 597–602.
58. Thomas D, Bloehs R, Koschny R, Ficker E, Sykora J, Kiehn J *et al*. Doxazosin induces apoptosis of cells expressing hERG K⁺ channels. *Eur J Pharmacol* 2008; **579**: 98–103.
59. Davey M. Mechanism of alpha blockade for blood pressure control. *Am J Cardiol* 1987; **59**: 18G–28G.
60. The ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. Major cardiovascular events in hypertensive patients randomized to doxazosin vs chlorthalidone: the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *JAMA* 2002; **283**: 1967–1975.
61. Kang PM, Izumo S. Apoptosis and heart failure: a critical review of the literature. *Circ Res* 2000; **86**: 1107–1113.

62. Kyprianou N, Vaughan TB, Michel MC. Apoptosis induction by doxazosin and other quinazoline α 1-adrenoceptor antagonists: a new mechanism for cancer treatment? *Naunyn Schmiedeberg's Arch Pharmacol* 2009; **380**: 473–477.
63. Partin JV, Anglin IE, Kyprianou N. Quinazoline-based α 1-adrenoceptor antagonists induce prostate cancer cell apoptosis via TGF- β signalling and I kappa B α induction. *Br J Cancer* 2003; **88**: 1615–1621.
64. Eiras S, Fernandez P, Pineiro R, Iglesias MJ, Gonzalez-Juanatey JR, Lago F. Doxazosin induces activation of GADD153 and cleavage of focal adhesion kinase in cardiomyocytes en route to apoptosis. *Cardiovasc Res* 2006; **71**: 118–128.
65. Oyadomari S, Mori M. Roles of CHOP/GADD153 in endoplasmic reticulum stress. *Cell Death Differ* 2004; **11**: 381–389.
66. Kornberg LJ. Focal adhesion kinase and its potential involvement in tumor invasion and metastasis. *Head Neck* 1998; **20**: 745–752.
67. Walden PD, Globina Y, Nieder A. Induction of anoikis by doxazosin in prostate cancer cells is associated with activation of caspase-3 and a reduction of focal adhesion kinase. *Urol Res* 2004; **32**: 261–265.
68. Zhao H, Lai F, Nonn L, Brooks JD, Peehl DM. Molecular targets of doxazosin in human prostatic stromal cells. *Prostate* 2005; **62**: 400–410.
69. Pillozzi S, Brizzi MF, Bernabei PA, Bartolozzi B, Caporale R, Basile V et al. VEGFR-1 (FLT-1), β 1 integrin, and hERG K⁺ channel for a macromolecular signaling complex in acute myeloid leukemia: role in cell migration and clinical outcome. *Blood* 2007; **110**: 1238–1250.
70. Shao XD, Wu KC, Hao ZM, Hong L, Zhang J, Fan DM. The potent inhibitory effects of cisapride, a specific blocker for human ether-a-go-go-related gene (HERG) channel, on gastric cancer cells. *Cancer Biol Ther* 2005; **4**: 295–301.
71. Thomas D, Wendt-Nordahl G, Röckl K, Ficker E, Brown AM, Kiehn J. High-affinity blockade of HERG human cardiac potassium channels by the novel antiarrhythmic drug BRL-32872. *J Pharmacol Exp Ther* 2001; **297**: 753–761.
72. Thomas D, Gut B, Wendt-Nordahl G, Kiehn J. The antidepressant drug fluoxetine is an inhibitor of human ether-a-go-go-related gene (HERG) potassium channels. *J Pharmacol Exp Ther* 2002; **300**: 543–548.
73. Zhang S, Zhou Z, Gong Q, Makielski JC, January CT. Mechanism of block and identification of the verapamil binding domain to HERG potassium channels. *Circ Res* 1999; **84**: 989–998.

Cell Death and Disease is an open-access journal published by **Nature Publishing Group**. This work is licensed under the **Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License**. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>