Correspondence

Mammalian ncRNA-disease repository: a global view of ncRNA-mediated disease network

Y Wang^{1,6}, L Chen^{1,6}, B Chen^{1,6}, X Li^{1,6}, J Kang¹, K Fan¹, Y Hu¹, J Xu¹, L Yi¹, J Yang¹, Y Huang¹, L Cheng², Y Li³, C Wang⁴, K Li¹, X Li^{*,1}, J Xu^{*,1,5} and D Wang^{*,1}

Cell Death and Disease (2013) 4, e765; doi:10.1038/cddis.2013.292; published online 8 August 2013

Subject Category: Cancer

Dear Editor

Recently, substantial studies have begun to explore the functional diversity and mechanistic roles of ncRNAs in mammals.¹ Now, it has become increasingly apparent that ncRNAs are involved in multiple major biological processes, such as developmental timing, fat metabolism and cell death.² Furthermore, the epigenetic and genetic defects in ncRNAs and their processing machinery have been implicated in the etiology of many forms of diseases.³ Several databases that documented the relevance of the microRNAs(miRNAs) to diseases have been constructed and provided useful results.^{4,5} However, miRNAs are just the tip of the iceberg, other ncRNAs such as long non-coding RNAs (lncRNAs), PIWI-interacting RNAs (piRNAs) and small nucleolar RNAs (snoRNAs) have also been demonstrated to contribute to diseases.^{3,6} Accumulated evidence suggest the diverse non-coding RNAs (ncRNAs) involved in a wide variety of diseases progression.^{3,6,7} It is a key challenge for understanding the precise behavior of diverse ncRNAs in mammalian diseases and deciphering the cross-regulations among disease-associated ncRNAs. Because there was no repository focused on diverse ncRNA-disease relationships in mammals, we have developed a manually curated diverse ncRNA-disease repository (MNDR, www.rna-society.org/mnldr/) by integrating evidence in three mammals. Totally, 807 lncRNA-associated, 229 miRNA-associated, 13 piRNA-associated and 100 snoRNA-associated entries for 1149 curated entries were documented for three mammals (866 *Homo sapiens*-associated, 251 *Mus musculus*-associated and 32 *Rattus norvegicus*-associated entries) (Table 1).

Recent investigations indicated there are complex regulations among diverse ncRNAs and protein-coding genes. Such as, *PTEN* gene and the *PTEN* pseudogenes (*ptenp1*, one of lncRNAs) share a high degree of sequence homology, changes in *ptenp1* expression levels indirectly affect *PTEN* expression by sequestering *PTEN*-targeting miRNAs.⁸

Table 1 The statistics of the ncRNA-disease entries in MNDR database

Species	lncRNA	miRNA	piRNA	snoRNA	Total
<i>Homo sapiens</i>	753	NA	13	100	866
<i>Mus musculus</i>	50	201	0	0	251
<i>Rattus norvegicus</i>	4	28	0	0	32
Total	807	229	13	100	1149

Thus understanding the mutual regulating pattern among diverse ncRNAs and protein-coding genes, particularly in disease conditions, is a key challenge. Thus, MNDR is not only a knowledge depository but providing us a good opportunity to view the ncRNA-mediated disease network globally (in visualization page: www.rna-society.org/mnldr/visualization.html). Diverse ncRNAs and interaction genes were represented as nodes and the regulations were denoted as edges. Based on such a simplified ncRNA-mediated disease network, interesting observations have been achieved. The result showed that snoRNA *htr*, as a hub node, has intensively linked to 21 interaction genes in the network. More important, through *BCL2*, *BCL2L1* and *BAX*, the snoRNA *htr* can communicate with the lncRNA *malat1* (Supplementary Figure 1). Another example is snoRNA *htr* and lncRNA *h19* are linked by *E2F1* and *MYC*. When combined with human disease-associated miRNA evidence from mir2disease database, lncRNAs, miRNAs and snoRNAs, together with their interaction/target genes, can be integrated into bigger expanding ncRNA-mediated disease network. The biggest sub-network has 129 nodes and 149 edges, involving 33 lncRNAs, 1 snoRNA, 19 miRNAs and 76 interaction protein-coding genes (Supplementary Figure 2). In this network, more regulations among diverse ncRNAs directly or indirectly via intermediate genes, lncRNA *dgcr5*, *har1a* and *har1b*, were connected with *hsa-mir-21* via intermediate gene *REST*. Interestingly, *hsa-mir-21* and snoRNA *htr* were linked by key anti-apoptosis gene *BCL2*. Similar results were observed that

¹College of Bioinformatics Science and Technology, Harbin Medical University, Harbin, China; ²State Key Laboratory of Emerging Infectious Diseases, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China; ³Institute of Cardiovascular Sciences and Key Laboratory of Molecular Cardiovascular Sciences, Ministry of Education, Peking University Health Science Center, Beijing, China; ⁴Department of General Surgery, Huaihe Hospital of Henan University, Kaifeng, China and ⁵College of Bioengineering, Henan University of Technology, Zhengzhou, China

*Corresponding author: D Wang, J Xu and X Li, College of Bioinformatics Science and Technology, Harbin Medical University, Harbin 150081, China. Fax: 86 045186615922; E-mail: wangdong@ems.hrbmu.edu.cn (WD) or xujz0451@gmail.com (XJ) or lixia@hrbmu.edu.cn (LX)

⁶These authors contributed equally to this work

lncRNA *dgcr5*, *har1a* and *har1b* can also communicate with snoRNA *htr* through alternative route *NFKB1-hsa-mir-9-REST*. Hence, according to current data, the two pivot protein-coding genes (*BCL2* and *NFKB1*) and several ncRNAs (lncRNA *malat1*, snoRNA *htr* and miRNA *hsa-mir-21*, *has-mir-9*) collectively play an important role in the ncRNA-mediated disease network (Supplementary Figure 2). Importantly, the crosstalk between lncRNA *malat1* and miRNA *hsa-mir-21* can be found conserved in mouse ncRNA-mediated disease network. Above observations indicated diverse ncRNAs could communicate with each other in disease state through some disease-associated genes in mammals, highlighting the complexity, conservative and plasticity of the regulatory relationships between diverse ncRNAs and protein-coding genes in diseases.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. This work was supported by the National Natural Science Foundation of China (31100901, 91129710, 61170154 and 61073136),

China Postdoctoral Science Foundation (2013M531064), Natural Science Foundation of Heilongjiang Province of China (QC2010012 and ZD201114), Heilongjiang Postdoctoral Foundation (LBH-Z12171), Henan University of Technology (2009BS040 and 11JCYJ11) and China Undergraduate Training Programs for Innovation (201210226016).

1. Frazer KA. *Genome Res* 2013; **22**: 1599–1601.
2. Qureshi IA, Mehler MF. *Nat Rev Neurosci* 2012; **13**: 528–541.
3. Esteller M. *Nat Rev Genet* 2011; **12**: 861–874.
4. Jiang Q *et al.* *Nucleic Acids Res* 2009; **37**(Database issue): D98–D104.
5. Xu J, Li YH. *Cell Death Differ* 2012; **19**: 1571.
6. Sana J *et al.* *J Transl Med* 2012; **10**: 103.
7. Li L *et al.* *Proc Natl Acad Sci USA* 2009; **106**: 12956–12961.
8. Poliseno L *et al.* *Nature* 2010; **465**: 1033–1038.

Cell Death and Disease is an open-access journal published by *Nature Publishing Group*. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies this paper on Cell Death and Disease website (<http://www.nature.com/cddis>)