Dual role of the p38 MAPK/cPLA₂ pathway in the regulation of platelet apoptosis induced by ABT-737 and strong platelet agonists

N Rukoyatkina^{1,2}, I Mindukshev², U Walter³ and S Gambaryan^{*,1,2}

p38 Mitogen-activated protein (MAP) kinase is involved in the apoptosis of nucleated cells. Although platelets are anucleated cells, apoptotic proteins have been shown to regulate platelet lifespan. However, the involvement of p38 MAP kinase in platelet apoptosis is not yet clearly defined. Therefore, we investigated the role of p38 MAP kinase in apoptosis induced by a mimetic of BH3-only proteins, ABT-737, and in apoptosis-like events induced by such strong platelet agonists as thrombin in combination with convulxin (Thr/Cvx), both of which result in p38 MAP kinase phosphorylation and activation. A p38 inhibitor (SB202190) inhibited the apoptotic events induced by ABT-737 but did not influence those induced by Thr/Cvx. The inhibitor also reduced the phosphorylation of cytosolic phospholipase A₂ (cPLA₂), an established p38 substrate, induced by ABT-737 or Thr/Cvx. ABT-737, but not Thr/Cvx, induced the caspase 3-dependent cleavage and inactivation of cPLA₂. Thus, p38 MAPK promotes ABT-737-induced apoptosis by inhibiting the cPLA₂/arachidonate pathway. We also show that arachidonic acid (AA) itself and in combination with Thr/Cvx or ABT-737 at low concentrations prevented apoptotic events, whereas at high concentrations it enhanced such events. Our data support the hypothesis that the p38 MAPK-triggered arachidonate pathway serves as a defense mechanism against apoptosis under physiological conditions.

Cell Death and Disease (2013) 4, e931; doi:10.1038/cddis.2013.459; published online 21 November 2013

Subject Category: Experimental Medicine

Platelets play a key role in normal and pathological hemostasis through their ability to rapidly adhere to activated or injured endothelium and subendothelial matrix proteins (platelet adhesion), and to other activated platelets (platelet aggregation).¹ Platelets are also considered as key mediators of thrombosis, vascular inflammation, and atherosclerosis.^{2,3}

Apoptosis play a significant role in platelet production from megakaryocytes, and circulating platelets contain many components of the apoptotic machinery.⁴ Indeed, cytochrome c, caspase 9, caspase 3, apoptotic protease-activating factor, and Bcl-2 family proteins (for example, BAK, BAX, Bcl-X_L, Bcl-2, Bcl_w, Bim, Bid) are all expressed in platelets.^{5–8}

A variety of stimuli can induce apoptotic or apoptotic-like events in platelets, such as ABT-737, thrombin, collagen, and A23187. ABT-737, a potent mimetic of Bcl-2 homology BH3-only proteins (including Bim, Bid, and other proteins that can be important for binding and neutralizing antiapoptotic Bcl-2 family proteins), induces caspase-dependent platelet apoptosis that is associated with BAX translocation from the cytosol to mitochondria and homo-oligomerization.⁷ In highly activated platelets, several structural and functional changes, such as phosphatidylserine (PS) externalization, cell shrinkage, loss of mitochondrial membrane potential ($\Delta\Psi_m$), microparticle (MP) formation, and cleavage of gelsolin and

protein kinase C- δ , are reminiscent of nucleated cell apoptosis.^{9–13} However, strong agonists either did not activate caspase 3¹⁴ or did so very weakly.^{15–17} A total of 13 types of cell death have been characterized for nucleated cells, including caspase-dependent and caspase-independent intrinsic apoptosis, autophagic cell death, necroptosis and others.¹⁸ Although highly activated platelets exhibit the characteristic features of dying cells, there is no consensus in the literature clearly defining the type of strong activated platelet death. Some authors define them as necrotic cells¹⁹ whereas others^{11,20,21} consider them as apoptotic platelets.

Mitogen-activated protein (MAP) kinases represent a family of threonine/tyrosine-activated serine/threonine kinases that control many cellular responses, such as proliferation, migration, differentiation, and apoptosis. In platelets, p38 MAP kinase is phosphorylated and activated by different physiological agonists, including thrombin, collagen, and thromboxane A₂ (TxA₂).^{22–25} However, the mechanisms of p38 activation and their downstream effects are controversial and have not been clearly defined. p38 has been suggested to regulate platelet adhesion to collagen²⁶ and aggregation.^{24,27} In contrast, using different p38 inhibitors, other authors did not find any significant effect of p38 on platelet activation.^{28–31}

¹Institute of Clinical Biochemistry and Pathobiochemistry, University of Würzburg, Würzburg D-97080, Germany; ²Sechenov Institute of Evolutionary Physiology and Biochemistry, Russian Academy of Sciences, St Petersburg 194223, Russia and ³Center for Thrombosis and Hemostasis (CTH), Johannes Gutenberg University, Mainz 55131, Germany

*Corresponding author: S Gambaryan, Institute of Clinical Biochemistry and Pathobiochemistry, University of Würzburg, Grombühlstr. 12, Würzburg D-97080, Germany. Tel: +49 931 32 93617; Fax: +49 931 32 93630; E-mail: gambaryan@klin-biochem.uni-wuerzburg.de

Keywords: platelet; apoptosis; p38 MAP kinase; cPLA₂

Abbreviations: AA, arachidonic acid; cPLA₂, cytosolic phospholipase A₂; $\Delta\Psi_m$, mitochondrial membrane potential; MP, microparticle; ROS, reactive oxygen species; Thr/Cvx, thrombin/convulxin; TxA₂, thromboxane A₂; WP, washed platelets

Received 16.9.13; revised 21.9.13; accepted 22.10.13; Edited by A Stephanou

One of the established p38 substrates in platelets is cytosolic phospholipase A₂ (cPLA₂), which is phosphorylated by p38 at serine505 in agonist-stimulated platelets.^{32–34} In platelets, cPLA₂ activity is responsible for the release of arachidonic acid (AA) from membrane phospholipids.³⁵ AA is then metabolized by cyclooxygenase to prostaglandins, which are further converted to TxA₂ by thromboxane synthase (TxS). In addition, AA is involved in the regulation of reactive oxygen species (ROS) production. cPLA₂-induced AA is a prominent requirement for the activation of NADPH oxidase and generation of ROS in phagocytes.³⁶ ROS may also be generated as a by-product during the oxidation of AA by cyclooxygenase or lipoxygenase.^{37,38} Therefore, platelet cPLA₂ is involved in the regulation of two different functions, ROS generation and eicosanoid production. Excess ROS generation and AA release play essential roles in the initiation of apoptosis in platelets.^{21,39} As p38/cPLA₂ is involved in the regulation of nucleated cell apoptosis^{40–43} and the possible involvement of p38/cPLA₂ in the induction of platelet apoptotic events is not known, we compared the mechanism of p38/cPLA₂-mediated signaling induced by platelet agonists and ABT-737.

In this study, we show that p38 plays a significant role in ABT-737-induced platelet apoptosis, whereas it has no effect on thrombin in combination with convulxin(Thr/Cvx)-induced apoptotic-like events. ABT-737, but not Thr/Cvx, induced the caspase-dependent cleavage and inactivation of cPLA₂. We also assessed multiple influences of AA/ROS on apoptotic-like events in platelets and suggest that the p38-triggered arachidonate pathway serves as a defense mechanism under physiological conditions.

Results

Activation of p38 MAP kinase plays a significant role in ABT-737-induced platelet apoptosis. To clarify the function of p38 in platelet apoptosis induced by two independent pathways, the BH3 mimetic compound ABT-737 was used as an activator of the BAK/BAX-caspase pathway and a combination of Thr/Cvx as agonists of the second caspase-independent pathway.

In our study, we used terms apoptosis-like events for platelets activated by strong agonists and apoptosis for ABT-737-induced platelet death. ABT-737 at concentrations ranging from 0.1 to 1 μ M induced dose- and time-dependent increases in Annexin V binding (Figures 1a and c). Simultaneously, ABT-737 (0.1–1 μ M) time dependently increased the phosphorylation of p38 (Figure 1b). In contrast to Thr/Cvx, ABT-737 did not activate α IIb β 3 integrins (Figure 1c) and did not induce MP formation (Figures 6a and b). The activation of p38 by thrombin and convulxin alone or in combination (Thr/Cvx) was transient, reaching a maximum after 5 min of incubation and then gradually decreasing to a level comparable with that of the control after 60 min (Figure 1d). Conversely, activation of p38 by ABT-737 was detectable only after 20 min of stimulation and remained constant even up to 90 min of stimulation (Figure 1a). A p38 inhibitor (SB202190, 1 μ M) significantly decreased the amount of Annexin V-positive platelets (67 $\pm$ 3% compared with

Figure 1 BH3 mimetic (ABT-737) phosphorylates p38. (a) FACS analysis of Annexin V binding and (c) PAC-1 binding (integrin α IIb β 3 activation, mean fluorescence, AU) to WP (1×10^7 /ml) incubated with ABT-737. Annexin V and PAC-1 binding are presented as the fold increase compared with the control taken as 1. Data are presented as means $\pm$ S.E.M, $n = 5$; $+ P < 0.05$ compared with the control. (b and d) Western blot analysis using WP (4×10^6 /ml) of p38 phosphorylation after incubation with thrombin (5 mU/ml) and convulxin (5 ng/ml) or ABT-737 (0.1–1 μ M) (data shown are representative of three independent experiments)

ABT-737-stimulated platelets taken as 100%) and prevented decreases in $\Delta\Psi_m$ ($32 \pm 4\%$) stimulated by ABT-737 (Figures 2a and b). Similarly, SB202190 inhibited the cleavage of procaspase 3 by $31 \pm 3\%$ (Figures 2c and d), indicating that p38 is partly involved in ABT-737-induced caspase 3 activation and apoptosis.

p38 MAP kinase is not involved in the generation of apoptotic-like events triggered by Thr/Cvx. We next investigated whether the activation of p38 is involved in Thr/Cvx-induced platelet death. The activation of platelets by Thr/Cvx significantly increased Annexin V binding 7.6 ± 1.5

compared with control 1 ± 0.14) and $\alpha\text{IIb}\beta 3$ integrin activation (Figure 3a). The inhibition of p38 (SB202190, $1 \mu\text{M}$), as assessed by the phosphorylation of its established substrate (HSP27) (Figures 3c and d), did not significantly inhibit Annexin V binding (Figure 3a) or decreases in $\Delta\Psi_m$ (Figure 3b) and had no effect on $\alpha\text{IIb}\beta 3$ integrin activation (Figure 3a), indicating that p38 is not involved in Thr/Cvx-induced platelet apoptotic-like events.

Platelet cPLA₂ cleavage by ABT-737 is caspase dependent.

The stimulation of platelets by Cvx and Thr/Cvx (both 5 min) and by ABT-737 (30 min) induced p38 and cPLA₂ phosphorylation, and the latter was partly (in the case of Cvx and Thr/Cvx) and fully (in the case of ABT-737) prevented by preincubation with SB202190 (Figures 4a and b). However, in contrast to Cvx and Thr/Cvx, ABT-737 induced the cleavage of cPLA₂, which was prevented by a procaspase 3 inhibitor (z-DEVD-fmk) at concentrations of 50–100 μM (Figures 4c and d). We then tested whether the cleavage of cPLA₂ would inhibit its activity. One of the main functions of cPLA₂ in platelets is associated with the production of thromboxane (TxA₂) TxS. ABT-737 ($1 \mu\text{M}$) induced only a weak activation of TxS (4 ± 0.3 -fold increase compared with the control taken as 1), Cvx and Thr/Cvx (Figure 4e) strongly activated TxS (20 ± 1.3 - and 52 ± 1.3 -fold, respectively). The inhibition of p38 (SB202190, $1 \mu\text{M}$) decreased the TxS activity induced by Cvx or Thr/Cvx to $75 \pm 1.7\%$ and $87 \pm 1.8\%$, respectively, compared with the agonist-induced activity taken as 100% (Figure 4f).

Arachidonic acid prevents apoptosis in platelets at low concentrations and induces apoptosis in platelets at high concentrations.

In nucleated cells, AA has been shown to contribute to mitochondrial dysfunction during apoptosis;^{44,45} however, AA was used at very high concentrations (more than $10 \mu\text{M}$) in these studies. The depletion of AA prevents apoptotic development in cold-stored platelets whereas incubation with extremely high concentrations of AA (50–100 μM) induces platelet apoptosis.³⁹ In the present study, we tested whether AA would contribute to apoptosis in freshly isolated platelets. Washed platelets (WP) were incubated with different concentrations of AA (10^{-7} to 10^{-4} M), and apoptosis was assessed by decreases in $\Delta\Psi_m$, PS exposure, and procaspase 3 cleavage. At high concentrations (10^{-5} and 10^{-4} M), AA did indeed induce a decline in $\Delta\Psi_m$ (Figure 5a) and an increase in Annexin V binding (Figure 5b) and cleavage of procaspase 3 after 24 h of incubation (Figure 5c). However, at low concentrations (10^{-6} and 10^{-7} M), AA appeared to have antiapoptotic effects: it did not affect Annexin V binding, slightly increased

Figure 2 p38 MAP kinase inhibition prevents ABT-737-induced platelet apoptosis (a) FACS analysis of Annexin V-positive platelets ($1 \times 10^7/\text{ml}$) and (b) mitochondrial membrane potential ($\Delta\Psi_m$) in WP incubated with ABT-737 ($0.1 \mu\text{M}$, 30 min) or pretreated (10 min, $1 \mu\text{M}$) with p38 inhibitor (SB202190). (c) Western blot and (d) densitometry analyses of caspase 3 in human WP ($4 \times 10^8/\text{ml}$) incubated with ABT-737 ($0.1 \mu\text{M}$) or pretreated (10 min, $1 \mu\text{M}$) with p38 inhibitor (SB202190). In (a), ABT-737 was taken as 100%; in (b) and (d), the control was taken as 100%. Data are presented as means $\pm$ S.E.M., $n = 4$; + $P < 0.05$ compared with the control, * $P < 0.05$ compared with ABT-737

Figure 3 p38 inhibitor does not prevent PS surface exposure, decreased $\Delta\Psi_m$, and integrin $\alpha IIb\beta 3$ activation in platelets stimulated with Thr/Cvx (a) FACS analysis of Annexin V-positive cells and PAC-1 binding and (b) $\Delta\Psi_m$ in WP (1×10^7 /ml) stimulated by Thr/Cvx or pretreated (10 min, $1 \mu\text{M}$) with p38 inhibitor (SB202190). In (a) and (d), Thr/Cvx was taken as 100%; in (b), control was taken as 100%. Data are presented as means $\pm$ S.E.M, $n=6$; + $P<0.05$ compared with the control, * $P<0.05$ compared with platelets incubated with Thr/Cvx. (c) Western blot and (d) densitometry analyses of p38 and HSP27 phosphorylation in WP (3×10^8 /ml) stimulated with Thr/Cvx or pretreated (10 min, $1 \mu\text{M}$) with p38 inhibitor (SB202190). The data shown are representative of three independent experiments

$\Delta\Psi_m$ (Figure 5a), and decreased the procaspase 3 cleavage (Figures 5d and e) induced by ABT-737 and procaspase 3 cleavage in stored platelets after 24 h (Figure 5c). At low concentrations, AA also decreased the Annexin V binding induced by Thr/Cvx and ABT-737. In accordance with these data, AA induced MP formation only at high concentrations (Figures 6a and b).

ROS generated by activated platelets are not directly correlated with apoptotic events. AA can be involved in the regulation of ROS production as a by-product during the oxidation of AA by cyclooxygenase or lipoxygenase.³⁷ In addition, AA can directly activate NADPH oxidase, inducing ROS generation.⁴⁶ As MP formation in platelets may be induced by ROS,^{47,48} we compared MP formation and ROS production in platelets activated by ABT-737 ($1 \mu\text{M}$), AA (10^{-6} and 10^{-4} M), and Thr/Cvx (5 mU/5 ng/ml). AA at a concentration of 10^{-4} M induced the highest amount of

MPs (2.7 ± 0.34 fold compared with the control taken as 1), and Thr/Cvx induced 1.7 ± 0.2 ; neither a low concentration of AA (10^{-6} M) nor $1 \mu\text{M}$ ABT-737 induced significant MP formation (Figures 6a and b).

ROS production was strongly induced by AA at a low but not at a high concentration (27 ± 5.6 and 3 ± 1.5 -fold respectively, compared with the control taken as 1) (Figure 6c). Thr/Cvx stimulation also significantly increased ROS production (20 ± 4 -fold, compared with the control taken as 1), whereas ABT-737 had no effect on platelet ROS production (Figure 6c).

Discussion

In nucleated cells, stress-activated p38 MAPK is involved in apoptosis, cytokine production, cytoskeleton reorganization, and transcriptional regulation. p38 MAPK also appears to play a role in the pathogenesis of heart ischemia, sepsis, arthritis,

Figure 4 ABT-737 cleavage of platelet cPLA₂ is caspase dependent. (a) Western blot and (b) densitometry analysis of p-cPLA₂ in WP incubated with Cvx (5 ng/ml), Thr/Cvx (5 mU/5 ng/ml), or ABT-737 (1 μM) or pretreated (10 min, 1 μM) with p38 inhibitor (SB202190). (c) Western blot and (d) densitometry analysis of cPLA₂ in WP after stimulation with ABT-737 or pretreated with the indicated concentrations of the caspase 3 inhibitor: z-DEVD. Western blot data are representative of three independent experiments of p38 and cPLA₂ phosphorylation in WP (4×10^8 /ml). (e and f) ELISA data for the TxS activity in WP (4×10^8 /ml) activated by Cvx (5 ng/ml, 5 min), Thr/Cvx (5 mU/5 ng/ml, 5 min), or ABT-737 (1 μM, 30 min) and after preincubation with SB202190 (1 μM, 10 min)

human immunodeficiency virus infection, and Alzheimer's disease.⁴⁹ In platelets, p38 MAPK has been shown to be activated by almost all platelet agonists, including thrombin, collagen, ADP, and TxA₂.^{22–25} Nonetheless, the involvement of p38 in the mechanisms that trigger apoptosis in platelets has remained unclear. Therefore, we investigated the functions of p38 in the regulation of apoptosis induced by ABT-737 and the apoptosis-like events induced by strong agonists (Thr/Cvx). We found that ABT-737 dose- and time-dependently resulted in the phosphorylation of p38 (Figure 1). The inhibition of p38 significantly prevented platelet apoptosis, as assessed by PS-positive cells, a decrease in mitochondrial potential, and cleavage of caspase 3 (Figure 2). cPLA₂ is an established p38 substrate in platelets,^{23,32–34} and we show that the ABT-737-induced activation of p38 triggered cPLA₂ phosphorylation. In nucleated cells undergoing apoptosis, cPLA₂ is cleaved and inactivated by caspase 3 activation.^{50,51} In platelets, cPLA₂ was also cleaved upon ABT-737 activation and inactivated by a caspase 3-dependent mechanism (Figures 4a and b). In contrast, Thr/Cvx treatment induced the phosphorylation of cPLA₂ but did not promote its degradation (Figure 4a). Consistent with these data,

Thr/Cvx strongly enhanced cPLA₂ activity compared with the effect of ABT-737 (Figure 4e). It is important to note that the inhibition of p38 significantly prevented the apoptosis induced by ABT-737 but had no effect on the apoptotic-like events triggered by Thr/Cvx stimulation (Figure 3). As ABT-737-induced platelet apoptosis was directly correlated with the cleavage and inactivation of cPLA₂ and because Thr/Cvx stimulation strongly activated cPLA₂, which was responsible for AA generation, we tested whether AA itself was involved in apoptosis. In nucleated cells, high concentrations of AA contribute to mitochondrial dysfunction and apoptotic cell death,^{44,45,52} and AA also promotes apoptosis in stored platelets.³⁹ In addition, the concentration of AA can increase up to 10-fold during cerebral or heart ischemia compared with normal physiological conditions.^{53,54} Therefore, we tested the effects of different AA concentrations on hallmarks of platelet apoptosis. AA at low concentrations (10^{-7} – 10^{-6} M) enhanced $\Delta\Psi_m$, but had no influence on PS externalization and did not cleave caspase 3. In contrast, AA at high concentrations (10^{-5} – 10^{-4} M) induced a decrease in $\Delta\Psi_m$, enhanced PS externalization, and activated caspase 3 (Figure 5). Thus, the amount of AA might be an essential

Figure 5 Arachidonic acid at low concentrations prevents, and at high concentrations induces apoptosis in platelets. (a) FACS analysis of $\Delta\Psi_m$ and (b) Annexin V binding in WP (1×10^7 /ml) stimulated with AA for 5 min at the indicated concentrations. (f) Inhibition of Annexin V binding by AA (10^{-7} – 10^{-6} M) in platelets stimulated with Thr/Cvx (5 min) and ABT-737 (1μ M, 30 min). (c and d) Western blot analysis (data shown are representative of three independent experiments) of caspase 3 in WP (4×10^8 /ml) after stimulation with AA at the indicated concentrations for 2 and 24 h (c) and after stimulation with ABT-737 alone and in combination with AA for 30 min (d) and (e) densitometry analysis of (d). In (e), the control was taken as 100%. Data are presented as means $\pm$ S.E.M, $n = 6$; + $P < 0.05$ compared with the control, * $P < 0.05$ compared with platelets incubated with ABT-737

factor for mediating pro or antiapoptotic-like effects in platelets. We next addressed whether different concentrations of AA could prevent or enhance the apoptotic events induced by Thr/Cvx and ABT-737. Accordingly, we found that AA at low concentrations (10^{-7} for Thr/Cvx and 10^{-6} for ABT-737) inhibited the apoptotic events induced by both stimuli (Figure 5). The higher concentrations of AA required for the inhibitory effects of ABT-737-induced apoptosis were most likely associated with the inactivation of cPLA₂ and consequently a reduced basal AA concentration in this case.

ROS are involved in cellular responses to stress⁵⁵ and may contribute to various cellular pathological events, including apoptosis⁵⁶ and, in particular, MP formation.^{47,48} We investigated whether MP formation induced by ABT-737, Thr/Cvx, and AA correlates with ROS production and found increases only in the Thr/Cvx- and 10^{-6} M AA-treated platelets (Figure 6c), whereas MPs were enhanced only in the Thr/Cvx- and 10^{-4} M AA-stimulated platelets (Figures 6a and b), indicating that ROS are most likely not directly linked to apoptotic events.

In summary, we show here that p38 MAP kinase plays a significant role in platelet apoptosis induced by ABT-737, but is not involved in the generation of the apoptotic-like events

triggered by physiological stimuli (Thr/Cvx). Consistently, under physiological conditions, the p38-triggered activation of cPLA₂, release of AA, and ROS generation may serve as mechanisms to protect cells from apoptosis. In contrast, a high concentration of AA (as often observed under pathological conditions) may promote proapoptotic events.

Materials and Methods

Materials. Thrombin was obtained from Roche (Mannheim, Germany), and convulxin (Cvx) (ligand of glycoprotein VI from the snake venom *Crotalus durissus terrificus*) was purchased from LOXO (Dossenheim, Germany). SB202190 and z-DEVD-fmk were purchased from Calbiochem (Schwalbach, Germany), and ABT-737 was provided by Selleck Chemicals (Munich, Germany). Antibodies against p38, phospho-p38, and actin were obtained from Sigma (Munich, Germany). Antibodies against phospho-HSP27, phospho-cPLA₂, cPLA₂, and caspase 3 were from Cell Signaling (Frankfurt, Germany). Anti-rabbit and anti-mouse IgG conjugated with horseradish peroxidase were obtained from Amersham, Pharmacia Biotech (Freiburg, Germany). The fluorescent dyes Annexin-V-PE and PAC-1 were obtained from BD-Bioscience (Heidelberg, Germany). JC1 was obtained from Invitrogen (Eugen, Germany), and H₂DCF-DA was purchased from Molecular Probes (Göttingen, Germany).

Platelet preparation. Human platelets were prepared and used as previously reported,⁵⁷ with minor modifications. Blood was obtained from healthy volunteers according to our institutional guidelines and the Declaration of Helsinki. Our studies with human platelets were approved and recently (24 September, 2008) reconfirmed by the local ethics committee of the University of Würzburg (Studies Number 67/92 and 114/04).

Blood was collected into 1/7 volume of ACD solution (12 mM citric acid, 15 mM sodium citrate, 25 mM D-glucose, and 2 μ M EGTA, final concentrations). Platelet rich plasma was obtained by a 5-min centrifugation at $330 \times g$. To reduce leukocyte contamination, Platelet rich plasma was diluted 1:1 with PBS and centrifuged at $240 \times g$ for 10 min. The supernatant was centrifuged for 10 min at $430 \times g$, and the pelleted platelets were washed once in CGS buffer (120 mM sodium chloride, 12.9 mM trisodium citrate, and 30 mM D-glucose, pH 6.5) and resuspended in HEPES buffer (150 mM sodium chloride, 5 mM potassium chloride, 1 mM magnesium chloride, 10 mM D-glucose, and 10 mM HEPES, pH 7.4) without stirring; 1 mM CaCl₂ was added directly before platelet stimulation.

WP were used at a concentration of 3×10^6 /ml for western blot analysis and TxS activity, and a concentration of 1×10^7 /ml was used for flow cytometry analysis.

Flow cytometry (FACS) analysis. FACS analysis was performed using a Becton Dickinson FACS Calibur with CELLQuest software, version 3.1f (Becton Dickinson, Heidelberg, Germany). For the detection of surface PS or activated α IIb β 3 integrins, WP (50 μ l) were labeled with Annexin V-PE or PAC-1-FITC (both 1:10 dilution) for 10 min at RT after agonist stimulation. The platelets were then diluted 10 times with Annexin V-binding solution (140 mM NaCl, 10 mM HEPES, and 2.5 mM CaCl₂) for Annexin V or PBS for PAC-1 and immediately analyzed by flow cytometry.

Analysis of mitochondrial membrane potential. The $\Delta\Psi_m$ in human platelets was analyzed with JC1 dye (1.5 μ M final concentration). JC1 was incubated with WP (50 μ l) for 10 min at RT, and the samples were diluted (1:10) in PBS and analyzed by FACS. Green fluorescence (FL1) and red fluorescence (FL2) were measured in logarithmic scales using voltage settings of 710 and 588, respectively. Compensation settings (FL1-%FL2 and FL2-%FL1) were performed using FITC and PE beads (BD-Bioscience). JC1 is sensitive to $\Delta\Psi_m$, and the ratio of fluorescence in FL2 to FL1 corresponds to changes in $\Delta\Psi_m$.

Measurement of ROS production. Intracellular ROS production was analyzed by H₂DCF-DA, as described previously.⁵⁸ Briefly, WP were preloaded with 10 μ M H₂DCF-DA for 30 min at 37 °C in HEPES buffer. H₂DCF-DA is a cell-permeable, nonfluorescent dye converted intracellularly to cell-impermeable H₂DCF by intracellular esterases; the dye can then be oxidized by ROS to the fluorescent species (DCF) that is measured by flow cytometry. After incubation with agonists, the reaction was stopped by diluting the platelets in PBS for FACS analysis.

Measurement of thromboxane synthase activity. The TxS activity (TxA₂ synthase) was measured as described⁵⁹ using a method based on

Figure 6 Microparticle generation and ROS production from platelets activated by different stimulators. Human WP were stimulated with ABT-737 (1 μ M, 60 min), Thr/Cvx (5 mU/5 ng/ml, 2 min), and AA at the indicated concentrations (5 min) and processed for FACS analysis of MP formation, as assessed by the SSC/FSC plot (a and b) and (c) ROS production (DCF-DA fluorescence). The MP counts and DCF-DA fluorescence are presented as a fold increase compared with the control taken as 1. Data are presented as means $\pm$ S.E.M., $n = 4$; * $P < 0.05$ compared with the control. (a) Representative plots (SSC/FSC) of the original data

malondialdehyde formation by TxA_2 synthase. For the experiments, 0.2-ml aliquots of WP ($3 \times 10^9/\text{ml}$) were incubated with the reagents at 37°C with gentle shaking; the reaction was stopped by the addition of 0.115 ml of ice-cold trichloroacetic acid (20% wt/vol). The samples were kept on ice for 10 min and then centrifuged for 10 min at 14 000 r.p.m. Equal volumes of thiobarbituric acid (0.53% wt/vol) and supernatant were combined and incubated for 30 min at 70°C . The mixture was then incubated for 30 min at room temperature, and fluorescence was measured with a Wallac Victor2 1420 Multilabel counter (Perkin Elmer Wallac Life and Analytical Sciences, Boston, MA, USA) at excitation and emission wavelengths of 520 nm and 550 nm, respectively. Standards were prepared from the stable malondialdehyde derivative tetraethoxypropane with thiobarbituric acid in the presence of trichloroacetic acid.

Western blot analysis. For western blot analysis, WP were added directly to sodium dodecyl sulfate (SDS) gel-loading buffer and analyzed by SDS-polyacrylamide gel electrophoresis (PAGE), as described.⁵⁷ The separated proteins were transferred to a nitrocellulose membrane and incubated with the primary antibody overnight at 4°C . For visualization of the signal, goat anti-rabbit or anti-mouse IgG conjugated with horseradish peroxidase was used as a secondary antibody, followed by ECL detection (Amersham, Pharmacia Biotech).

Data analysis. All experiments were performed with at least $n=4$, and the combined data are expressed as means $\pm$ S.E.M. Differences between groups were analyzed by ANOVA, followed by Bonferroni's test; Student's t -test was used when appropriate. A value of $P<0.05$ was considered statistically significant.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. This study was supported by the DFG (SFB688, TP A2) and BMBF (01EO1003).

- Versteeg HH, Heemskerk JW, Levi M, Reitsma PH. New fundamentals in hemostasis. *Physiol Rev* 2013; **93**: 327–358.
- Semple JW, Italiano JE Jr, Freedman J. Platelets and the immune continuum. *Nat Rev Immunol* 2011; **11**: 264–274.
- Projahn D, Koenen RR. Platelets: key players in vascular inflammation. *J Leukoc Biol* 2012; **92**: 1167–1175.
- Geddis AE. The regulation of proplatelet production. *Haematologica* 2009; **94**: 756–759.
- Li J, Xia Y, Bertino AM, Coburn JP, Kuter DJ. The mechanism of apoptosis in human platelets during storage. *Transfusion* 2000; **40**: 1320–1329.
- Zhang H, Nimmer PM, Tahir SK, Chen J, Fryer RM, Hahn KR et al. Bcl-2 family proteins are essential for platelet survival. *Cell Death Differ* 2007; **14**: 943–951.
- Kodama T, Takehara T, Hikita H, Shimizu S, Shigekawa M, Li W et al. BH3-only activator proteins Bid and Bim are dispensable for Bak/Bax-dependent thrombocyte apoptosis induced by Bcl-xL deficiency: molecular requisites for the mitochondrial pathway to apoptosis in platelets. *J Biol Chem* 2011; **286**: 13905–13913.
- Burkhardt JM, Vaudel M, Gambaryan S, Radau S, Walter U, Martens L et al. The first comprehensive and quantitative analysis of human platelet protein composition allows the comparative analysis of structural and functional pathways. *Blood* 2012; **120**: e73–e82.
- Pereira J, Soto M, Palomo I, Ocqueteau M, Coetzee LM, Astudillo S et al. Platelet aging in vivo is associated with activation of apoptotic pathways: studies in a model of suppressed thrombopoiesis in dogs. *Thromb Haemost* 2002; **87**: 905–909.
- Rand ML, Wang H, Bang KW, Poon KS, Packham MA, Freedman J. Procoagulant surface exposure and apoptosis in rabbit platelets: association with shortened survival and steady-state senescence. *J Thromb Haemost* 2004; **2**: 651–659.
- Remenyi G, Szasz R, Friese P, Dale GL. Role of mitochondrial permeability transition pore in coated-platelet formation. *Arterioscler Thromb Vasc Biol* 2005; **25**: 467–471.
- Rukoyatkina N, Begonja AJ, Geiger J, Eigenthaler M, Walter U, Gambaryan S. Phosphatidylserine surface expression and integrin $\alpha\text{IIb}\beta_3$ activity on thrombin/convulxin stimulated platelets/particles of different sizes. *Br J Haematol* 2009; **144**: 591–602.
- Tonon G, Luo X, Greco NJ, Chen W, Shi Y, Jamieson GA. Weak platelet agonists and U46619 induce apoptosis-like events in platelets, in the absence of phosphatidylserine exposure. *Thromb Res* 2002; **107**: 345–350.
- Wolf BB, Goldstein JC, Stennicke HR, Beere H, Amarante-Mendes GP, Salvesen GS et al. Caspase functions in a caspase-independent manner to promote apoptosis-like events during platelet activation. *Blood* 1999; **94**: 1683–1692.
- Leytin V, Allen DJ, Mykhaylov S, Lyubimov E, Freedman J. Thrombin-triggered platelet apoptosis. *J Thromb Haemost* 2006; **4**: 2656–2663.
- Lin KH, Chang HC, Lu WJ, Jayakumar T, Chou HC, Fong TH et al. Comparison of the relative activities of inducing platelet apoptosis stimulated by various platelet-activating agents. *Platelets* 2009; **20**: 575–581.
- Winkler J, Rand ML, Schmugge M, Speer O. Omi/HtrA2 and XIAP are components of platelet apoptosis signalling. *Thromb Haemost* 2013; **109**: 532–539.
- Galluzzi L, Vitale I, Abrams JM, Alnemri ES, Baehrecke EH, Blagosklonny MV et al. Molecular definitions of cell death subroutines: recommendations of the Nomenclature Committee on Cell Death 2012. *Cell Death Differ* 2012; **19**: 107–120.
- Jackson SP, Schoenwaelder SM. Procoagulant platelets: are they necrotic? *Blood* 2010; **116**: 2011–2018.
- Leung R, Gwozdz AM, Wang H, Bang KW, Packham MA, Freedman J et al. Persistence of procoagulant surface expression on activated human platelets: involvement of apoptosis and aminophospholipid translocase activity. *J Thromb Haemost* 2007; **5**: 560–570.
- Lopez JJ, Salido GM, Gomez-Arteta E, Rosado JA, Pariente JA. Thrombin induces apoptotic events through the generation of reactive oxygen species in human platelets. *J Thromb Haemost* 2007; **5**: 1283–1291.
- Kramer RM, Roberts EF, Striffler BA, Johnstone EM. Thrombin induces activation of p38 MAP kinase in human platelets. *J Biol Chem* 1995; **270**: 27395–27398.
- Kramer RM, Roberts EF, Um SL, Borsch-Haubold AG, Watson SP, Fisher MJ et al. p38 mitogen-activated protein kinase phosphorylates cytosolic phospholipase A_2 (cPLA2) in thrombin-stimulated platelets. Evidence that proline-directed phosphorylation is not required for mobilization of arachidonic acid by cPLA2. *J Biol Chem* 1996; **271**: 27723–27729.
- Saklatvala J, Rawlinson L, Waller RJ, Sarsfield S, Lee JC, Morton LF et al. Role for p38 mitogen-activated protein kinase in platelet aggregation caused by collagen or a thromboxane analogue. *J Biol Chem* 1996; **271**: 6586–6589.
- Adam F, Kauskot A, Rosa JP, Bryckaert M. Mitogen-activated protein kinases in hemostasis and thrombosis. *J Thromb Haemost* 2008; **6**: 2007–2016.
- Mazharian A, Roger S, Maurice P, Berrou E, Popoff MR, Hoylaerts MF et al. Differential involvement of ERK2 and p38 in platelet adhesion to collagen. *J Biol Chem* 2005; **280**: 26002–26010.
- Canobbio I, Reineri S, Sinigaglia F, Balduini C, Torti M. A role for p38 MAP kinase in platelet activation by von Willebrand factor. *Thromb Haemost* 2004; **91**: 102–110.
- Begonja AJ, Geiger J, Rukoyatkina N, Rauchfuss S, Gambaryan S, Walter U. Thrombin stimulation of p38 MAP kinase in human platelets is mediated by ADP and thromboxane A_2 and inhibited by cGMP/cGMP-dependent protein kinase. *Blood* 2007; **109**: 616–618.
- Borsch-Haubold AG, Pasquet S, Watson SP. Direct inhibition of cyclooxygenase-1 and -2 by the kinase inhibitors SB 203580 and PD 98059. SB 203580 also inhibits thromboxane synthase. *J Biol Chem* 1998; **273**: 28766–28772.
- Kato H, Takai S, Matsushima-Nishiwaki R, Adachi S, Minamitani C, Otsuka T et al. HSP27 phosphorylation is correlated with ADP-induced platelet granule secretion. *Arch Biochem Biophys* 2008; **475**: 80–86.
- Kuliopulos A, Mohanlal R, Covic L. Effect of selective inhibition of the p38 MAP kinase pathway on platelet aggregation. *Thromb Haemost* 2004; **92**: 1387–1393.
- Lin LL, Wartmann M, Lin AY, Knopf JL, Seth A, Davis RJ. cPLA2 is phosphorylated and activated by MAP kinase. *Cell* 1993; **72**: 269–278.
- Borsch-Haubold AG, Kramer RM, Watson SP. Phosphorylation and activation of cytosolic phospholipase A_2 by 38-kDa mitogen-activated protein kinase in collagen-stimulated human platelets. *Eur J Biochem* 1997; **245**: 751–759.
- Waterman WH, Molski TF, Huang CK, Adams JL, Sha'afi RI. Tumour necrosis factor- α -induced phosphorylation and activation of cytosolic phospholipase A_2 are abrogated by an inhibitor of the p38 mitogen-activated protein kinase cascade in human neutrophils. *Biochem J* 1996; **319**, Pt 1: 17–20.
- Lin LL, Lin AY, Knopf JL. Cytosolic phospholipase A_2 is coupled to hormonally regulated release of arachidonic acid. *Proc Natl Acad Sci USA* 1992; **89**: 6147–6151.
- Shmelzer Z, Haddad N, Admon E, Pessach I, Leto TL, Eitan-Hazan Z et al. Unique targeting of cytosolic phospholipase A_2 to plasma membranes mediated by the NADPH oxidase in phagocytes. *J Cell Biol* 2003; **162**: 683–692.
- Kukreja RC, Kontos HA, Hess ML, Ellis EF. PGH synthase and lipoxygenase generate superoxide in the presence of NADH or NADPH. *Circ Res* 1986; **59**: 612–619.
- Edderkaoui M, Hong P, Vaquero EC, Lee JK, Fischer L, Friess H et al. Extracellular matrix stimulates reactive oxygen species production and increases pancreatic cancer cell survival through 5-lipoxygenase and NADPH oxidase. *Am J Physiol Gastrointest Liver Physiol* 2005; **289**: G1137–G1147.
- van der Wal DE, Gitz E, Du VX, Lo KS, Koekman CA, Versteeg S et al. Arachidonic acid depletion extends survival of cold-stored platelets by interfering with the (glycoprotein Ib α -14-3-3 ζ) association. *Haematologica* 2012; **97**: 1514–1522.
- De Genaro P, Simon MV, Rotstein NP, Politi LE. Retinoic acid promotes apoptosis and differentiation in photoreceptors by activating the P38 MAP kinase pathway. *Invest Ophthalmol Vis Sci* 2013; **54**: 3143–3156.
- Mayr M, Hu Y, Hainaut H, Xu Q. Mechanical stress-induced DNA damage and rap-p38MAPK signal pathways mediate p53-dependent apoptosis in vascular smooth muscle cells. *Faseb J* 2002; **16**: 1423–1425.
- Shou Y, Li L, Prabhakaran K, Borowitz JL, Isom GE. p38 mitogen-activated protein kinase regulates Bax translocation in cyanide-induced apoptosis. *Toxicol Sci* 2003; **75**: 99–107.
- Tripathi T, Smith AD, Abdi M, Alizadeh H. Acanthamoeba-cytopathic protein induces apoptosis and proinflammatory cytokines in human corneal epithelial cells by cPLA2 α activation. *Invest Ophthalmol Vis Sci* 2012; **53**: 7973–7982.

44. Penzo D, Tagliapietra C, Colonna R, Petronilli V, Bernardi P. Effects of fatty acids on mitochondria: implications for cell death. *Biochim Biophys Acta* 2002; **1555**: 160–165.
45. Scorrano L, Penzo D, Petronilli V, Pagano F, Bernardi P. Arachidonic acid causes cell death through the mitochondrial permeability transition. Implications for tumor necrosis factor- α apoptotic signaling. *J Biol Chem* 2001; **276**: 12035–12040.
46. Shiose A, Sumimoto H. Arachidonic acid and phosphorylation synergistically induce a conformational change of p47phox to activate the phagocyte NADPH oxidase. *J Biol Chem* 2000; **275**: 13793–13801.
47. Brill A, Chauhan AK, Canault M, Walsh MT, Bergmeier W, Wagner DD. Oxidative stress activates ADAM17/TACE and induces its target receptor shedding in platelets in a p38-dependent fashion. *Cardiovasc Res* 2009; **84**: 137–144.
48. Nardi MA, Gor Y, Feinmark SJ, Xu F, Karparkin S. Platelet particle formation by anti GPIIb/IIIa 49-66 Ab, Ca^{2+} ionophore A23187, and phorbol myristate acetate is induced by reactive oxygen species and inhibited by dexamethasone blockade of platelet phospholipase A_2 , 12-lipoxygenase, and NADPH oxidase. *Blood* 2007; **110**: 1989–1996.
49. Obata T, Brown GE, Yaffe MB. MAP kinase pathways activated by stress: the p38 MAPK pathway. *Crit Care Med* 2000; **28**: N67–N77.
50. Adam-Klages S, Schwandner R, Luschen S, Ussat S, Kreder D, Kronke M. Caspase-mediated inhibition of human cytosolic phospholipase A_2 during apoptosis. *J Immunol* 1998; **161**: 5687–5694.
51. Atsumi G, Tajima M, Hadano A, Nakatani Y, Murakami M, Kudo I. Fas-induced arachidonic acid release is mediated by Ca^{2+} -independent phospholipase A_2 but not cytosolic phospholipase A_2 , which undergoes proteolytic inactivation. *J Biol Chem* 1998; **273**: 13870–13877.
52. Kriem B, Sponne I, Ffire A, Malaplate-Armand C, Lozac'h-Pillot K, Koziel V *et al*. Cytosolic phospholipase A_2 mediates neuronal apoptosis induced by soluble oligomers of the amyloid-beta peptide. *FASEB J* 2005; **19**: 85–87.
53. Prinzen FW, Van der Vusse GJ, Arts T, Roemen TH, Coumans WA, Reneman RS. Accumulation of nonesterified fatty acids in ischemic canine myocardium. *Am J Physiol* 1984; **247**: H264–H272.
54. Westerberg E, Deshpande JK, Wieloch T. Regional differences in arachidonic acid release in rat hippocampal CA1 and CA3 regions during cerebral ischemia. *J Cereb Blood Flow Metab* 1987; **7**: 189–192.
55. Jiang F, Zhang Y, Dusting GJ. NADPH oxidase-mediated redox signaling: roles in cellular stress response, stress tolerance and tissue repair. *Pharmacol Rev* 2011; **63**: 218–242.
56. Lambeth JD. NOX enzymes and the biology of reactive oxygen. *Nat Rev Immunol* 2004; **4**: 181–189.
57. Gambaryan S, Geiger J, Schwarz UR, Butt E, Begonja A, Obergfell A *et al*. Potent inhibition of human platelets by cGMP analogs independent of cGMP-dependent protein kinase. *Blood* 2004; **103**: 2593–2600.
58. Begonja AJ, Gambaryan S, Geiger J, Aktas B, Pozgajova M, Nieswandt B *et al*. Platelet NAD(P)H-oxidase-generated ROS production regulates $\alpha\text{IIb}\beta_3$ -integrin activation independent of the NO/cGMP pathway. *Blood* 2005; **106**: 2757–2760.
59. Aktas B, Utz A, Hoenig-Liedl P, Walter U, Geiger J. Dipyridamole enhances NO/cGMP-mediated vasodilator-stimulated phosphoprotein phosphorylation and signaling in human platelets: *in vitro* and *in vivo/ex vivo* studies. *Stroke* 2003; **34**: 764–769.

Cell Death and Disease is an open-access journal published by Nature Publishing Group. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>