Correspondence

Male sterility in Mcl-1-flox mice is not due to enhanced Mcl1 protein stability

Cell Death and Disease (2016) 7, e2490; doi:10.1038/cddis.2016.391; published online 1 December 2016

Dear Editor,

The myeloid cell leukemia sequence 1 (*Mcl1*) gene is essential for development and tissue homeostasis throughout life. Homozygous deletion of *Mcl1* alleles leads to the death of the embryos at the time of implantation. We have previously engineered a conditional (*Mcl1*-flox) allele of the gene and reported that the insertion of a loxP site in the first exon had inadvertently led to the addition of 13 amino acids at the N-terminus of the coding sequence.¹ The Mcl1 protein is turned over rapidly, and several mechanisms have been implicated in the regulation of Mcl-1 levels.^{2–4} Interestingly, the extended Mcl1 protein (termed N⁺Mcl1) encoded by the *Mcl1*-flox allele appeared more stable than the wild-type protein. Significantly, homozygous *Mcl1*^{fl/fl} males were sterile, showing impaired sperm development and complete absence of mature sperm in the epididymis. This phenotype was indistinguishable from that observed in mice lacking both Bim and Blk.⁵ Since proper testis development and sperm maturation appear to require a fine balance between cell death and cell survival, we surmised that a more stable Mcl1 protein in *Mcl1*^{fl/fl} males somehow altered this balance and cause the testicular phenotype.

Since a more stable Mcl-1 protein may influence the outcome of some experiments in unpredictable ways, we have now used the CRISPR/Cas9 technology to remove the sequence encoding the 13 extra amino acids from the *Mcl1*-flox allele, giving rise to the *Mcl1*-flox-del strain. As shown in Figure 1a, and confirmed by sequencing of the locus, the Mcl1 protein produced from the *Mcl1*-flox-del allele had the same size as the WT protein, while the addition of 13 amino acids in the N⁺Mcl1 protein resulted in a higher molecular weight protein as observed by western blot. Even though the Mcl-1 protein in *Mcl1*-flox-del homozygous animals is normal, the males were still infertile. As observed in the *Mcl1*-flox strain, seminiferous tubules of homozygous *Mcl1*-flox-del males contained some round spermatids as well as spermatocytes, but lacked 1N spermatids and had essentially no mature sperm. Accordingly, their epididymides were devoid of sperm, like those of homozygous *Mcl1*-flox mice (Figure 1b). These results thus exclude that the enhanced stability of the N⁺Mcl1 protein is the cause of the testicular phenotype. Interestingly, an independent conditional Mcl1 KO strain has been generated, in which the first loxP site was introduced upstream of the first exon, and the coding sequence unmodified.⁶ Homozygous males of this strain were also infertile, confirming our present result.

Although our results fail to provide an explanation for the impairment of spermatogenesis in mice carrying a floxed *Mcl1* allele, it seems reasonable to assume that the introduction of a

Figure 1 Spermatogenesis defect in *Mcl1*-flox-del mice. (a) Detection of Mcl1 protein in thymic extracts from *Mcl1*-flox-del and *Mcl1*-flox mice. Note the size correction in *Mcl1*-flox-del mice compared to the protein from *Mcl1*-flox mice. (b) Sections through testis (i – iii) and epididymis (iv – vi) from WT, *Mcl1*-flox and *Mcl1*-flox-del homozygous mice. Note the complete absence of mature sperm in the epididymides of both floxed strains

loxP site close to *Mcl1* first exon may have unintended effects on the expression of the gene, at least in the testis. Surprisingly, several studies have shown that the floxed alleles do not provide any survival advantage to the many hematopoietic cells studied to date in *in vitro* death assays,^{6–8} suggesting that this effect may be specific to the testis. More work remains to be done to fully understand the intricacies governing *Mcl1* expression in different tissues.

Conflict of Interest

The authors declare no conflict of interest.

Casey Ah-Cann^{1,2}, Maximilien Tailler^{1,2},
 Andrew J Kueh^{1,2}, Marco J Herold^{1,2},
 Joseph T Opferman³, Marie-Liesse Asselin-Labat^{*,1,2}
 and Philippe Bouillet^{*,1,2}

¹ Molecular Genetics of Cancer, The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia;

² Department of Medical Biology, The University of Melbourne, Melbourne, Victoria 3052, Australia and

³ Department of Cell and Molecular Biology, St. Jude Children's Research Hospital, Memphis, TN, USA

* Corresponding authors: M-L Asselin-Labat or P Bouillet, Molecular Genetics of Cancer, The Walter and Eliza Hall Institute of Medical Research, 1G Royal Parade, Parkville, Victoria 3052, Australia, Tel: +61 3 9345 2334, Fax: +61 3 9347 0852, E-mail: labat@wehi.EDU.AU or bouillet@wehi.edu.au.

1. Okamoto T *et al. Proc Natl Acad Sci USA* 2014; **111**: 261–266.

2. Czabotar PE *et al. Proc Natl Acad Sci USA* 2007; **104**: 6217–6222.

3. Schwickart M *et al. Nature* 2010; **463**: 103–107.

4. Inuzuka H *et al. Nature* 2011; **471**: 104–109.

5. Coultas L *et al. EMBO J* 2005; **24**: 3963–3973.

6. Opferman JT *et al. Nature* 2003; **426**: 671–676.

7. Vikstrom I *et al. Science* 2010; **330**: 1095–1099.

8. Kelly GL *et al. Genes Dev* 2014; **28**: 58–70.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016