

Efficient genome editing in filamentous fungus *Trichoderma reesei* using the CRISPR/Cas9 system

Rui Liu*, Ling Chen*, Yanping Jiang, Zhihua Zhou, Gen Zou

CAS-Key Laboratory of Synthetic Biology, Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, China

Filamentous fungi have wide applications in biotechnology. The CRISPR/Cas9 system is a powerful genome-editing method that facilitates genetic alterations of genomes in a variety of organisms. However, a genome-editing approach has not been reported in filamentous fungi. Here, we demonstrated the establishment of a CRISPR/Cas9 system in the filamentous fungus *Trichoderma reesei* by specific codon optimization and *in vitro* RNA transcription. It was shown that the CRISPR/Cas9 system was controllable and conditional through inducible Cas9 expression. This system generated site-specific mutations in target genes through efficient homologous recombination, even using short homology arms. This system also provided an applicable and promising approach to targeting multiple genes simultaneously. Our results illustrate that the CRISPR/Cas9 system is a powerful genome-manipulating tool for *T. reesei* and most likely for other filamentous fungal species, which may accelerate studies on functional genomics and strain improvement in these filamentous fungi.

Keywords: genome-editing; CRISPR/Cas9; filamentous fungus; *Trichoderma reesei*

Cell Discovery (2015) 1, 15007; doi:10.1038/celldisc.2015.7; published online 12 May 2015

Introduction

Filamentous fungi, such as *Trichoderma reesei*, *Aspergillus niger*, *A. oryzae*, and *Penicillium chrysogenum*, have been used to produce diverse enzymes or metabolites for centuries [1]. To obtain hyper-producers, repeated classical mutagenesis and screening have been applied for long-term strain improvements. Genetic engineering approaches for filamentous fungi have been well developed. However, these approaches are not as efficient as those available for yeast and bacteria due to the additional complexity of filamentous fungi [2], such as multicellular morphology, cellular differentiation, thick chitinous cell walls, and the lack of suitable plasmids.

Genome-editing is a type of genetic engineering in which DNA is inserted, replaced, or removed from a genome using artificially engineered nucleases, or 'molecular scissors'. It is a useful tool to elucidate the function and effect of a gene or protein in a sequence-specific manner [3]. Recently, developed engineering nucleases, zinc-finger nucleases [4] and transcription activator-like effector nucleases [5], are believed to facilitate precise genome modifications. However, the time- and labor-intensive process of using zinc-finger nucleases and the requirements of specific enzyme engineering for different targets in transcription activator-like effector nucleases limit their applications. The type II clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated gene (Cas) system is the most popular genome-editing tool at this time. A single multidomain Cas9 catalyzes a double-strand break (DSB) in the target DNA composed of a 20-bp sequence matching the protospacer of the guide RNA (gRNA) and an adjacent downstream 5'-NGG nucleotide sequence (termed as the protospacer-adjacent motif (PAM)) [6]. Over the past 2 years, many studies have demonstrated that the

*These two authors contributed equally to this work.

Correspondence: Zhihua Zhou

Tel: +86 21 5492 4050; Fax: +86 21 5492 4049

E-mail: zhouzhihua@sippe.ac.cn

or Gen Zou

Tel: +86 21 5492 4049; Fax: +86 21 5492 4049

E-mail: zougen@sibs.ac.cn

Received 28 December 2014; accepted 28 February 2015

CRISPR/Cas9 system is a powerful genome-editing method that facilitates genetic alterations in genomes in a variety of organisms. Until now, there have been no reports on the CRISPR/Cas9 system or other genome-editing approaches in filamentous fungi, even in the model organism *Neurospora crassa*, despite the successful application of this technique in yeast [2, 7].

The filamentous fungus *T. reesei*, which is recognized by its Generally Recognized as Safe status by the US Food and Drug Administration, is the most widely used producer of commercial lignocellulolytic enzyme preparations [1] and is expected to be a potential cell factory for different heterologous proteins with its powerful ability to synthesize and secrete proteins with high quantities. However, the genetic basis of desired domestication traits is poorly understood, because robust genetic tools do not exist for this industrial host. The establishment of a genome-editing system can be used to develop *T. reesei* as a super cell factory for lignocellulolytic enzyme preparations and other heterologous proteins, as well as to characterize the regulatory mechanisms for induction, synthesis, and secretion of proteins in *T. reesei*.

In this study, we constructed a CRISPR/Cas9 system in *T. reesei* that allowed us to induce mutagenesis or introduce a new gene by homologous recombination into a target site of the *T. reesei* genome. In addition, it can be used to generate multiple genome modifications simultaneously by co-transforming gRNAs and donor DNAs (dDNAs) for different targets. This system

demonstrates great potential as a tool for genome engineering in filamentous fungi, owing to user-designated site specificity of Cas9 endonuclease activity and the simplicity of gRNA construction.

Results

Expression of codon-optimized Cas9 in *T. reesei*

Construction of the CRISPR/Cas9 system in an organism requires successful expression of Cas9. Unlike in *Saccharomyces cerevisiae* [7], the codon-optimized *Cas9* gene for human cells does not function in *T. reesei* Qm6a or Rut-C30 under control of the constitutive promoter Ppdc (the promoter of *pdc*, the gene encoding pyruvate decarboxylase, which participates in glucose metabolism [8]) or the inducible promoter Pcbh1 (the promoter of *cbh1*, the gene encoding cellobiohydrolase I, which is repressed by glucose but induced by a series of oligosaccharides or cellulose [9]). On the basis of the codon frequency of *T. reesei* (<http://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=51453>), we optimized the *Cas9* gene of *Streptococcus pyogenes* and the sequence of the SV40 nuclear localization signal (Supplementary Data S1). To detect the expression and localization of the optimized Cas9, the enhanced green fluorescent protein (eGFP) gene was fused to *toCas9* (optimized *Cas9* gene) and expressed in both the wild-type strain Qm6a and the mutant strain Rut-C30 under control of the Ppdc and Pcbh1 promoter, respectively (Supplementary




Figure 1 Expression and localization of Cas9-eGFP in *T. reesei* strains. (a) Detection of Cas9-eGFP expression in transformants 6a-pe and C30-ce by western blots. Transformants 6a-gfp and C30-gfp expressing eGFP were used as controls. Lanes C30-ce and 6a-pe were loaded with 0.05 mg protein. Lanes C30-gfp and 6a-gfp were loaded only with 0.01 mg protein to avoid overexposure. Anti-GFP antibody (Genscript, Nanjing, China) was used for detection. (b) Fluorescence microscopic assessment of the localization of Cas9-eGFP in C30-ce. The bars were 5 μ m. DAPI, 4',6-diamidino-2-phenylindole; eGFP, enhanced green fluorescent protein.

Data S1). The expression of *toCas9* in each of the selected transformants was detected based on western blots (Figure 1a). *toCas9* was expressed in both the transformant 6a-pe (introduced Ppdc-*toCas9-eGFP*-Tpdc into Qm6a) and the transformant C30-ce (introduced Pcbh1-*toCas9-eGFP*-Tpdc into Rut-C30).

To determine whether the SV40 nuclear localization signal localizes Cas9 to the nucleus, C30-ce was assessed using fluorescence microscopy (Figure 1b). The mycelia of C30-ce glowed with punctiform green fluorescence after 2 days of growth on the inducing culture media containing 1% Avicel. After staining with 4',6-diamidino-2-phenylindole ($1 \mu\text{g ml}^{-1}$), the majority of the green fluorescence spots overlapped with nuclei (blue fluorescence). This demonstrated that Cas9 was located mostly in the nucleus of *T. reesei*, where Cas9 is believed to generate DSBs within the target gene.

To build Cas9-expressing chassis, the Ppdc-*toCas9*-Tpdc cassette and Pcbh1-*toCas9*-Tpdc cassette were introduced into Qm6a and Rut-C30, respectively. To choose the Cas9 expression chassis for further investigations, we examined the growth and cellulase activities of the selected transformant 6a-pc from Qm6a and C30-cc from Rut-C30 after three consecutive subcultures for genetic stability, which did not show any differences compared with those of the parent strains (Supplementary Figure S1).

CRISPR/Cas9 system directed URA5 mutagenesis in T. reesei

Using appropriately designed gRNAs, the genome can be cut at any desired location by the Cas9 protein. Owing to the lack of confirmed RNA polymerase III-based promoters in *T. reesei*, such as the U6 promoter in *Arabidopsis thaliana* [10] and SNR52 promoter in *S. cerevisiae* [7], we preferred to introduce gRNAs into cells after transcription *in vitro* [11]. As *ura5* has undergone negative selection, its inactivation can be observed in 5-FOA-containing minimal medium (MM) plates (cells expressing wild-type *ura5* convert 5-FOA into the toxic substance 5' fluorouridine monophosphate, which severely limits cell growth). Thus, gRNA was designed to target *ura5* and was delivered into Cas9-expressing *T. reesei* cells through protoplast transformation. To ensure that Cas9 is active in cells, we first chose transformant 6a-pc (a transformant constitutively expressing Cas9) as the Cas9-expressing chassis to examine the mutation frequency. The *ura5* genes of 14 selected 5-FOA-resistant transformants were sequenced to confirm whether the mutations were directed by CRISPR/Cas9 (Table 1). Each *ura5* gene of

the 14 selected 6a-pc transformants showed a frameshift mutation at the expected target site, proximally located upstream of the PAM sequence (Figure 2a). This was consistent with the fact that CRISPR/Cas9-guided DSBs can be repaired through nonhomologous end joining (NHEJ), which generates insertions and deletions in the vicinity of the cleavage site. By contrast, no colony was observed in 5-FOA-containing MM plates from the control transformation of 6a-pc without gRNA. It confirmed that gRNA might be the essential component to result in mutations.

Controllability of the CRISPR/Cas9 system in T. reesei

In industries, the majority of filamentous fungi produce metabolites or enzymes at a specific stage or under unique conditions. A spatiotemporally controllable genome-editing tool precisely controls the metabolic shift in industry bioprocesses. We tested the conditionality of CRISPR/Cas9-guided mutagenesis in the Cas9-expressing chassis, C30-cc (a transformant expressing Cas9 only under inducing conditions) (Supplementary Data S1). As expected, mutagenesis was performed conditionally by an inducer (such as lactose or cellulose) in C30-cc. In the inducible medium (lactose or cellulose as the sole carbon resource), all 15 selected transformants (AR1–AR15) showed mutations in the target gene (Figure 2a). However, in the repressing medium (containing glucose, a repressor for promoter Pcbh1 that controls the *Cas9* gene expression), none of the colonies showed mutations. Besides, no colony was observed in 5-FOA-containing MM plates from the control transformation of 6a-pc without gRNA. The above results demonstrated that the CRISPR/Cas9 system functions in *T. reesei* as an efficient and controllable genome-editing tool in a sequence-specific manner. Besides, the expression of Cas9 under an inducible promoter or other conditionally activated promoters renders the CRISPR/Cas9 system a spatiotemporal-controller of genome-editing of filamentous fungi.

CRISPR/Cas9 system stimulated homologous recombination in T. reesei

Similarly to inducing mutagenesis in a sequence-specific manner, the CRISPR/Cas9 system also induces homologous recombination between endogenous and exogenous DNAs. We used *poua5* (an exogenous *ura5* gene, encoding URA5 in *P. oxalicum* [12]) as a selectable marker and the endogenous putative methyltransferase *lae1* gene as a target to test the incidence rate of homologous recombination (Figure 2b) by the CRISPR/Cas9 system [13]. The uridine-

Table 1 Primers and oligonucleotides used in this study

<i>Name</i>	<i>Sequence (5' to 3')</i>	<i>Experiment</i>
ToCas9F	AGCGCAGCTACAGCACAATCATGGACAAGAAGTACAGCAT	Cloning of <i>toCas9</i>
ToCas9R	TTAGACCTTGCCTTCTTCTTGGGGTCGCGCGGGAGTCG	Cloning of <i>toCas9</i>
PpdcF	ACGACGGCCAGTGCCAAGCTTAGGACTTCCAGGGCTACTTG	Cloning of <i>pdc</i> promoter
PpdcR	GATTGTGCTGTAGCTGCGCTGCTTTGATCGTTTTGAGGTGC	Cloning of <i>pdc</i> promoter
pcbhlcasF	ACGACGGCCAGTGCCAAGCTTTTTCCCTGATTCAGCGTACC	Cloning of <i>cbh1</i> promoter
pcbhlcasR	TTGACTATTGGGTTTCTGTGCCTC	Cloning of <i>cbh1</i> promoter
eGFPF	GTGGAGACAGCAGGGCTGACGTACCGGTCGCCACCATGGTG	Cloning of <i>eGFP</i>
eGFPR	TTACACCTTCTCTTCTTCTTGGGCTTGACAGCTCGTCCATG	Cloning of <i>eGFP</i>
TpdcF	AGAAGAAGAGGAAGGTGTGACCCGGCATGAAGTCTGACCG	Cloning of <i>pdc</i> terminator
TpdcR	TAATTGCGCGGATCCTCTAGATGGACGCCTCGATGTCTTCC	Cloning of <i>pdc</i> terminator
ToCGFPF	GCGGCGACTCCCGCGCCGACGTACCGGTCGCCACCATGGTG	Overlapping <i>eGFP</i> to <i>toCas9</i>
GFPTTrCasR	GTCGGCGCGGGAGTCGCCG	Overlapping <i>eGFP</i> to <i>toCas9</i>
lae1 up F	TGCAGGTCGACGATTGAAGCCAATCAGCAAGGTTGAACTG	Cloning the 5' flanking region of <i>lae1</i>
lae1 up R	TACTGGCTTAACTATGCGGCTTGGCATACTGAAAAATGT	Cloning the 5' flanking region of <i>lae1</i>
lae1-0.2 F	TGCAGGTCGACGATTATGTCTTTCACAAGTATTC	Cloning the 5' flanking region (0.2 kb) of <i>lae1</i>
lae1-0.4 F	TGCAGGTCGACGATTAACCGGTCAGCGAGCC	Cloning the 5' flanking region (0.4 kb) of <i>lae1</i>
lae1-0.6 F	TGCAGGTCGACGATTGCACTGGACTCTGGAAC	Cloning the 5' flanking region (0.6 kb) of <i>lae1</i>
lae1-0.8 F	TGCAGGTCGACGATTTCGTTATACTGTACTTAACAAG	Cloning the 5' flanking region (0.8 kb) of <i>lae1</i>
THRUra5 F	GCCGCATAGTTAAGCCAGTATACACTCCGCTATCGCTACG	Cloning the 5' flanking region of donor DNA
THRUra5 R	CAGGGCTGGTGACGGAATTTTCATAGTCAAGCTATCAGAG	Cloning the 5' flanking region of donor DNA
lae1 down F	AAATTCCGTCACCAGCCCTGCGGCAGCATAAGAGATGACAA	Cloning the 3' flanking region of <i>lae1</i>
lae1 down R	GGATCCTCTAGAGATGTGCTTCATTCGAATGAGACAGAGC	Cloning the 3' flanking region of <i>lae1</i>
lae1-0.2 R	GGATCCTCTAGAGATTGAAGAGCCGAATGTCTG	Cloning the 3' flanking region (0.2 kb) of <i>lae1</i>
lae1-0.4 R	GGATCCTCTAGAGATGGCTGAAGGCGAGGTTGAAC	Cloning the 3' flanking region (0.4 kb) of <i>lae1</i>
lae1-0.6 R	GGATCCTCTAGAGATAAATGTGTCTGCAAGCGTGATTAG	Cloning the 3' flanking region (0.6 kb) of <i>lae1</i>
lae1-0.8 R	GGATCCTCTAGAGATCCGCAGAACGCTGGCTCTC	Cloning the 3' flanking region (0.8 kb) of <i>lae1</i>
Ulae1 F	GGCAACCCACCCGTACCACA	Verifying the 5' flanking region of <i>lae1</i>
HR ura5 R	TGCTCGATGATCCCGACAGC	Cloning the 3' flanking region of <i>pura5</i> cassette
HR ura5 F	AATCGCCCTACTTCTTCACA	Cloning the 5' flanking region of <i>pura5</i> cassette
Dlae1 R	CAGGCTCTTATCGCCATCTC	Verifying the 3' flanking region of <i>lae1</i>
Tura5 F	GCGGCGTCCCAAGTTTGGC	Sequencing and verifying <i>ura5</i>
Tura5 R	CGGTAATCCTCCGTGTTCTT	Sequencing and verifying <i>ura5</i>




Figure 2 CRISPR/Cas9-based mutagenesis and homologous recombination in *T. reesei*. **(a)** Alignments of *ura5* mutants from 5-FOA-resistant colonies constitutive expression or post-lactose induction (the gene encoding Cas9 under control of Pcbh1 can be induced by oligosaccharides such as lactose, sophorose, or cellodextrin, in C30-cc) of Cas9 in gRNA transformants (14 6a-pc transformants and 15 C30-cc transformants). Tura5 is the wild-type reference *ura5* gene from *T. reesei* Qm6a and Rut-C30 (the PAM sequence is highlighted in blue and the guide RNA (gRNA)-guiding sequence in green). Deletions or insertions were found near the PAM sequence. **(b)** Schematic for homologous recombination (HR) of *lae1*, *vib1*, and *clr2* mediated by Cas9 and donor DNAs. Results of PCR analyses revealed HR events.

dependent transformant AR3 (derived from chassis C30-cc, Figure 2a) was used as a host. After co-transformation of gRNA (gRNA-*lae1*, targeting *lae1* site) with dDNA (dDNA-*lae1*, containing Pgpda-*poua5*-*Ttrpc* cassette and the 3' and 5' flanking regions of *lae1*) (Table 1), the transformants could be selected in MM plates. The sequence of the selectable marker cassette replaced *lae1* in all 14 transformants (Supplementary Figure S2). All Δ *lae1* transformants showed weakened sporulation (Figure 3a) and cellulytic capability (Figure 3b) compared with the controls, as expected [13]. Compared with other genetic approaches reportedly used for homologous recombination in filamentous fungi [14], the efficiency and precision of the CRISPR/Cas9 system supported its significant advantages.

The effect of the length of homology arms on the recombination frequency was also investigated. The uridine-dependent Cas9 expression chassis AR3 was transformed with gRNA-*lae1* and a set of dDNAs with various lengths of homology arms ranging from 0.2 to 1.0 kb at 0.2-kb intervals (dDNA-*lae1*-0.2k, dDNA-*lae1*-0.4k, dDNA-*lae1*-0.6k, dDNA-*lae1*-0.8k and dDNA-*lae1*) (Supplementary Data S1). Nine, 14 or 15 transformants (monoconidial cultures) from each transformation were selected and then checked using diagnostic PCR. These results are summarized in Table 2. The homologous recombination frequencies for all tested homology arms were $\geq 93\%$. These results demonstrated that a pair of 200-bp homology arms is sufficient to achieve efficient homologous integration stimulated by the CRISPR/Cas9 system in *T. reesei*.




Figure 3 Reduction of sporulation and cellulolytic capability in $\Delta lae1$ strains. **(a)** The observed sporulation of mutants and controls in potato dextrose agar plates after 7 days of growth. Sporulation was reduced in all three randomly selected $\Delta lae1$ strains compared with controls. **(b)** Assays for filter paper activity of $\Delta lae1$ strains and controls in inducing medium with 1% lactose after 3 days of fermentation. The cellulolytic capability was weakened significantly ($P < 0.01$, t -test). Error bars represent the s.d. among three independent measurements.

Table 2 Homologous recombination frequencies at the *lae1* locus using homology arms of various lengths

Homology arm size (kb)	No. of analyzed transformants	No. of deletants	Homologous recombination frequency (%)
0.2	15	14	93
0.4	15	14	93
0.6	15	15	100
0.8	9	9	100
1.0	14	14	100

Simultaneous homologous recombination of multigenes using the CRISPR/Cas9 system in T. reesei

Disruptions or homologous recombinations of multigenes are required during strain modification or to investigate regulatory systems for a unique metabolic process in filamentous fungi. A genetic approach that could target multiple genes simultaneously would greatly benefit such studies. In the uridine-dependent Cas9 expression chassis AR3, we tested the efficiency of multiple genome modifications directed by the CRISPR/Cas9 system. The selectable marker (Pgpda-poura5-Ttrpc cassette) was used to simultaneously target *lae1* and *vib1* (coding VIB1, a putative link between glucose signaling and carbon catabolite repression [15]). The gRNA and dDNA targeting *vib1* were termed gRNA-*vib1* and dDNA-*vib1*, respectively (containing Pgpda-poura5-Ttrpc cassette and the 3' and 5' flanking regions of *vib1*)

(Supplementary Data S1). After co-transformation of the two sets of gRNA and dDNA at the same molar concentration (gRNA-*vib1*:dDNA-*vib1*:gRNA-*lae1*:dDNA-*lae1* = 1:1:1:1), 24 transformants were selected in the uridine-free medium. All 18 positive transformants integrated the selectable marker cassette into the *lae1* locus. However, only three showed homologous recombination in both *lae1* and *vib1* loci. The frequency for double recombination was only 16% (Supplementary Figure S3).

As all the single-deletion transformants were $\Delta lae1$ strains, the molar concentration ratio of the gRNA and dDNA targeting *vib1* to those targeting *lae1* was adjusted to 1.5:1:1:1 (gRNA-*vib1*:dDNA-*vib1*:gRNA-*lae1*:dDNA-*lae1*) in the next co-transformation test. Eleven of the 24 randomly selected transformants were double-deletion strains. The frequency of double recombination was ~45%, and the remaining strains were all single-*lae1* deletion transformants (Supplementary Figure S3).

To test the efficiency of simultaneous homologous recombination of triple genes, *chr2* used as the third target, which is a conserved and essential transcription factor for cellulase gene expression in ascomycete fungi [16]. After co-transformation of the three sets of gRNA and dDNA at the same molar concentration (gRNA-*lae1*:dDNA-*lae1*:gRNA-*vib1*:dDNA-*vib1*:gRNA-*chr2*:dDNA-*chr2* = 1:1:1:1:1:1), 24 transformants were selected in uridine-free medium, in which there was one triple-gene deletion (4.2%), seven $\Delta lae1\Delta chr2$ double deletions, 13 $\Delta lae1$ single deletions, and two $\Delta chr2$ single deletions (Supplementary Figure

S4). Although the frequency of triple recombination was quite low, the CRISPR/Cas9 system is an applicable and promising approach to target multiple genes simultaneously.

Discussion

High-efficiency genetic manipulation approaches facilitate fungal strain improvement, as well as the elucidation of fungal molecular mechanisms behind the industrial applications. The first reports of genetic transformation in *Aspergillus* were in 1983 [17, 18], which lagged behind those for *Escherichia coli* and *S. cerevisiae*. Over the past three decades, scientists have attempted to develop various tools or strategies to improve genetic manipulation approaches in filamentous fungi [19–21]. Different genetic transformation approaches have been applied for homologous recombination in many filamentous fungi; however, the low homologous recombination frequency (< 5%) has limited studies on their functional genomics through targeted gene replacement [22]. Deletion of the Ku70 or Ku80 heterodimer (Ku complex) increased the frequency of homologous recombination by up to 65% in many filamentous fungi by eliminating the NHEJ pathway [23]. However, Δku strains were more sensitive to growth environments with specific chemicals such as phleomycin, bleomycin, and methyl/ethyl methane-sulfonate. Besides, the strategy to delete the Ku70 or Ku80 heterodimer is not suitable for *Agrobacterium*-mediated fungal transformation because of its incompatibility in NHEJ-deficient strains [24]. In this study, the CRISPR/Cas9 system showed the highest frequency of single homologous recombination in *T. reesei* (almost 100%) using a pair of ≥ 600 -bp homology arms (Table 2). Even when the length of the homology arms was decreased to 200 bp, the CRISPR/Cas9 system still reached a 93% homologous recombination frequency (Table 2), higher than the frequency in NHEJ-deficient *T. reesei* using 500-bp homology arms (63%) [25].

Establishing the CRISPR/Cas9 system in filamentous fungi is based on basic transformation systems. High frequencies of multicopy insertion and ectopic integration have been observed in the application of these filamentous fungal transformation systems [25, 26]. Compared with PEG/CaCl₂ protoplast, electroporation and biolistic transformation systems, *A. tumefaciens*-mediated transformation alleviates variable multicopy insertion and ectopic integration with its high efficiency (> 90%) of single-copy integration in *T. reesei* [27]. In this study, we introduced the Cas9-encoding gene into the genome of *T. reesei* using

random *A. tumefaciens*-mediated transformation. To ensure that the introduction and expression of the Cas9-encoding gene does not disturb the expression and function of other genes, the morphology and important phenotypes such as growth and cellulase activities of the selected transformants were compared with their parent strains (Supplementary Figure S1). Although the confirmation process for Cas9 expression typically takes more than 1 month, similarly to previous transformation systems, the Cas9 expression chassis (including the uridine-dependent Cas9 expression chassis) could be used in the following studies for different gene mutations or gene knockout /knock-in by changing the 20-bp protospacer of gRNA to the target locus according to the fungal genomic sequence information. Thus, the CRISPR/Cas9 system in filamentous fungi would save time and labor in the study of fungal life activity regulatory systems, such as reproduction, development, metabolic pathways, protein synthesis, and secretion, which involve multiple unknown functional genes. For example, it took more than a decade to construct genome-wide single-gene deletion libraries for functional genomics research in *N. crassa* and *A. nidulans* using traditional genetic manipulation approaches [28, 29]. The CRISPR/Cas9 system is expected to shorten the process for establishing a genome-wide single-gene deletion library for a filamentous fungal species with available genomic sequence information.

Owing to the site specificity of the Cas9 endonuclease and gRNA, the CRISPR/Cas9 system is believed to introduce a genetic alteration or a heterologous gene into the target locus precisely and efficiently. In this study, variations in the target *ura5* gene were detected in all selected mutagenesis transformants (Figure 2a), and *poura5* was shown to replace the target gene *lae1* in all homologous recombination transformants (Supplementary Figure S2). Even in the screened homologous recombination transformants simultaneously targeting triple loci, the heterologous gene *poura5* was shown to replace target genes *lae1*, *vib1*, and *clr2* or at least one or two of these genes (Supplementary Figure S4). Our results confirmed that the CRISPR/Cas9 system was successful in editing the genome of a filamentous fungus precisely and efficiently, similarly to other organisms.

In our simultaneous homologous recombination experiments targeting double genes, almost all of the selected single-deletion transformants were $\Delta lae1$ strains (Supplementary Figure S3). We did not obtain a $\Delta vib1$ strain, and DSBs were not observed in the gRNA-directed *vib1* locus. Similarly, in the

simultaneous homologous recombination experiments targeting triple genes, the selected single-deletion transformants were $\Delta lae1$ or $\Delta clr2$ strains, and the selected double deletion was $\Delta lae1/\Delta clr2$. No $\Delta vib1$, $\Delta lae1/\Delta vib1$, or $\Delta lae1/\Delta clr2$ strains were detected among the selected transformants (Supplementary Figure S4). The CRISPR/Cas9 system prefers to target the *lae1* or *clr2* locus rather than the *vib1* locus. The gene *vib1* may be located in a region of low transcriptional activity [30], which complicates the process of gRNA directing Cas9 to the *vib1* locus. The increase in the molar concentration ratio of gRNA-*vib1* (the gRNA targeting *vib1*) during co-transformation resulted in an increase in frequency for double recombination from 16 to 45% (Supplementary Figure S3). This suggests that the proportion of gRNA and dDNA of multiple genes requires optimization to achieve a high frequency of simultaneous multirecombination.

This study demonstrates precise editing of *T. reesei* genome through the repair of Cas9-induced DSBs in an NHEJ or homology-recombination method. However, unwanted insertion and deletion mutations were observed at off-target sites, which share high sequence similarity with the on-target site in other organisms [31–33]. We tried to avoid the selection of target sites sharing high sequence similarity with other sites in the *T. reesei* genome in this study. However, this is only one approach [34–36] used to reduce rather than eliminate off-target sites. Thus, it is possible that off-target might also occur in our *T. reesei* CRISPR/Cas9 system. It is an important issue to eliminate the off-target effect in the CRISPR/Cas9 system as well as the other genome-editing tools [37, 38]; specific experiments to test the off-target effect in *T. reesei* CRISPR/Cas9 system should be designed in a future study.

In this study, the primary applications of the CRISPR/Cas9 system have been demonstrated as model genome-engineering methods in filamentous fungi, which can accelerate research on the functional genomics of filamentous fungi. Numerous applications of the CRISPR/Cas9 system have been developed in other organisms within the past year. Similarly, more CRISPR/Cas9-mediated systems could be examined in filamentous fungi. For example, a CRISPR/Cas9-mediated large fragment deletion would facilitate genetic modification of filamentous fungal chassis [31]. A catalytically deactivated version of Cas9 can be used for genome-scale CRISPR-mediated control of gene repression and activation [10]. However, until now gRNAs could not be synthesized *in vivo* easily in *T. reesei*,

which may limit its application fields. It is important to allow gRNA to be synthesized *in vivo* by identifying effective RNA polymerase-III-based promoters in *T. reesei*.

Materials and Methods

Strains and plasmids

E. coli DH5 α served as the cloning host (Novagen, Gibbstown, NJ, USA). *A. tumefaciens* AGL1 was used to transform the *Cas9* gene to *T. reesei* strains [39]. The *T. reesei* strain Qm6a (13631) and Rut-C30 (56765) purchased from American Type Culture Collection (ATCC, Manassas, VA, USA), were used as the hosts for Cas9 expression. A T-DNA binary vector, pDHT/sk, containing the *hph* coding for hygromycin B phosphotransferase (under control of the *A. nidulans* *trpC* promoter and terminator) was used to construct the transformation vectors. pMD-18T vector (Takara, Dalian, China) was used to maintain gRNA templates and dDNAs in DH5 α .

Expression of *Cas9* in *T. reesei*

The *Cas9* gene with a SV40 nuclear localization signal was codon-optimized for expression in *T. reesei* (*toCas9*, Supplementary Data S1). To express *toCas9* efficiently, a constitutive promoter *Ppdc* and an inducible promoter *Pcbh1* were used to express *toCas9* in Qm6a and Rut-C30, respectively. The *eGFP* gene was fused to *toCas9* as a tag. All the cassettes with *Tpdc* as terminator fused into *XbaI* site of the linearized pDHT/sk using the ClonExpress MultiS One Step Cloning Kit (Vazyme, Nanjing, China) to generate four vectors, pDHT/sk-*Ppdc-toCas9-Tpdc* (pDHT/sk-PC), pDHT/sk-*Ppdc-toCas9-eGFP-Tpdc* (pDHT/sk-PE), pDHT/sk-*Pcbh1-toCas9-Tpdc* (pDHT/sk-CC), and pDHT/sk-*Pcbh1-toCas9-eGFP-Tpdc* (pDHT/sk-CE) (Table 1). We also constructed vectors only bearing the *eGFP* gene (pDHT/sk-*Ppdc-eGFP-Tpdc* and pDHT/sk-*Pcbh1-eGFP-Tpdc*) as controls. All of the expression vectors were transformed into the recipient *T. reesei* using *Agrobacterium*-mediated fungal transformation [40]. Transformants were selected using hygromycin B (10 $\mu\text{g ml}^{-1}$) and cefotaxime (300 μM) on potato dextrose agar. Corresponding to the expression vector, the transformants derived from Qm6a (with constitutive promoter) were named 6a-pc, 6a-pe, and 6a-gfp and ones from Rut-C30 (with inducible promoter) were named C30-cc, C30-ce, and C30-gfp. Each positive transformant was used to create monoconidial cultures for genetic stability test.

The positive transformants with eGFP tag under *Ppdc* and *Pcbh1* were grown on MM [40] with 2% glucose and 1% Avicel as carbon resources, respectively. The mycelium was collected by filtration and used to extract intracellular proteins after grinding in liquid nitrogen. Fusion protein production was examined by SDS-polyacrylamide gel electrophoresis and western blots using an anti-GFP antibody (Genscript, Nanjing, China).

Protoplast transformation of gRNA

The gRNA cassette including synthetic gRNA sequence and target DNA of the *ura5* gene (5'-GGCGAGGGCGGCAAC ATCGT cgg -3', PAM is shown in italics) was controlled by T7

promoter (Supplementary Data S1). This cassette was ligated into pMD-18T and maintained in DH5 α as template for the following transcription. Before transformation, the cassette was transcribed into RNA *in vitro* using the MEGAscript T7 Kit (Ambion, Austin, TX, USA). The generated RNA fragments (≥ 10 μ g) were transformed into the Cas9-expressing chassis (6a-pc and C30-cc, respectively) using a modified polyethylene glycol-mediated protoplast transformation procedure [41]. To express Cas9 as efficiently as possible, different carbon resources were used in screening MM plates corresponding to the promoters as mentioned in the previous section. Finally, the non-sense mutants of URA5 were selected in these modified MM plates including the 1.5 mg ml⁻¹ 5-FOA (Sangon, Shanghai, China) and 10 mM Uridine (Sangon). The genomic DNAs of selected transformants were extracted using the FastPrep instrument (MP Biomedicals, Santa Ana, CA, USA) with a standard protocol and used for the amplification of the gene *ura5*. Sequence analysis of the partial *ura5* from each transformant was carried out by BioSune (Shanghai, China).

Co-transformation of gRNA with donor DNA

The generated uridine-dependent Cas9 expression chassis (AR3) derived from C30-cc was used as a host to construct an efficient CRISPR-Cas-directed homologous recombination system. Similar to URA5 mutagenesis, the CRISPR-Cas target site (5'-GGATGCCGAAATCATGGCCG*ttg*-3', PAM is shown in italics) (Supplementary Data S1) was located in *lae1* [13]. The dDNA (dDNA-*lae1*) containing the 5' and 3' flanking sequences of *lae1* and the selectable marker cassette (the *ura5* gene from *P. oxalicum* controlled by the Pgpda promoter and Ttrpc terminator, Pgpda-*poura5*-Ttrpc) (Supplementary Data S1) was generated by overlapping PCR using Q5 (NEB, Beijing, China) and ligated into the pMD-18T vector. The generated vector was propagated in DH5 α and purified using the Plasmid Midi Kit (Qiagen, Hilden, Germany). Accompanying the RNA fragments including gRNA and the target sequence of *lae1*, dDNA was introduced into *T. reesei* using the protoplast transformation procedure. The transformants were selected using MM plates with 1% lactose as an inducer of Cas9.

Construction of dDNAs with different homology arm sizes

To examine the effect of homology arm length on targeted gene replacements in *T. reesei*, dDNAs were constructed using homology arms at various sizes. In addition to dDNA-*lae1*, Pgpda-*poura5*-Ttrpc was used as a selectable marker. Besides the 1-kb flanking sequences (dDNA-*lae1*), the 0.2-, 0.4-, 0.6-, and 0.8-kb flanking sequences of *lae1* were fused to the selectable marker. The generated dDNAs were named dDNA-*lae1*-0.2k, dDNA-*lae1*-0.4k, dDNA-*lae1*-0.6k, and dDNA-*lae1*-0.8k (Supplementary Data S1) and were ligated into the pMD-18T vector. The vector was propagated in DH5 α and purified using the Plasmid Midi Kit (Qiagen). The co-transformation procedure was similar to that described above.

Co-transformation of multiple fragments

Multiple genome modification was also performed in the uridine-dependent Cas9-expressing chassis AR3 through

co-transformation of two sets of gRNA and dDNA. In addition to the locus of *lae1*, another target site (5'-GGCTTCTGTCAGACAACCT*ttg*-3', PAM is shown in italics) (Supplementary Data S1) was located in the gene encoding VIB1, a putative link between glucose signaling and carbon catabolite repression [15]. The dDNA (dDNA-*vib1*) containing the 5' and 3' flanking sequences of *vib1* and the selectable marker cassette (Pgpda-*poura5*-Ttrpc) (Supplementary Data S1) was constructed as similarly to dDNA-*lae1*. In the triple-loci assay, *clr2* (5'-GGA CGAGGAACGAAGAGACC*ggg*-3', PAM is shown in italics) (Supplementary Data S1) was used as the third target, which was a conserved and essential transcription factor for cellulase gene expression in ascomycete fungi [16]. The dDNA (dDNA-*clr2*) containing the 5' and 3' flanking sequences of *clr2* and the selectable marker cassette (Pgpda-*poura5*-Ttrpc) (Supplementary Data S1) was constructed similarly to dDNA-*lae1*. After quantitating to the same molar concentration, two or three sets of gRNA and dDNA were introduced into *T. reesei* using the protoplast transformation procedure. The transformants were also selected using the MM plate with an inducer.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

This work was financially supported by National Basic Research Program of China (973: 2011CB707403), High-tech Research and Development Program of China (863: 2013AA102806) and the National Natural Science Foundation of China (31300073, 31470201).

References

- 1 Punt PJ, van Biezen N, Conesa A, Albers A, Mangnus J, van den Hondel C. Filamentous fungi as cell factories for heterologous protein production. *Trends Biotechnol* 2002; **20**: 200–206.
- 2 Jiang D, Zhu W, Wang Y, Sun C, Zhang KQ, Yang J. Molecular tools for functional genomics in filamentous fungi: recent advances and new strategies. *Biotechnol Adv* 2013; **31**: 1562–1574.
- 3 Burgess DJ. Technology: a CRISPR genome-editing tool. *Nat Rev Genet* 2013; **14**: 80.
- 4 Carroll D. Genome engineering with zinc-finger nucleases. *Genetics* 2011; **188**: 773–782.
- 5 Cermak T, Doyle EL, Christian M *et al.* Efficient design and assembly of custom TALEN and other TAL effector-based constructs for DNA targeting. *Nucleic Acids Res* 2011; **39**: e82.
- 6 Cong L, Ran FA, Cox D *et al.* Multiplex genome engineering using CRISPR/Cas systems. *Science* 2013; **339**: 819–823.
- 7 DiCarlo JE, Norville JE, Mali P, Rios X, Aach J, Church GM. Genome engineering in *Saccharomyces cerevisiae*

- using CRISPR-Cas systems. *Nucleic Acids Res* 2013; **41**: 4336–4343.
- 8 Li J, Wang J, Wang S, Xing M, Yu S, Liu G. Achieving efficient protein expression in *Trichoderma reesei* by using strong constitutive promoters. *Microb Cell Fact* 2012; **11**: 84.
 - 9 Ilmén M, Onnela ML, Klemsdal S, Keränen S, Penttilä M. Functional analysis of the cellobiohydrolase I promoter of the filamentous fungus *Trichoderma reesei*. *Mol Gen Genet* 1996; **253**: 303–314.
 - 10 Gilbert LA, Larson MH, Morsut L *et al*. CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* 2013; **154**: 442–451.
 - 11 Wang Y, Li Z, Xu J *et al*. The CRISPR/Cas system mediates efficient genome engineering in *Bombyx mori*. *Cell Res* 2013; **23**: 1414–1416.
 - 12 Liu G, Zhang L, Wei X *et al*. Genomic and secretomic analyses reveal unique features of the lignocellulolytic enzyme system of *Penicillium decumbens*. *PLoS One* 2013; **8**: e55185.
 - 13 Seiboth B, Karimi RA, Phatale PA *et al*. The putative protein methyltransferase LAE1 controls cellulase gene expression in *Trichoderma reesei*. *Mol Microbiol* 2012; **84**: 1150–1164.
 - 14 Meyer V, Arentshorst M, El-Ghezal A *et al*. Highly efficient gene targeting in the *Aspergillus niger kusA* mutant. *J Biotechnol* 2007; **128**: 770–775.
 - 15 Xiong Y, Sun J, Glass NL. VIB1, a link between glucose signaling and carbon catabolite repression, is essential for plant cell wall degradation by *Neurospora crassa*. *PLoS Genet* 2014; **10**: e1004500.
 - 16 Coradetti ST, Craig JP, Xiong Y, Shock T, Tian CG, Glass NL. Conserved and essential transcription factors for cellulase gene expression in ascomycete fungi. *Proc Natl Acad Sci USA* 2012; **109**: 7397–7402.
 - 17 Ballance DJ, Buxton FP, Turner G. Transformation of *Aspergillus nidulans* by the orotidine-5'-phosphate decarboxylase gene of *Neurospora crassa*. *Biochem Biophys Res Commun* 1983; **112**: 284–289.
 - 18 Tilburn J, Scazzocchio C, Taylor GG, Zabicky-Zissman JH, Lockington RA, Davies RW. Transformation by integration in *Aspergillus nidulans*. *Gene* 1983; **26**: 205–221.
 - 19 Penttilä M, Nevalainen H, Rättö M, Salminen E, Knowles J. A versatile transformation system for the cellulolytic filamentous fungus *Trichoderma reesei*. *Gene* 1987; **61**: 155–164.
 - 20 Herzog RW, Daniell H, Singh NK, Lemke PA. A comparative study on the transformation of *Aspergillus nidulans* by microprojectile bombardment of conidia and a more conventional procedure using protoplasts treated with polyethyleneglycol. *Appl Microbiol Biot* 1996; **45**: 333–337.
 - 21 Frandsen RJN. A guide to binary vectors and strategies for targeted genome modification in fungi using *Agrobacterium tumefaciens*-mediated transformation. *J Microbiol Methods* 2011; **87**: 247–262.
 - 22 Kanaar R, Hoeijmakers JH, van Gent DC. Molecular mechanisms of DNA double strand break repair. *Trends Cell Biol* 1998; **8**: 483–489.
 - 23 Weld RJ, Plummer KM, Carpenter MA, Ridgway HJ. Approaches to functional genomics in filamentous fungi. *Cell Res* 2006; **16**: 31–44.
 - 24 van Attikum H, Bundock P, Hooykaas PJ. Non-homologous end-joining proteins are required for *Agrobacterium* T-DNA integration. *EMBO J* 2001; **20**: 6550–6558.
 - 25 Zhang G, Hartl L, Schuster A *et al*. Gene targeting in a nonhomologous end joining deficient *Hypocrea jecorina*. *J Biotechnol* 2009; **139**: 146–151.
 - 26 Krappmann S, Sasse C, Braus GH. Gene targeting in *Aspergillus fumigatus* by homologous recombination is facilitated in a nonhomologous end-joining-deficient genetic background. *Eukaryot Cell* 2006; **5**: 212–215.
 - 27 Fu K, Liu L, Fan L, Liu T, Chen J. Accumulation of copper in *Trichoderma reesei* transformants, constructed with the modified *Agrobacterium tumefaciens*-mediated transformation technique. *Biotechnol Lett* 2010; **32**: 1815–1820.
 - 28 Colot HV, Park G, Turner GE *et al*. A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors. *Proc Natl Acad Sci USA* 2006; **103**: 10352–10357.
 - 29 Park G, Colot HV, Collopy PD *et al*. High-throughput production of gene replacement mutants in *Neurospora crassa*. *Methods Mol Biol* 2011; **722**: 179–189.
 - 30 Bi X. Heterochromatin structure: Lessons from the budding yeast. *IUBMB Life* 2014; **66**: 657–666.
 - 31 Essletzbichler P, Konopka T, Santoro F *et al*. Megabase-scale deletion using CRISPR/Cas9 to generate a fully haploid human cell line. *Genome Res* 2014; **24**: 2059–2065.
 - 32 Hruscha A, Krawitz P, Rechenberg A *et al*. Efficient CRISPR/Cas9 genome editing with low off-target effects in zebrafish. *Development* 2013; **140**: 4982–4987.
 - 33 Veres A, Gosis BS, Ding QR *et al*. Low incidence of off-target mutations in individual CRISPR-Cas9 and TALEN targeted human stem cell clones detected by whole-genome sequencing. *Cell Stem Cell* 2014; **15**: 254–254.
 - 34 Ran FA, Hsu PD, Lin CY *et al*. Double nicking by RNA-guided CRISPR Cas9 for enhanced genome editing specificity. *Cell* 2013; **155**: 479–480.
 - 35 Ansai S, Kinoshita M. Targeted mutagenesis using CRISPR/Cas system in medaka. *Biol Open* 2014; **3**: 362–371.
 - 36 Fu Y, Sander JD, Reyon D, Cascio VM, Joung JK. Improving CRISPR-Cas nuclease specificity using truncated guide RNAs. *Nat Biotechnol* 2014; **32**: 279–284.
 - 37 Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem* 1959; **31**: 426–428.
 - 38 Grau J, Boch J, Posch S. TALENoffer: genome-wide TALEN off-target prediction. *Bioinformatics* 2013; **29**: 2931–2932.
 - 39 Ram AFJ, Michielse CB, Hooykaas PJJ, van den Hondel CAMJJ. *Agrobacterium*-mediated transformation of the filamentous fungus *Aspergillus awamori*. *Nat Protoc* 2008; **3**: 1671–1678.
 - 40 Zou G, Shi S, Jiang Y *et al*. Construction of a cellulase hyper-expression system in *Trichoderma reesei* by

promoter and enzyme engineering. *Microb Cell Fact* 2012; **11**: 21.

- 41 Liu T, Wang T, Li X, Liu X. Improved heterologous gene expression in *Trichoderma reesei* by cellobiohydrolase I gene (*cbh1*) promoter optimization. *Acta Bioch Bioph Sin* 2008; **40**: 158–165.

(**Supplementary Information** is linked to the online version of the paper on the *Cell Discovery* website.)



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>