OCEAN SCIENCE

Shifting sink

DEMIS

Global Biogeochemical Cycles **22**, BG4027 (2008)

Over the past decade, the North Atlantic Ocean has been absorbing less and less atmospheric carbon dioxide. Now scientists report that blame may lie with changes in the region's dominant climate pattern, known as the North Atlantic Oscillation, rather than anthropogenic warming.

The time series of carbon dioxide observations in the North Atlantic extends back only as far as 1995. To reconstruct the patterns of carbon dioxide uptake

back through 1979, Helmuth Thomas at Dalhousie University in Halifax, Nova Scotia, and colleagues created a 'hindcast', essentially the opposite of a model forecast, for the region using a complex model of North Atlantic Ocean chemistry and physics. Close correspondence of their model simulations with the few available ten-year observational records gave the researchers increased confidence in the results.

They found that the decrease in CO₂ absorption reported since the mid 1990s coincided with a shift in the North Atlantic Oscillation index, a statistic that describes the location of air pressure anomalies, to more negative values, which altered patterns of circulation within the North Atlantic basin. They suggest that a return to a more positive index in the future could spur increased carbon dioxide uptake in the eastern North Atlantic.

Alicia Newton

JOY SINGARAYER

a 'bio-engineering' approach involving the careful selection of crop varieties could be used to mitigate surface warming. The team used a global climate model to simulate the warming that would result from doubling atmospheric carbon dioxide concentrations and heightening albedo, or solar reflectivity, of vegetation over specific cropland areas. Their results show that increasing the albedo of vegetation in these areas by 20 per cent could result in regional summertime cooling of up to 1 °C across much of mid-latitude Eurasia and North America. This would cancel out up to one-fifth of the regional seasonal warming caused by a doubling of atmospheric carbon dioxide.

The authors say that, unlike growing biofuels, choosing the more reflective crop varieties would not disrupt global food production. They note that further temperature reductions could be achieved by genetic modification of plants for high albedo.

Olive Heffernan

EXTREME EVENTS

The hydrate hazard

J. Geophys. Res. **113**, C12023 (2008)

A moderate increase in sea-floor temperature could trigger the widespread release of methane from ocean hydrates, finds new research. Large quantities of the potent greenhouse gas are stored beneath the sea in solid crystalline structures, known as hydrates, that could potentially be destabilized by ocean warming.

Matthew Reagan and George Moridis of the Lawrence Berkeley National Laboratory in Berkeley, California, used computer simulations to examine the effect of 100 years of moderate ocean warming on the stability of deep and shallow ocean hydrates. They

found that deep ocean deposits remained stable over the 100-year period, but that shallow ocean deposits were highly unstable, releasing significant quantities of dissolved and gaseous methane in response to just 1 °C of warming at the sea floor. Of the shallow deposits examined, cold hydrates, representative of the Arctic continental shelf, released nearly three times more methane than those representative of warm regions such as the Gulf of Mexico.

An increase in deep-sea pressure resulting from sea level rise would delay methane release but would not prevent its inevitable escape into the sea, say the researchers. Given that the release of methane could amplify climate change, the authors call for a detailed assessment of the hydrate hazard.

Anna Armstrong

US GEOLOGICAL SURVEY

MITIGATION

Cooling crops

Curr. Biol. doi:10.1016/j.cub.2008.12.025 (2009)

Cultivating only crop varieties that have a high capacity for reflecting sunlight could lower regional temperatures by as much as 1 °C during summertime, shows new research.

A team from the University of Bristol, UK, led by Andy Ridgwell tested whether

BIODIVERSITY AND ECOLOGY

Factoring in fish

PETER MUMBY/EXETER UNIVERSITY

Science **323**, 359–362 (2009)

Fish may contribute to the ocean carbon cycle in a previously unsuspected way. Traditionally, scientists have attributed the production of calcium carbonate in surface waters to marine plankton. But a study led by Rod Wilson at the University of Exeter, UK,

now shows that fish excrete the mineral as a by-product of drinking seawater.

Wilson and his colleagues measured the rate at which fish excrete calcium carbonate from their intestines and then combined this with two independent estimates of global fish biomass. They calculated that bony fish contribute 3 to 15 per cent of the total estimated carbonate in the ocean. Their study helps to unravel a mystery that has perplexed oceanographers for decades — namely, why the upper 1,000 metres of ocean is more alkaline than predicted by computer models. The carbonate produced by fish dissolves more easily than other forms, say the authors, and may explain this puzzling distribution.

The authors expect that fish will have an even greater role in the oceanic carbon cycle in the future. But calculating their exact contribution will be complex: rising ocean temperatures will increase the metabolic rate of fish — and their excretion of carbonate — but continued overfishing would decrease their overall carbonate output.

Olive Heffernan

REGIONAL CLIMATE

Fading forests

NATE STEPHENSON / AAAS

Science **323**, 521–524 (2009)

Regional warming may be killing off trees in old forests across the western United States, finds new research. Until now, large-scale studies of tree mortality have been confined to the tropics, where new tree growth has counterbalanced a recent surge in tree mortality.

Phillip van Mantgem and Nathan Stephenson of the US Geological Survey and colleagues analysed mortality data from 76 undisturbed old-growth coniferous forest plots censused between 1955 and 2007 in the western United States. During this period, tree mortality increased in 87 per cent of the plots examined and doubled within 17 to 29 years, depending on region. Forest die-back was not confined to any one location

or tree type. Rather, mortality increased at low and high elevations, in young and old trees, and in all major genera. The density of forests decreased over the study period, suggesting that competition for resources was not the main cause of increasing mortality. Rather, regional warming — which approached 0.5 °C per decade over the study period — and an associated rise in drought stress may have been responsible.

The researchers warn that the steep rise in tree mortality may be symptomatic of increasing stress on forests and could serve as a taste of things to come.

Anna Armstrong

ANTHROPOGENIC CHANGE

Early seasons

PA-PHOTOS / ISTOCKPHOTO.COM

Nature **457**, 435–441 (2009)

Spring now arrives sooner than it did 50 years ago, say scientists, who suggest that this shift is linked to an increase in atmospheric greenhouse gas concentrations.

Alexander Stine of the University of California at Berkeley and colleagues reconstructed seasonal cycles in the world's temperate zones in the last century, using updated records from the Climate Research Unit at the University of East Anglia, UK. Over land, they found that the onsets of all four seasons have shifted forward by an average of about 1.7 days between 1954 and 2007. During this period, the temperature difference between summer and winter narrowed by about six per cent over land because of warming winters. Both these changes are significantly larger than those observed in the first half of this century from 1900–1953, leading the researchers to infer that human influence, and not natural variability alone, caused the changes.

Warmer winters combined with a premature turn towards spring may be causing earlier spring blooms and bird migrations, effects long reported by biologists. The group's results suggest that these plants and animals are indeed directly responding to a changing climate.

Alicia Newton

authors & referees @ npg

All the information you need for publishing your research with Nature Publishing Group

www.nature.com/authors

- **Editorial policies** – publication policies for authors and peer-reviewers, to help you find an appropriate journal for publication of your work
- **Author services** – including author and peer-reviewer guidelines and submission pages for our journals
- **Peer reviewers** – peer review policies, advice for peer-reviewers
- **Author benefits** – the many benefits of publishing in an NPG journal

Join the discussion:

Join the Nautilus blog for present and future NPG authors

Join the Peer-to-Peer blog discussing the peer review system

www.nature.com/authors

nature publishing group