npg

Argonaute-2 regulates the proliferation of adult stem cells in planarian

Yong-Qin Li^{1, 3}, An Zeng^{1, 3}, Xiao-Shuai Han¹, Chen Wang¹, Ge Li^{1, 4}, Zhen-Chao Zhang¹, Jian-Yong Wang¹, Yong-Wen Qin², Qing Jing^{1, 2}

¹Key Laboratory of Stem Cell Biology and Laboratory of Nucleic Acid and Molecular Medicine, Institute of Health Sciences, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences and Shanghai Jiao-Tong University School of Medicine, Shanghai 200025, China; ²Department of Cardiology, Changhai Hospital, Shanghai 200433, China; ³Graduate School of Chinese Academy of Sciences, Beijing 100049, China; ⁴Present address: Department of Pediatrics, University of Alberta, Edmonton, AB, Canada T6G 2S2

Cell Research (2012) 22:275. doi:10.1038/cr.2012.8; published online 5 January 2012

Correction to: Cell Research (2011) 21:1750-1754. doi:10.1038/cr.2011.151; published online 6 September 2011

In our discussion of the role of Argonaute proteins in stem cell regulation, we inadvertently omitted an important earlier contribution by Agata's group [1]. Through testing conserved post-transcriptional regulators in the planarian *Dugesia japonica*, their study reports that *Djago2* is required to maintain stem cell identity.

We regret any inconvenience caused by this omission.

Reference

 Rouhana L, Shibata N, Nishimura O, Agata K. Different requirements for conserved post-transcriptional regulators in planarian regeneration and stem cell maintenance. *Dev Biol* 2010; **341**:429-443.