PKR activation and eIF2 α phosphorylation mediate human *globin* mRNA splicing at spliceosome assembly

Lena Ilan^{1,*}, Farhat Osman^{1,*}, Lise Sarah Namer^{1,*}, Einav Eliahu¹, Smadar Cohen-Chalamish¹, Yitzhak Ben-Asouli¹, Yona Banai¹, Raymond Kaempfer¹

¹Department of Biochemistry and Molecular Biology, Institute of Medical Research Israel-Canada, The Hebrew University-Hadassah Medical School, Jerusalem, Israel

Short elements in mammalian mRNA can control gene expression by activating the RNA-dependent protein kinase PKR that attenuates translation by phosphorylating cytoplasmic eukaryotic initiation factor 2 α (eIF2 α). We demonstrate a novel, positive role for PKR activation and eIF2 α phosphorylation in human *globin* mRNA splicing. PKR localizes in splicing complexes and associates with splicing factor SC35. Splicing and early-stage spliceosome assembly on β -*globin* pre-mRNA depend strictly on activation of PKR by a codon-containing RNA fragment within exon 1 and on phosphorylation of nuclear eIF2 α on Serine 51. Nonphosphorylatable mutant eIF2 α S51A blocked β -*globin* mRNA splicing in cells and nuclear extract. Mutations of the β -*globin* RNA activator abrogated PKR activation and profoundly affected mRNA splicing efficiency. PKR depletion abrogated splicing and spliceosome assembly; recombinant PKR effectively restored splicing. Excision of the first intron of β -*globin* induces strand displacement within the RNA activator of PKR by a sequence from exon 2, a structural rearrangement that silences the ability of spliced β -*globin* mRNA to activate PKR. Thus, the ability to activate PKR is transient, serving solely to enable splicing. α -*Globin* pre-mRNA splicing is controlled likewise but positions of PKR activator and silencer are reversed, demonstrating evolutionary flexibility in how PKR activation regulates *globin* mRNA splicing through eIF2 α phosphorylation.

Keywords: *globin* mRNA splicing; eIF2 α phosphorylation; PKR; RNA activator of PKR; spliceosome assembly

Cell Research (2017) 27:688–704. doi:10.1038/cr.2017.39; published online 4 April 2017

Introduction

The RNA-dependent protein kinase R (PKR) is an intracellular sensor of stress that plays a central role in antiviral defense and innate immunity [1]. Double-stranded RNA generated during viral replication activates PKR, which inhibits protein synthesis by phosphorylating the α -subunit of eukaryotic translation initiation factor 2 (eIF2 α) on Ser51; phosphorylation of eIF2 α blocks GTP/GDP exchange needed for recycling of eIF2 between rounds of translation initiation [2]. eIF2 α phosphorylation is critical for mounting the integrated cellular stress response [3–5]. Activation of PKR in the cytoplasm leads

to eIF2 α phosphorylation that results in a block of translation of mRNAs such as *globin* mRNA [6, 7].

Effective engagement of the tandem RNA-binding domains of PKR needed for *trans*-autophosphorylation of the PKR dimer and the subsequent kinase activation [8, 9] requires double-helical RNA of minimally 16–18 base pairs (bp) [10] and optimally 40–80 bp [11]. However, short elements within mammalian mRNA can regulate specific gene expression by activating PKR, as shown for the human genes encoding inflammatory cytokines, such as tumor necrosis factor- α (TNF- α) [12] and interferon- γ (IFN- γ) [13]. IFN- γ mRNA activates PKR through a 5'-proximal pseudoknot, resulting in eIF2 α phosphorylation and attenuation of its own translation through a negative feedback loop [13, 14]. Efficient splicing of TNF- α pre-mRNA is dependent on a short element in the 3'-untranslated region (3'-UTR) that potently activates PKR [12]. This *cis*-acting RNA element, conserved from teleost fish to humans, folds into a pseudoknot that is critical for PKR activation and mRNA splicing. Splicing enabled

*These three authors contributed equally to this work.

Correspondence: Raymond Kaempfer

Tel: +972 2 6758389

E-mail: kaempfer@hebrew.edu

Received 20 July 2016; revised 20 November 2016; accepted 30 December 2016; published online 4 April 2017

by the *TNF- α* RNA activator of PKR depends strictly on phosphorylation of eIF2 α on Ser51 that yet does not repress translation. Indeed, eIF2 α phosphorylation is not only necessary but also sufficient to achieve efficient splicing of *TNF- α* mRNA (manuscript in submission). eIF2 α phosphorylation positively controls *TNF- α* mRNA splicing in primary human immune cells, showing its physiological relevance, but the mechanism behind this need remains to be resolved.

Hemoglobin production depends on a massive expression of *globin* mRNA [15] that demands highly efficient splicing of precursor transcripts. We report here that splicing of α - and β -*globin* pre-mRNAs depends strictly on the activation of PKR by intragenic RNA activators that overlap with translated codons. Splicing of β -*globin* mRNA is abrogated by PKR depletion, which can be restored by adding back recombinant PKR. The RNA activator of PKR in β -*globin* pre-mRNA resides in the first exon; mutations of a short helix within this activator profoundly affected both PKR activation and mRNA splicing. Efficient *globin* mRNA splicing requires phosphorylation of eIF2 α . PKR activation and eIF2 α phosphorylation mediate splicing at an early step in β -*globin* spliceosome assembly.

The activator of PKR in β -*globin* pre-mRNA is generated from 5'-UTR sequence and the start of the open reading frame (ORF), as in the case of *IFN- γ* mRNA [13, 14]. However, we show that excision of the first intron of β -*globin* induces strand displacement within exon 1 by a sequence from exon 2 that affects base pairing in the short helix critical for PKR activation, resulting in silencing of the PKR activator within the spliced mRNA. In this manner, the ability to activate PKR is transient, solely to enable splicing.

Our results link a mechanism that controls mRNA translation in the cytoplasm, i.e., phosphorylation of the eIF2 α chain upon RNA-mediated activation of PKR, to a critical step in *globin* mRNA splicing in the nucleus. The finding that splicing of the *globin* mRNAs is mediated by intragenic RNA activators of PKR shows that this stress response kinase is used more widely beyond the immune system to effect post-transcriptional upregulation of specific genes.

Results

PKR inhibitors block splicing of β -globin mRNA

We followed excision of human β -*globin* introns 1 (Figure 1A) and 2 (Figure 1B) in transfected cells by means of antisense RNA probes. Addition of 2-aminopurine (2-AP), an inhibitor of eIF2 α kinases [16] that blocks splicing of *TNF- α* mRNA [12], induced accumulation of

unspliced β -*globin* pre-mRNA at the expense of spliced mRNA, leading to an increase in pre-mRNA/mRNA ratio (Figure 1A-1B). When we expressed a β -*globin* gene from which intron 2 had been deleted, intron 1 failed to be excised and only unspliced pre-mRNA was recovered (Figure 1C), in agreement with earlier observations [17]. This result implies that intron 1 must be excised before intron 2 deletion, rendering removal of intron 1 the critical first step in β -*globin* mRNA splicing. Once intron 1 has been excised, splicing is completed efficiently (Figure 1B, lanes without 2-AP).

We next created constructs containing β -*globin* exons 1 and 2 wherein the β -*globin* coding sequence is fused in-frame to green fluorescent protein (GFP) ORF (Figure 1D). When splicing is blocked, the construct containing intron 1 would generate an RNA product having a stop codon in the intron, precluding expression of GFP. Cells transfected with the intronless construct expressed GFP regardless of the addition of 2-AP or co-expression of PKR Δ 6, a dominant-negative mutant of human PKR (Figure 1E). By contrast, these treatments abolished GFP expression when intron 1 was present (Figure 1E). By forming heterodimers with wild-type (*wt*) PKR, kinase-dead PKR Δ 6 blocks its activation [18]. Therefore, 2-AP and PKR Δ 6 blocked excision of the β -*globin* intron in *globin-GFP* pre-mRNA. Unlike the case of β -*globin* pre-mRNA (Figure 1C), *globin-GFP* pre-mRNA splicing was not impeded by lack of the second intron; apparently, due to the lack of exon 3, the downstream sequence does not interfere with upstream splicing. Moreover, in a HeLa cell nuclear extract, splicing of a β -*globin* pre-mRNA template consisting of exon 1, intron 1 and exon 2 but lacking downstream sequence [19] was inhibited completely by 2-AP (Figure 1F). The effect of 2-AP resembled that of okadaic acid, a known splicing inhibitor [20] (Figure 1G). These results suggest that splicing of β -*globin* pre-mRNA requires activation of PKR.

To examine the hypothesis that β -*globin* pre-mRNA functions as an activator of PKR, we studied its ability to activate recombinant human holo-PKR expressed in an RNA-activatable form (rPKR) [14]. We used adenovirus *MINX* pre-mRNA, an efficient template for *in vitro* splicing [21], and the 5'-terminal fragment of human *IFN- γ* mRNA, which is a potent activator of PKR [13, 14] as negative and positive controls, respectively. In contrast to *MINX* pre-mRNA, the β -*globin* pre-mRNA template strongly activated PKR, manifested by autophosphorylation of the kinase (Figure 1H).

Splicing of β -globin mRNA is dependent on PKR activation

Similar to an antibody (Ab) against splicing factor SC35, an Ab against total PKR blocked splicing of β -*glo*-

Figure 1 Splicing of β -globin mRNA is blocked by PKR inhibitors. **(A, B)** Expression of unspliced pre-mRNA (reflected by the 150-nt **(A)** or 353-nt **(B)** band) and spliced mRNA of β -globin (reflected by the 90-nt **(A)** or 210-nt **(B)** band) in HEK293 cells transfected with the human β -globin gene, assayed by RNase protection analysis using RNA probes a **(A)** or b **(B)** (e, exon; i, intron). Where indicated, 15 mM 2-AP was added 18 h after the transfection. Ratio of pre-mRNA/mRNA in the absence ($\circ$) or presence of 2-AP ($\bullet$) is plotted. **(C)** Expression of β -globin RNA in HEK293 cells transfected with the β -globin gene lacking intron 2, assayed with probe a. **(D, E)** Inhibition of β -globin intron 1 excision by 2-AP and a dominant-negative mutant of PKR. BHK-21 cells were stably transfected with fusion constructs p β G(e1-e2)GFP and p β G(e1-i-e2)GFP lacking and containing intron 1, respectively, under the tetracycline-activatable promoter (pTet) **(D)**. Expression of β -globin/GFP protein was induced by the treatment with 12.5 pg/ml Tet for 6 h. Where indicated, 20 mM 2-AP was added 14 h before induction, or cells were transfected with 2 g pPKR Δ 6 DNA [9] 17 h before induction. Confocal microscopy of fluorescence is shown (GFP, orange); PKR Δ 6 panels are at lower magnification **(E)**. **(F, G)** *In vitro* splicing of β -globin pre-mRNA is blocked by 2-AP and okadaic acid. Splicing of capped, 32 P-labeled 477-nt β -globin precursor transcript in HeLa cell nuclear extract was analyzed as described in Materials and Methods section, in the presence of 2-AP **(F)** or okadaic acid **(G)**. Position of intron lariat is indicated in **F**. M, marker DNA; U, input unspliced RNA. Splicing efficiency (SE), defined as a ratio of spliced/unspliced RNA, is provided under each lane. **(H)** Activation of PKR by β -globin pre-mRNA. rPKR was incubated with the 477-nt β -globin pre-mRNA template or 226-nt MINX pre-mRNA (0.75, 2.25, 3.75 and 7.5 nM) or with 203-nt 5'-terminal fragment of human IFN- γ mRNA (0.75 and 2.25 nM) in the presence of [γ - 32 P]ATP as described in Materials and Methods section. Autoradiogram shows phosphorylated PKR.

bin pre-mRNA template *in vitro* (Figure 2A). Anti-PKR Abs against the catalytic domain (Figure 2B) or the RNA-binding domain of the kinase (Figure 2C) blocked splicing of β -globin pre-mRNA template, as did a monoclonal antibody (mAb) specific for phosphorylated PKR (Figure 2D). Thus, β -globin mRNA splicing not only requires PKR but the kinase must be phosphorylated and hence activated. Indeed, splicing was also abrogated by PKR Δ 6 (Figure 2E). Whereas anti-phospho-PKR mAb inhibited splicing of β -globin pre-mRNA template, it did

not affect splicing of *MINX* pre-mRNA, showing specificity (Figure 2F). In the presence of anti-phospho-PKR mAb, unspliced β -globin pre-mRNA accumulated yet there was no accumulation of unspliced *MINX* RNA (Figure 2F).

PKR is found in splicing complexes and is needed for spliceosome formation

To investigate whether PKR associates with the β -globin pre-mRNA splicing complex, we performed the *in*

Figure 2 Splicing of β -globin mRNA depends on PKR activation. (A–F) Splicing of β -globin pre-mRNA template is blocked by anti-PKR Abs and the dominant-negative mutant of PKR. *In vitro* splicing was performed as in Figure 1F. Where indicated, normal goat serum (NGS, 1 μ g) (A, C), polyclonal Ab against SC35 (α SC35), holo-PKR (α PKR) (A), the catalytic domain of PKR (α PKR^{cat}) (B) or the RNA-binding domain of PKR (α PKR^{rbd}) (C), anti-phospho-PKR mAb (α P-PKR) specific for phospho-Thr446 (D, F), or rPKR Δ 6 (E) was added in the amounts shown. Western blot of PKR in reaction mixtures is shown at bottom in E. Effect of α P-PKR on splicing of the β -globin pre-mRNA template and *MINX* pre-mRNA is compared in F. Buffer, corresponding protein storage buffer. SE, splicing efficiency.

in vitro splicing reaction for 30 min and then immunoprecipitated holo-PKR or SC35, the latter of which is present in active splicing complexes [22]. Abs against SC35 precipitated PKR from HeLa cell nuclear extract and vice versa, Abs against PKR precipitated SC35 (Figure 3A). Confocal microscopy of intact cells confirmed that PKR is present in both cytoplasm and nucleus, and it colocalizes with SC35 to the nuclear speckles (Supplementary information, Figure S1A), sites of mRNA splicing [23], supporting the inhibition of splicing by Abs against PKR and SC35 (Figure 2A) and the co-immunoprecipitation of these two proteins (Figure 3A). To investigate whether

PKR is essential for splicing complex formation, we studied kinetics of ribonucleoprotein formation during *in vitro* splicing using labeled β -*globin* pre-mRNA template. Appearance of the spliceosomal complex A preceded that of complexes B and C (Figure 3B). Preincubation of HeLa cell nuclear extract with Abs against the catalytic or RNA-binding domain of PKR sufficed to abrogate formation of complex A yet did not affect formation of early complex H, which is not committed to splicing [24] (Figure 3B). We conclude that PKR associates with splicing complexes and is needed early in spliceosome formation.

Figure 3 PKR is in splicing complexes and required for splicing and spliceosome formation. **(A)** β -*globin* pre-mRNA template was incubated in HeLa cell nuclear extract for 30 min before immunoprecipitation with NGS, α SC35 or α PKR. Western blots of PKR (top) and SC35 (bottom) are shown for soluble (Sup) and precipitated fractions (Pellet). **(B)** Anti-PKR Abs inhibit spliceosome formation. Labeled β -*globin* pre-mRNA template was incubated in HeLa cell nuclear extract for the times shown. Where indicated, α PKR^{cat} or α PKR^{rbd} was added 20 min before incubation. Ribonucleoprotein splicing complexes A, B, C and H were resolved by native PAGE. Left lane shows input unspliced RNA. **(C-D)** Recombinant PKR (rPKR) reconstitutes β -*globin* pre-mRNA splicing upon PKR depletion. **(C)** After adding NaCl to 500 mM, nuclear extract was incubated at 4 °C with α IgG or α PKR Ab for 18 h; the immunoprecipitate was removed using protein A/G Sepharose beads. The depleted extract was centrifuged through Sephadex G-25 before *in vitro* splicing was performed for 2 h, with rPKR or storage buffer (buffer) added as shown. **(D)** Plot of mRNA/pre-mRNA ratio in **C**; α IgG, grey bars; α PKR, black bars. **(E)** In parallel, splicing of MINX RNA was assayed as in **C**. U, input unspliced RNA. **(F)** Reconstitution of spliceosomes on β -*globin* pre-mRNA template with rPKR. Formation of spliceosomes was assayed as in **B**, using extract depleted of PKR as in **C**. Where indicated, 480 ng of rPKR was added back. Three lanes on right show MINX RNA, assayed in parallel.

Recombinant PKR (rPKR) reconstitutes β -globin mRNA splicing upon PKR depletion

Splicing of the β -globin pre-mRNA template was virtually eliminated when PKR was depleted by immunoprecipitation with anti-PKR Ab, whereas anti-IgG had no effect (Figure 3C). Splicing in PKR-depleted nuclear extract was restored in full by adding back rPKR (Figure 3C–3D). By contrast, splicing of *MINX* RNA was completely resistant to PKR depletion, showing specificity (Figure 3E). Likewise, formation of splicing complexes A–C on β -globin pre-mRNA template, but not on *MINX* RNA, was abolished by PKR depletion, and was restored by adding back rPKR (Figure 3F). Of note, to avoid co-immunoprecipitation of splicing factors with PKR, we performed the immunoprecipitation in the presence of 500 mM NaCl [25], and desalted the nuclear extract by gel filtration before the splicing reaction. Splicing and spliceosome formation likewise were restored in full by rPKR when the nuclear extract was desalted by dialysis (Supplementary information, Figure S1B–S1C). These reconstitution experiments validate the specificity of anti-PKR Ab that blocks splicing (Figures 2 and 3) and demonstrate directly that splicing of β -globin mRNA is PKR-dependent.

Addition of rPKR did not further stimulate splicing in α IgG-treated nuclear extract (Figure 3D), indicating that unlike the RNA-mediated activation of PKR, the level of this kinase, an abundant protein in the cell (Supplementary information, Figure S1A), does not limit β -globin mRNA splicing.

eIF2 α phosphorylation is required for splicing of β -globin mRNA

PKR inhibits translation initiation by phosphorylating its substrate eIF2 α . We considered the possibility that eIF2 α phosphorylation might play a role in β -globin mRNA splicing. First, we studied splicing of β -globin pre-mRNA template in nuclear extract in the presence of recombinant vaccinia virus K3L protein, which acts as a pseudosubstrate of PKR that binds the kinase and thus blocks its catalytic site [26, 27]. K3L effectively inhibited splicing of β -globin pre-mRNA template but not that of *MINX* pre-mRNA (Figure 4A).

Because K3L blocks autophosphorylation of PKR and phosphorylation of eIF2 α [26], we next examined the role of eIF2 α phosphorylation in β -globin mRNA splicing. Indeed, eIF2 α was phosphorylated during the splicing reaction (Figure 4B). The anti-phospho-eIF2 α mAb, which did not recognize unphosphorylated eIF2 α (Figure 4B, zero time point), blocked splicing of β -globin pre-mRNA template but not that of *MINX* pre-mRNA (Figure 4B) and completely inhibited formation of spli-

ceosomal complex A on β -globin pre-mRNA template, showing a need for phosphorylated eIF2 α in the early stages of spliceosome assembly (Figure 4C). By contrast, spliceosomal complex assembly on *MINX* RNA was not affected by this mAb (Supplementary information, Figure S2A), demonstrating the selective requirement for phospho-eIF2 in β -globin mRNA splicing. Consistently, addition of recombinant non-phosphorylatable mutant eIF2S51A strongly inhibited splicing of β -globin pre-mRNA template (Figure 4D–4E), whereas addition of *wt* eIF2 α did not (Figure 4D). Splicing of *MINX* pre-mRNA, by contrast, was unaffected by eIF2 α S51A (Figure 4E). Whereas eIF2 α S51A inhibited β -globin mRNA splicing, phosphomimetic mutant eIF2 α S51D did not (Figure 4F), reinforcing the need for phospho-eIF2 α .

We next followed splicing of β -globin mRNA in cells. Expression of non-phosphorylatable mutant eIF2 α S51A [13] strongly reduced the splicing efficiency of β -globin mRNA (Figure 4G and Supplementary information, Figure S2B), whereas expression of *wt* human eIF2 α did not (Figure 4G). These results extend the *in vitro* splicing data and show that phosphorylation of eIF2 α is needed for efficient splicing of full-length β -globin mRNA in the cell. We conclude that eIF2 α plays an essential role in splicing of β -globin mRNA and must be phosphorylated on Ser51 for splicing to occur.

The RNA activator of PKR resides in β -globin exon 1

The β -globin pre-mRNA splicing template activates PKR (Figure 1H), raising the question of where this property maps within the 477-nucleotide (nt) RNA molecule. Progressive truncation from the 3' end showed that the ability to activate PKR was fully maintained in the 5'-terminal 124-nt RNA fragment but lost almost completely in the 5'-terminal 118-nt fragment; removal of the first 4 nts from the 5' end compromised PKR activation (Figure 5A). The RNA activator of PKR thus maps to the first 124 nts of β -globin pre-mRNA comprising the 50-nt 5'-UTR and the first 74 nts of the ORF, and is made up solely of exon 1 sequence.

Probing of the 124-nt RNA element by spontaneous cleavage analysis [28] supports the structure shown in Figure 5B. rPKR induced increased cleavage or protection from cleavage at various positions in the RNA element (Figure 5B), indicating that large sections of the activator RNA interact directly with PKR or undergo conformational changes upon binding of the kinase.

Splicing efficiency is coupled to the ability of the RNA to activate PKR

Our results so far show that β -globin pre-mRNA splicing template potently activates PKR (Figure 1H) through

Figure 4 β -Globin pre-mRNA splicing depends on eIF2 α phosphorylation. **(A)** K3L inhibits splicing of β -globin mRNA. *In vitro* splicing was performed as in Figure 1F, using β -globin pre-mRNA template or MINX pre-mRNA. Before incubation, reaction mixtures were supplemented with recombinant K3L (rK3L) in the amounts shown. **(B)** Splicing of β -globin mRNA requires eIF2 phosphorylation. Right, splicing was assayed as in **A**, in the presence of anti-phospho-eIF2 α mAb (α PeIF2 α ; 3 μ g). Left, western blot of splicing reaction mixture with α PeIF2 α shows mAb specificity and eIF2 α phosphorylation at 70 min. **(C)** α PeIF2 α inhibits early spliceosome formation. Labeled β -globin pre-mRNA template was incubated in HeLa cell nuclear extract for the times shown. Where indicated, α PeIF2 α was added 30 min before addition of the RNA. Ribonucleoprotein complexes A, B, C and H were resolved by native PAGE. Left lane shows input unspliced RNA (Free RNA). **(D-F)** eIF2 α S51A inhibits splicing of β -globin mRNA. Splicing was assayed as in **B**, in the presence of yeast eIF2 α or eIF2 α S51A (**D**, **E**) in the amounts shown, or of 0.5 μ g eIF2 α S51A or eIF2 α S51D (**F**). SE, splicing efficiency. **(G)** Expression of non-phosphorylatable mutant eIF2 α S51A inhibits splicing of β -globin mRNA. HEK293T cells were co-transfected with equal amounts of β -globin gene vector and vector expressing human full-length eIF2 α S51A or eIF2 α WT, or empty vector (Control). At intervals after transfection, levels of spliced and unspliced β -globin transcripts were determined by qRT-PCR; splicing efficiency is expressed as the ratio of spliced RNA over unspliced RNA (means $\pm$ SEM; $n = 3$).

its 5'-terminal domain (Figure 5A) and that activation of PKR is needed for splicing of this RNA (Figures 2, 3B-3F and Supplementary information, Figure S1). We next asked whether during splicing, activation of PKR is accomplished by the splicing template itself. We mutated the short central helix *S1* consisting of 7 bp within the 124-nt RNA element (Figure 5B) to yield *s1a* and *s1b* mutations. The *s1a* and *s1b* mutations disrupt 3 and 5 bp in helix *S1*, respectively; this can explain why the *s1a* mutation partially reduced the ability of the 477-nt β -globin pre-mRNA template to activate PKR, as revealed by assays using rabbit reticulocyte ribosomal fraction containing native PKR (Figure 5C) or rPKR (Figure 5D), whereas the *s1b* mutation severely impaired PKR activation. Ability to activate PKR was restored, at least in part, in the double mutant *s1ab* which retains 5 out of 7 bp in helix *S1* (Figure 5C-5D).

The failure of the *s1b* pre-mRNA template to activate PKR (Figure 5C-5D) and to induce eIF2 α phosphorylation (Supplementary information, Figure S3A) was not due to a lack of the ability to bind PKR, because it competed as well as *wt* RNA for PKR binding in electrophoretic mobility shift analysis (Supplementary information, Figure S3B). Thus, binding of the β -globin RNA to PKR is insufficient for kinase activation, resembling the properties of the RNA activators of PKR in *IFN- γ* [13, 14] and *TNF- α* (manuscript in submission) genes. The phenotype of these mutations was conserved in the 124-nt RNA element with respect to the ability to activate PKR and to induce eIF2 α phosphorylation (Figure 5E). We conclude that helix *S1* shown in Figure 5B is maintained in the β -globin pre-mRNA splicing template.

We next analyzed the effect of these mutations on splicing of the β -globin pre-mRNA template. *s1a* and *s1b* mutations severely impaired splicing efficiency and this was more pronounced for *s1b*. In contrast, the double mutation *s1ab* restored splicing in part (Figure 5F-5G). Notably, the relative splicing efficiency for *wt*, *s1a*, *s1b* and *s1ab* β -globin pre-mRNA templates correlated closely with their ability to activate rPKR at the lower RNA concentration, which was more sensitive to mutations (Figure 5D and 5G). Thus, splicing efficiency in HeLa cell nuclear extract containing all components needed for spliceosome formation reflects the ability of the RNA to activate PKR in an assay where the only macromolecules present are RNA and rPKR.

Reflecting their similar ability to activate native PKR (Figure 5C and 5E), in an independent experiment, *wt* and *s1ab* mutant pre-mRNA templates were spliced with similar efficiency, which was inhibited by anti-phospho-eIF2 α mAb to a similar extent (Figure 5H). Thus, compensatory mutation in *s1ab* RNA not only restores

PKR activation and splicing to *s1b* mutant RNA but such splicing remains as dependent on eIF2 α phosphorylation as splicing of *wt* RNA. Even though *s1b* RNA is severely impaired in splicing, its low residual activity likewise requires phospho-eIF2 α (Figure 5H).

Mutation of the PKR activator also affected β -globin mRNA splicing in the cell, where the *s1b* mutation strongly impaired intron 1 excision (Figure 5I). The ratio of pre-mRNA/mRNA declined over time for the *wt* gene as expected yet increased for the *s1b* gene, attesting to a block in pre-mRNA processing. These results show that the splicing efficiency of the β -globin pre-mRNA template is regulated by its ability to activate PKR through the 124-nt exonic RNA element.

PKR activation and eIF2 α phosphorylation enable splicing of γ -globin mRNA

Fetal γ -globin genes are related to the adult β -globin gene. Except for nt 25 in the 5'-UTR, γ - and γ -globin gene sequences are identical through exon 2 [29]. We examined whether activation of PKR might also regulate excision of γ -globin intron 1. Indeed, as for β -globin (Figure 2E), PKR Δ 6 strongly inhibited splicing of γ -globin pre-mRNA template containing exon 1, intron 1 and the first 172 nts of exon 2 (Supplementary information, Figure S3C). γ -Globin and β -globin pre-mRNA templates activated PKR comparably (Supplementary information, Figure S3D). γ -Globin exon 1 RNA was sufficient to activate the kinase and induce eIF2 α phosphorylation (Supplementary information, Figure S3E). Splicing of γ -globin and β -globin pre-mRNA templates was similarly inhibited by anti-phospho-eIF2 α mAb (Supplementary information, Figure S3F). Thus, PKR activation and eIF2 phosphorylation also enable splicing of γ -globin mRNA.

The ability of β -globin pre-mRNA to activate PKR is silenced upon splicing

During erythroid development, the cellular globin content rises from < 0.1% of the total protein amount in proerythroblasts to 95% in reticulocytes, reflecting a massive translation of globin mRNA [15]. Erythropoiesis would be compromised severely if mature β -globin mRNA were to activate PKR and thereby inhibit its own translation, in a manner resembling that for *IFN- γ* mRNA [13]. In reticulocyte lysates, rabbit globin mRNA does not detectably activate PKR [30]. Because the activator of PKR in β -globin pre-mRNA splicing template is made up solely of exon sequence (Figure 5A), there is a need for a mechanism by which the ability to activate PKR is blocked in mature human β -globin mRNA. Indeed, whereas the 477-nt β -globin pre-mRNA template strong-

Figure 5 Splicing efficiency is coupled to activation of PKR by the activator in β -globin exon 1. **(A)** Activation of PKR by β -globin RNA. Phosphorylation of rPKR was assayed in the presence of [γ - 32 P]ATP, and 5 and 10 nM intact β -globin pre-mRNA template (nts 1-477) or truncated RNAs (nts 1-124, 5-124 and 1-118). **(B)** In-line structure probing of the 5'-terminal 124 nts of the β -globin pre-mRNA template. Gel shows representative pattern of spontaneous cleavages in 5' end-labeled RNA incubated alone (-) or with rPKR (nM) as shown. G (T1) and alkali (OH) ladders serve as size markers; NR, input RNA. Structure on left is based on repeated in-line probing. Red and green colors denote nt showing enhanced and reduced cleavage, respectively, induced by rPKR. In helix S1, boxes denote nts mutated in *s1a* and *s1b*. **(C-E)** In *s1a*, positions 47-51 were mutated to GUGGU; in *s1b*, positions 110-114 were mutated to GUACC (red sequences in **C**). Activation of rabbit reticulocyte PKR (**C**, **E**) or rPKR (**D**) by wt or mutant forms of the 477-nt (**C**, **D**) or 124-nt (**E**) RNA was assayed; eIF2 α phosphorylation is shown in **E**. For activation of rabbit reticulocyte PKR, 1.2 and 2 nM RNA was used, and for rPKR, 5 and 10 nM RNA was used. **(F, G)** Splicing efficiency is coupled to the ability to activate PKR. **(F)** Splicing of wt and mutant forms of β -globin pre-mRNA template in HeLa cell nuclear extract was analyzed as in Figure 1F; left four lanes, input unspliced RNA (U). **(G)** Comparison of the ability to activate PKR at the lower RNA concentration in **D** and splicing efficiency (ratio of spliced RNA over pre-mRNA) in **F**. **(H)** Splicing of *s1b* pre-mRNA template is restored by compensatory mutation. *In vitro* splicing was assayed in the absence or presence of α PelF2 α (1 μ g); splicing efficiency is plotted in the bar graph. **(I)** β -globin precursor RNA processing is coupled to the ability to activate PKR. HEK293T cells were transfected with wt- or *s1b*-mutant β -globin gene and excision of intron 1 was assayed by quantitative RT-PCR; ratio of unspliced over spliced RNA is plotted (means $\pm$ SEM; $n = 3$).

ly activated PKR (Figure 6A, *e-i-e*), the spliced mRNA product consisting of exon 1 and the first 205 nts of exon 2 activated PKR weakly (at high RNA concentration) or not at all (at low RNA concentration) (Figure 6A, *e/e*). Apparently, upon excision of intron 1, the RNA activator of PKR in β -globin pre-mRNA template is silenced.

To explore the silencing mechanism, we first truncated the second exon from the 3' end. β -globin mRNA fragment containing exon 1 and the first 21 nts of exon 2 failed to activate PKR significantly; even truncation that left only 10 nts of exon 2 fused to exon 1 sufficed to impair the ability to activate PKR (Figure 6A). This result suggests that when juxtaposed to exon 1, the first 10 nts of exon 2 are able to silence the PKR activator in exon 1. The first 10 nts of exon 2 sequence contains a 5'-UGGUG-3' motif (*Slc*) that in principle could anneal with the 3'-ACCAC-5' motif in *Slb*, forming 5 stable bp and displacing the corresponding *Slb* motif (Figure 6B, modeled in Supplementary information, Figure S4A). To test this concept, we mutated the 5'-CACCA-3' motif in *Slb* to 5'-GUGGU-3' and/or changed the *Slc* motif to 5'-AC-CAC-3' in the spliced *e/e* RNA. *e/e* RNA fragments with the resulting *slb* or *slc* mutation induced activation of PKR. The double mutation, *e/e slac*, which restores base pairing between *Slb* and *Slc*, also restored silencing of PKR activation (Figure 6C).

The 477-nt β -globin pre-mRNA template with the *slb* mutation, which partially retains the base pairing within helix *SI*, still possessed partial ability to activate PKR (Figure 5C). As the *slb* mutation abrogates base pairing with the *Slc* silencer sequence, this could prevent the disruption of base pairing between *Slb* and *Slc* sequences after splicing (Figure 6B), and thus allow for residual PKR activation, as is indeed observed in Figure 6C.

We conclude that upon excision of intron 1, base pairing between *Slb* and *Slc* sequences within helix *SI*, critical for activation of PKR by β -globin pre-mRNA template (Figure 5C-5E), is abrogated through displacement of *Slb* by *Slc* sequence located just downstream of the exon junction, within the first 10 nts of exon 2. This results in silencing of the ability of the mature mRNA product to activate PKR. Silencing thus requires strand replacement within helix *SI* and as shown by *e/e slac*, this process tolerates sequence inversion of the two complementary strands. Phylogenetic analysis shows that except for mouse β -globin major RNA, the potential for *Slb-Slc* base pairing through the AUG codon is conserved, even in teleost fish (Supplementary information, Figure S4B).

An RNA activator of PKR resides in α -globin exon 2

A hallmark of normal erythropoiesis is efficient syn-

thesis of α - and β -globin in an equimolar ratio. We therefore investigated whether splicing of α -globin mRNA is also regulated by activation of PKR and phosphorylation of eIF2 α . In cells transfected with the human $\alpha 2$ -globin gene, addition of 2-AP led to prompt accumulation of unspliced pre-mRNA (Figure 7A and Supplementary information, Figure S5). Splicing of the α -globin pre-mRNA template comprising exon 1, intron 1 and all but 9 nts of exon 2 in HeLa cell nuclear extract was blocked by anti-phospho-PKR mAb (Figure 7B) and was at least as sensitive as splicing of β -globin pre-mRNA template to anti-phospho-eIF2 α mAb (Figure 7C). α -Globin pre-mRNA template was as potent an activator of PKR as β -globin pre-mRNA template (Figure 7D). In contrast to β -globin, however, the α -globin RNA activator of PKR mapped to exon 2 (Figure 7E). 3'-truncation of exon 2 RNA showed that a full PKR activation requires the first 179 nts of exon 2 (nts 250-428) and was largely lost upon removal of five additional nts from the 3' end; 5'-truncations defined a 123-nt core (nts 306-428) as the minimal activator (Figure 7F).

A silencer of PKR activation resides in α -globin exon 1

Excision of intron 1 almost totally eliminated the ability of α -globin RNA to activate PKR (Figure 7G). PKR activation was restored in part by truncation of the first 36 nts of exon 1 and fully by deletion of the first 73 nts, placing the silencer of PKR activation within the 5'-proximal region of α -globin exon 1 (Figure 7G). Thus, for both α - and β -globin genes, splicing of intron 1 depends on PKR activation and eIF2 α phosphorylation, but the respective positions of activator and silencer are reversed between the adjoining exons.

Discussion

Our finding is that splicing of globin mRNA requires not only activation of PKR by an intragenic RNA activator but also phosphorylation of the PKR substrate, eIF2 α . There are similar requirements for the *TNF- α* gene encoding an inflammatory cytokine (manuscript in submission) suggesting the generality of our findings. Hemoglobin synthesis depends on optimal translation of globin mRNA in reticulocytes and globin reaches 95% of the total protein amount under conditions where no further transcription takes place [15]. Hence, local activation of PKR by globin mRNA, resulting in phosphorylation of eIF2 α and translational downregulation, would be detrimental. It is thus surprising that globin mRNA splicing depends strictly on activation of PKR as well as on phosphorylation of eIF2 α , with the RNA activator of PKR not only being intrinsic to the globin pre-mRNA transcript

On the basis of analysis in cells and *in vitro*, our data

SPRINGER NATURE | Cell Research | Vol 27 No 5 | May 2017

not activated by RNA [2] be capable of supporting β -globin mRNA splicing, these approaches would not have yielded an inhibition. Once PKR is activated, eIF2 α must be phosphorylated on Ser51 to allow formation of early-stage spliceosomal complex A. Splicing was inhibited by K3L, a viral eIF2 α pseudosubstrate of PKR, by anti-phospho-eIF2 α mAb and by non-phosphorylatable mutant eIF2 α S51A, yet not by phosphomimetic eIF2 α S51D. As for *TNF- α* (manuscript in submission), expression of eIF2 α S51A blocked splicing of full-length β -globin mRNA in cells. The dependence of splicing on eIF2 α phosphorylation early during β -globin spliceosome assembly provides a mechanistic explanation for the need for PKR activation in splicing.

Upon excision of β -globin intron 1, a 5-nt exon 2 sequence *c*, located just downstream of the exon junction, displaces strand *b* of short helix *a-b* in the RNA activator of PKR in exon 1, which is critical for the ability of β -globin pre-mRNA to activate PKR (Figures 6 and 8). This displacement results in silencing of the ability of the exonic RNA element to activate PKR. Mutation of either strand *a* or *c* in this helix was sufficient to allow the spliced RNA product to activate PKR, whereas double mutation that restored base pairing resulted in restoration of silencing of the ability to activate PKR. In this manner, activation of PKR is eliminated once intron 1 has been excised. This RNA-mediated silencing mechanism ensures that the local activation of PKR serves solely to enable splicing.

IFN- γ mRNA attenuates its own translation by locally activating PKR [13] through a structure comprised of the 5'-UTR and the first 26 codons, causing it to be disrupted by ribosomes during translation and requiring dynamic refolding to restore the activator [14]. Although the β -globin activator of PKR likewise spans the 5'-UTR and the first 25 codons of the ORF, β -globin mRNA does not activate PKR after splicing owing to the silencer mechanism.

α - and β -globin proteins are produced in an equimolar ratio but their genes differ in terms of chromosomal location, gene dosage, transcriptional control and mRNA stability [31, 32] as well as in the ability of their mRNAs to compete in translation [33, 34]. We have shown that α - and β -globin genes share a regulatory strategy, i.e., dependence of mRNA splicing on activation of PKR and phosphorylation of eIF2 α , but differ in the location of the elements that activate PKR and silence this property. Whereas the β -globin RNA activator of PKR is located in exon 1, the α -globin activator is composed solely of protein-encoding sequence in exon 2. Conversely, in β -globin mRNA the silencer of PKR activation is located in exon 2, whereas in α -globin mRNA it resides in the 5'-half of exon 1. These distinct solutions demonstrate

evolutionary flexibility in achieving the common aim of bringing mRNA splicing under the control of PKR activation. The first introns of α - and β -globin thus serve a novel function, to separate the PKR activator required for splicing from the silencer of PKR activation needed to enable translation.

Our experiments link eIF2 α phosphorylation to an early step in spliceosome assembly on *globin* mRNA, a complex and dynamic process in which the roles of multiple individual proteins remain to be delineated [35, 36]. Whereas eIF2 α phosphorylation inhibits translation, we demonstrate a positive role for phosphorylation of eIF2 α on Ser51: it is necessary for splicing of *globin* mRNA. This requirement is shared with splicing enabled by the RNA activator of PKR in the *TNF- α* 3'-UTR, which is likewise abrogated by eIF2 α S51A expression (manuscript in submission). Splicing efficiency of *TNF- β* pre-mRNA, which lacks a PKR activator, is strongly enhanced not only by inserting the *TNF- α* activator but also by merely increasing the level of phospho-eIF2 α globally in the cell by using an inhibitor of eIF2 α dephosphorylation, demonstrating that eIF2 α phosphorylation is both necessary and sufficient to achieve efficient splicing. Control of *TNF* mRNA splicing by eIF2 α phosphorylation operates in primary human immune cells, showing its physiological relevance (manuscript in submission).

Our results demonstrate a novel nuclear role for PKR and eIF2 α phosphorylation. A nuclear extract displays multiple important properties: the need for PKR in complex A formation on β -globin splicing template; inhibition of β -globin pre-mRNA splicing upon depletion of PKR and restoration of splicing by recombinant human PKR; restoration of complex A formation on a β -globin splicing template; phosphorylation of eIF2 α and the need for phospho-eIF2 α in complex A formation.

Differentiating human erythroleukemia cells do not express β -globin; very low levels of β -globin mRNA were detected, relative to α -globin mRNA, in conditions inhibiting cell proliferation [37]. Even before induction of differentiation, K562 cells expressed fully spliced α -globin mRNA, whereas pre-mRNA was not detectable (LSN, data not shown). We overcame the limitations of studying splicing of adult human *globin* mRNA in human cell lines by *in vitro* splicing assays and by genetic approaches, including gain-of-function mutations that provide mechanistic insight. Data from *in vitro* analysis of *globin* mRNA splicing were validated in transfected human and hamster cells.

Whereas human *TNF- α* and *IFN- γ* genes contain potent intragenic activators of PKR, these elements display structural defects in the mouse (manuscript in submission, [14]). The mouse also differs significantly in terms

Figure 7 Excision of $\alpha 2$ -globin intron 1 depends on PKR activation and eIF2 α phosphorylation. **(A)** 2-AP inhibits splicing of α -globin mRNA. HEK293T cells were transfected with the human $\alpha 2$ -globin gene. 2-AP (15 mM) was added at 18 h after transfection. At times shown, expression of unspliced and spliced RNA was assayed by RNase protection analysis (Supplementary information, Figure S5). Pre-mRNA/mRNA ratio is plotted. **(B, C)** mAbs against phospho-PKR and phospho-eIF2 α block splicing of α -globin pre-mRNA template. Splicing of capped, 32 P-labeled 445-nt α -globin pre-mRNA template in HeLa cell nuclear extract was analyzed as in Figure 2D, in the absence or presence of 1 μ g α P-PKR mAb **(B)** and for splicing of α - and β -globin pre-mRNA templates, in the absence or presence of 3 μ g α P-eIF2 α mAb **(C)**. SE, splicing efficiency. **(D, E)** The α -globin RNA activator of PKR maps to exon 2. Activation of rPKR (top) or rabbit reticulocyte PKR (bottom) was assayed in the presence of 445-nt α -globin or 477-nt β -globin pre-mRNA template (e1-i-e2) **(D)** or in the presence of 445-nt pre-mRNA template (e1-i-e2), 132-nt exon 1 RNA (e1) or 196-nt exon 2 RNA (e2) of α -globin **(E)**. For activation of rabbit reticulocyte PKR, 1.2 and 2 nM RNA was used and for rPKR, 5 and 10 nM RNA was used. **(F)** Definition of the minimal α -globin RNA activator of PKR. Activation of rPKR was analyzed in the presence of 5 and 10 nM α -globin exon 2 RNA truncated to the positions shown (see map at bottom). **(G)** Excision of α -globin intron 1 causes silencing of the ability to activate PKR. Activation of rPKR was analyzed in the presence of 5 and 10 nM α -globin RNA as shown (e/e, 328-nt spliced form). The silencer maps to nts 1-74 in exon 1, shown in red in the map.

Figure 8 Model for regulation of β -globin mRNA splicing by an RNA activator of PKR in exon 1 and silencing of the PKR activator upon intron 1 excision. The activator of PKR in exon 1 relies on 5 bp *a-b* pairing in helix *S1*. Activation of PKR results in phosphorylation of eIF2 α , a prerequisite for excision of intron 1 (top). Excision of intron 1 juxtaposes exon 2 sequence *c* and exon 1 sequence *a*, causing strand displacement in helix *S1* that silences the ability to activate PKR (bottom).

of β -globin gene organization, expressing major and minor species [38], impeding facile comparison with the human gene.

Synthesis of heme and globin is coordinated through a feedback loop wherein heme is needed to block activation of the eIF2 kinase HRI, thereby preventing attenuation of globin mRNA translation [39, 40]. Phosphorylation of eIF2 α by two distinct kinases, PKR and HRI, thus regulates separate steps in globin gene expression in nucleus and cytoplasm, i.e., mRNA splicing and translation, respectively. In the cytoplasm, eIF2 α is found in complex with eIF2 β and γ chains, forming holo-eIF2 active as translation factor [41]. Whether eIF2 β and γ subunits are involved in globin mRNA splicing is not yet known.

Materials and Methods

Gene transfection

Human embryonic kidney HEK293T cells were transiently transfected with 2 μ g pBS vector carrying the 5 kb *PstI-PstI*

fragment of genomic β -globin DNA including the upstream and downstream regulatory sequences. β -globin gene lacking intron 2 was constructed by replacing sequence downstream from the *Pml*I site in exon 2 with cDNA sequence. pBG(e1-i-e2)GFP and pBG(e1-e2)GFP contain β -globin exons 1 and 2, with and without intron 1, respectively, under the tetracycline-inducible promoter, exon 2 being fused in-frame to the *EGFP* ORF at the *Bam*HI site 18 bp upstream from the 3' end of exon 2. These two vectors were used to stably transfect BHK-21 cells together with pTet-tTAK (Gibco BRL) in a ratio of 1:10. After four passages in DMEM medium, transfected cells were induced with 12.5 pg/ml tetracycline; cells showing over 70% fluorescence were selected. Expression of PKR Δ 6 was performed as described [13].

Splicing assay by RNase protection analysis

To generate β -globin intron 1 splicing probe (probe a), genomic DNA was digested with *Nco*I and *Nsp*I and the filled-in fragment ligated into *Hinc*II-digested pBS vector. To generate β -globin intron 2 splicing probe (probe b), 5'-CGGAATTCTACCCTTGACCCAGAG-3' and 5'-CCCAAGCTTCGATCCTGAGACTTCAC-3' were used as PCR primers on genomic DNA; the *Eco*RI-*Hind*III fragment was excised from the product and inserted into pBS vector. Total RNA was isolated with TRIzol Reagent (Invitrogen) and subjected to RNase protection analysis with RNases A and T1 [12, 42]. Band intensity in autoradiograms was quantitated with a Fujifilm BAS 1800II Phosphorimager.

Splicing assay by quantitative real-time PCR

Total cellular RNA was harvested using TRIzol Reagent (Invitrogen) and cDNA synthesized from 1 μ g RNA using Verso RT-PCR Kit (ABgene) in a 20- μ l reaction volume. Quantitative real-time PCR was performed using 0.25 μ g cDNA, 300 nM primers/probe Double-Dye (Taqman style) mix and Absolute Blue QPCR Mix (ABgene). Primers and probes specific for spliced (primers: 5'-AAGGTGAACGTGGATGAAGT-3', 5'-AGCATCAGGAGTGGACAGAT-3'; probe: 5'-CCTGGGCAGGCTGCTGGTG-GTC-3') and unspliced (primers: 5'-TATCAAGGTTACAAGACAGGTTTAAG-3', 5'-GCCTAAGGGTGGGAAAATAGAC-3'; probe: 5'-CTCTGGGTTTCTGTATAGGCACTGACTCT-3') forms of β -globin RNA and normalizing human β -actin gene were from PrimerDesign. Reactions were performed for 40 cycles (95 $^{\circ}$ C for 15 s, 60 $^{\circ}$ C for 30 s and 72 $^{\circ}$ C for 15 s) using a Rotor-Gene 6000 instrument (Corbett Life Science). The $\Delta\Delta$ Ct method was used to evaluate mRNA-to-pre-mRNA ratio. Amplification efficiency of spliced and unspliced forms, determined by CT slope, was similar and at least 99%.

In vitro splicing

HeLa cell nuclear extract (Promega) and capped, 32 P-labeled 477-nt human β -globin precursor SP6 transcript generated from *Bam*HI-cut DNA template (0.17 pmol, 1.25×10^5 cpm/pmol) or adenovirus *MINX* SP6 transcript generated from *Bam*HI-cut DNA template [21] were incubated for 3 h at 30 $^{\circ}$ C in splicing buffer (reaction volume, 25 μ l) following instructions of the manufacturer (Promega). When the effect of α P-PKR (Calbiochem) or α P-eIF2 α (Abcam) was studied, the reaction was preincubated with Ab for 15 min at 30 $^{\circ}$ C and 15 min at 4 $^{\circ}$ C before RNA and RNasin were added; for α PKR (Cell Signaling), α PKR^{cat}, and α PKR^{rbd} (Santa Cruz), preincubation was carried out for 15 min at 4 $^{\circ}$ C. RNA was

extracted, precipitated and analyzed in 4% polyacrylamide/8 M urea gels. For analysis of spliceosome formation, 4- μ l samples were collected at time intervals, incubated for 10 min with 50 μ g heparin at 4 °C, made 10% glycerol and separated on 3.75% native polyacrylamide gels (polyacrylamide:bis, 80:1; 50 mM Tris-base, 50 mM glycine, pH 8.8). The α -*globin* pre-mRNA template for *in vitro* splicing, generated from *HincII*-digested human $\alpha 2$ -*globin* DNA, was a 445-nt SP6 transcript. The γ -*globin* pre-mRNA template was a 439-nt T3 transcript. 5'-end labeled *HinfI*-digested ϕ X174 single-stranded DNA (Fermentas) served as size marker, except in Figures 1G and 2A where boiled 100-bp DNA ladder (New England Biolabs) was used.

Co-immunoprecipitation of PKR and SC35

Antibodies were bound to protein A-Sepharose (GE Healthcare) by incubation for 2 h at 4 °C. Immunoprecipitation from HeLa nuclear extract was performed by addition of 12 μ g α PKR, 26.4 μ g α SC35 (Santa Cruz) or 60 μ g NGS followed by incubation for 5 h at 4 °C. Complexes were washed five times with the buffer containing 50 mM Tris-HCl (pH 7.4), 5 mM EDTA, 150 mM NaCl, 1 M KCl, 0.2% bovine serum albumin, and 0.5% Triton X-100.

Mutagenesis

β -*globin* *s1a* mutant was generated by replacing the *AleI*-*AleI* fragment in *wt* 1-477 bp DNA with 5'-CTGTGTTCACTAGCAACCTCAAACAGAGTGGT-3' annealed to 5'-ACCACTCTGTTTGAGGTTGCTAGTGAACACAG-3'. *s1b* was created by PCR using phosphorylated primers 5'-GAAGTACCATGAAGTTGGTGGTGAGG-3' and 5'-ACCTGCCCCACAGGG-3' with the *wt* 1-477 bp DNA template and KOD polymerase (Novagen) to generate linear full-length plasmid DNA carrying blunt ends; the product was purified on a 1% agarose gel and self-ligated. Mutant *e/e s1c* was generated from β -*globin* cDNA under the SP6 promoter by PCR, using primers 5'-TGCACCACGTCTACCCTTGGACCCAGAGG-3' and 5'-GCCTGCCCAGG-GCCTCA-3'. Truncations were created using restriction enzyme digestion, ligation to annealed complementary oligonucleotides, or PCR; primer sequences will be supplied on request. Truncation of $\alpha 2$ -*globin* and γ -*globin* exon 1/intron 1/exon 2 DNA was done by PCR using phosphorylated DNA primers with KOD polymerase as above.

In-line structure probing of RNA

For in-line probing of RNA, we adapted a previously described protocol [28]. About 1 nM 5' 32 P-labeled RNA was incubated for 40 h at 25 °C in the buffer containing 20 mM MgCl₂/50 mM Tris-HCl pH 8.3, 100 mM KCl, 1.5 mM dithiothreitol and 6% (v/v) glycerol in the absence or presence of rPKR. RNA cleavage products were resolved by 10% PAGE.

Recombinant proteins

PKR, PKRA6 and K3L were cloned into pHTT7K [43] and expressed in *Escherichia coli* Rosetta(DE3)pLysS (Invitrogen) as full-length N-terminally hexahistidine-tagged proteins. PKR was expressed in RNA-activatable form and was > 98% pure on SDS-12% PAGE [14]. K3L was recovered from inclusion bodies by solubilization in 6 M urea buffer and loaded onto a Ni-column that was eluted stepwise with imidazole buffer containing 6 M urea; next, urea concentration was reduced 10-fold in six dialysis steps.

K3L was renatured by dialysis against 0.5 M arginine, 20% glycerol, phosphate-buffered saline before arginine concentration was reduced five-fold in four dialysis steps; it was stored in the final buffer. *Saccharomyces cerevisiae* eIF2 α , eIF2 α S51A and eIF2 α S51D were cloned into pET15b and expressed as C-terminally truncated hexahistidine-tagged 25 kDa proteins (manuscript in submission).

Activation of PKR

Uncapped RNA transcripts were purified by Sephadex G-50 gel chromatography, washed with ethanol and eluted with water as described [44]. Rabbit reticulocyte ribosomal fraction was prepared and activation of PKR by RNA was assayed as described [45]. Activation of 100 ng rPKR by RNA was determined as described [14]. Unless otherwise noted, RNA concentrations used were 1.2 and 2 nM for rabbit reticulocyte ribosome fraction and 5 and 10 nM for rPKR.

Immunodepletion and reconstitution of PKR

For immunodepletion of PKR from HeLa nuclear extract, 200 μ l of the extract was brought to 500 mM NaCl [25], first incubated with 12 μ g of α IGG or α PKR Ab for 18 h at 4 °C with head-over-tail rotation, and then incubated with washed protein A/G Sepharose beads for 2 h at 4 °C with head-over-tail rotation before the nuclear extract was separated from the beads. To remove NaCl, the depleted nuclear extract was either centrifuged through Sephadex G-25 or desalted by overnight dialysis in buffer (20 mM Hepes pH 7.9, 0.5 mM dithiothreitol, 20% (v/v) glycerol, and 0.5 mM phenylmethylsulfonyl fluoride) before *in vitro* splicing was performed for 2 h with or without rPKR or storage buffer.

Acknowledgments

We thank Blanka Shanitzki for assistance in quantitative real-time PCR. This work was supported by grants from the Israel Science Foundation.

Author Contributions

LI, FO, LSN and RK designed research; LI, FO, LSN, EE, SC-C, YB-A and YB performed research and analyzed data; RK, LI and LSN wrote the manuscript.

Competing Financial Interests

The authors declare no competing financial interests.

References

- 1 Nallagatla SR, Toroney R, Bevilacqua PC. Regulation of innate immunity through RNA structure and the protein kinase PKR. *Curr Opin Struct Biol* 2011; **21**:119-127.
- 2 Sonenberg N, Hinnebusch AG. Regulation of translation initiation in eukaryotes: mechanisms and biological targets. *Cell* 2009; **136**:731-745.
- 3 Harding HP, Zhang Y, Zeng H, *et al.* An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell* 2003; **11**:619-633.
- 4 Muaddi H, Majumder M, Peidis P, *et al.* Phosphorylation of eIF2 α at serine 51 is an important determinant of cell survival and adaptation to glucose deficiency. *Mol Biol Cell* 2010;

- 21:3220-3231.
- 5 Tsaytler P, Harding HP, Ron D, Bertolotti A. Selective inhibition of a regulatory subunit of protein phosphatase 1 restores proteostasis. *Science* 2011; **332**:91-94.
- 6 Kaempfer R. Identification and RNA-binding properties of an initiation factor capable of relieving translational inhibition induced by heme deprivation or double-stranded RNA. *Biochem Biophys Res Commun* 1974; **61**:591-597.
- 7 Farrell PJ, Balkow K, Hunt T, Jackson RJ, Trachsel H. Phosphorylation of initiation factor eIF-2 and the control of reticulocyte protein synthesis. *Cell* 1977; **11**:187-200.
- 8 Zhang F, Romano PR, Nagamura-Inoue T, et al. Binding of double-stranded RNA to protein kinase PKR is required for dimerization and promotes critical autophosphorylation events in the activation loop. *J Biol Chem* 2001; **276**:24946-24958.
- 9 Dey M, Cao C, Dar AC, et al. Mechanistic link between PKR dimerization, autophosphorylation, and eIF2 α substrate recognition. *Cell* 2005; **122**:901-913.
- 10 Bevilacqua PC, Cech TR. Minor-groove recognition of double-stranded RNA by the double-stranded RNA-binding domain from the RNA-activated protein kinase PKR. *Biochemistry* 1996; **35**:9983-9994.
- 11 Manche L, Green SR, Schmedt C, Mathews MB. Interactions between double-stranded RNA regulators and the protein kinase DAI. *Mol Cell Biol* 1992; **12**:5238-5248.
- 12 Osman F, Jarrous N, Ben-Asouli Y, Kaempfer R. A cis-acting element in the 3'-untranslated region of human TNF- α mRNA renders splicing dependent on the activation of protein kinase PKR. *Genes Dev* 1999; **13**:3280-3293.
- 13 Ben-Asouli Y, Banai Y, Pel-Or Y, Shir A, Kaempfer R. Human interferon-gamma mRNA autoregulates its translation through a pseudoknot that activates the interferon-inducible protein kinase PKR. *Cell* 2002; **108**:221-232.
- 14 Cohen-Chalamish S, Hasson A, Weinberg D, et al. Dynamic refolding of IFN- γ mRNA enables it to function as PKR activator and translation template. *Nat Chem Biol* 2009; **5**:896-903.
- 15 Nienhuis AW, Benz EJ. Regulation of hemoglobin synthesis during the development of the red cell. *N Engl J Med* 1977; **297**:1318-1328.
- 16 Samuel CE. Role of the RNA-dependent protein kinase in the regulated expression of genes in transfected cells. *Pharmacol Ther* 1992; **54**:307-317.
- 17 Custódio N, Carvalho C, Condado I, et al. In vivo recruitment of exon junction complex proteins to transcription sites in mammalian cell nuclei. *RNA* 2004; **10**:622-633.
- 18 Koromilas AE, Roy S, Barber GN, Katze MG, Sonenberg N. Malignant transformation by a mutant of the IFN-inducible dsRNA-dependent protein kinase. *Science* 1992; **257**:1685-1689.
- 19 Krainer AR, Maniatis T, Ruskin B, Green MR. Normal and mutant human β -globin pre-mRNAs are faithfully and efficiently spliced *in vitro*. *Cell* 1984; **36**:993-1005.
- 20 Mermoud JE, Cohen P, Lamond AI. Ser/Thr-specific protein phosphatases are required for both catalytic steps of pre-mRNA splicing. *Nucleic Acids Res* 1992; **20**:5263-5269.
- 21 Hartmuth K, Urlaub H, Vornlocher HP, et al. Protein composition of human prespliceosomes isolated by a tobramycin affinity-selection method. *Proc Natl Acad Sci USA* 2002; **99**:16719-16724.
- 22 Fu XD, Maniatis T. The 35-kDa mammalian splicing factor SC35 mediates specific interactions between U1 and U2 small nuclear ribonucleoprotein particles at the 3' splice site. *Proc Natl Acad Sci USA* 1992; **89**:1725-1729.
- 23 Spector DL, Fu XD, Maniatis T. Associations between distinct pre-RNA splicing components and the cell nucleus. *EMBO J* 1991; **10**:3467-3481.
- 24 Michaud S, Reed R. An ATP-independent complex commits pre-mRNA to the mammalian spliceosome assembly pathway. *Genes Dev* 1991; **5**:2534-2546.
- 25 Makarova OV, Makarov EM, Lührmann R. The 65 and 110 kDa SR-related proteins of the U4/U6.U5 tri-snRNP are essential for the assembly of mature spliceosomes. *EMBO J* 2001; **20**:2553-2563.
- 26 Carroll K, Elroy-Stein O, Moss B, Jagus R. Recombinant vaccinia virus K3L gene product prevents activation of double-stranded RNA-dependent, initiation factor 2 α -specific protein kinase. *J Biol Chem* 1993; **268**:12837-12842.
- 27 Dar AC, Sichieri F. X-ray crystal structure and functional analysis of vaccinia virus K3L reveals molecular determinants for PKR subversion and substrate recognition. *Mol Cell* 2002; **10**:295-305.
- 28 Soukup GA, Breaker RR. Relationship between internucleotide linkage geometry and the stability of RNA. 1999; *RNA* **5**:1308-1325.
- 29 Slightom JL, Blechl AE, Smithies O. Human fetal $\epsilon\gamma$ - and $\alpha\gamma$ -globin genes: complete nucleotide sequences suggest that DNA can be exchanged between these duplicated genes. *Cell* 1980; **21**:627-638.
- 30 Rosen H, Di Segni G, Kaempfer R. Translational control by messenger RNA competition for eukaryotic initiation factor 2. *J Biol Chem* 1982; **257**:946-952.
- 31 Higgs DR, Garrick D, Anguita E, et al. Understanding α -globin gene regulation: aiming to improve the management of thalassemia. *Ann NY Acad Sci* 2005; **1054**:92-102.
- 32 Mahajan MC, Karmakar S, Weissman SM. Control of beta globin genes. *J Cell Biochem* 2007; **102**:801-810.
- 33 Lodish HF. Alpha and beta globin messenger ribonucleic acid. Different amounts and rates of initiation of translation. *J Biol Chem* 1971; **246**:7131-7138.
- 34 Di Segni G, Rosen H, Kaempfer R. Competition between α - and β -globin messenger ribonucleic acids for eukaryotic initiation factor 2. *Biochemistry* 1979; **18**:2847-2854.
- 35 Wahl MC, Will CL, Lührmann R. The spliceosome: design principles of a dynamic RNP machine. *Cell* 2009; **136**:701-718.
- 36 Hoskins AA, Moore MJ. The spliceosome: a flexible, reversible macromolecular machine. *Trends Biochem Sci* 2012; **37**:179-188.
- 37 Shiraishi M, Yamamoto Y, Hirooka N, et al. A high concentration of triiodothyronine attenuates the stimulatory effect on hemin-induced erythroid differentiation of human erythroleukemia K562 cells. *Endocr J* 2015; **62**:431-440.
- 38 Hoffmann FG, Opazo JC, Storz JF. New genes originated via multiple recombinational pathways in the beta-globin gene family of rodents. *Mol Biol Evol* 2008; **25**:2589-2600.
- 39 Kaempfer R, Kaufman J. Translational control of hemoglobin

- synthesis by an initiation factor required for recycling of ribosomes and for their binding to messenger RNA. *Proc Natl Acad Sci USA* 1972; **69**:3317-3321.
- 40 Chen JJ. Regulation of protein synthesis by the heme-regulated eIF2 α kinase: relevance to anemias. *Blood* 2007; **109**:2693-2699.
- 41 Aitken CE, Lorsch JR. A mechanistic overview of translation initiation in eukaryotes. *Nat Struct Mol Biol* 2012; **19**:568-576.
- 42 Jarrous N, Osman F, Kaempfer R. 2-Aminopurine selectively inhibits splicing of tumor necrosis factor alpha mRNA. *Mol Cell Biol* 1996; **16**:2814-2822.
- 43 Guerrier-Takada C, Eder PS, Gopalan V, Altman S. Purification and characterization of Rpp25, an RNA-binding protein subunit of human ribonuclease P. *RNA* 2002; **8**:290-295.
- 44 Circle DA, Neel OD, Robertson HD, Clarke PA, Mathews MB. Surprising specificity of PKR binding to delta agent genomic RNA. *RNA* 1997; **3**:438-448.
- 45 Ben-Asouli Y, Banai Y, Hauser H, Kaempfer R. Recognition of 5'-terminal TAR structure in human immunodeficiency virus-1 mRNA by eukaryotic translation initiation factor 2. *Nucleic Acids Res* 2000; **28**:1011-1018.

(**Supplementary information** is linked to the online version of the paper on the *Cell Research* website.)