

## CORRIGENDUM

# Dietary glycemic index, dietary glycemic load and mortality among men with established cardiovascular disease

EB Levitan, MA Mittleman and A Wolk

*European Journal of Clinical Nutrition* (2010) **64**, 439; doi:10.1038/ejcn.2010.11

**Correction to:** *European Journal of Clinical Nutrition* (2009) **63**, 552–557; doi:10.1038/sj.ejcn.1602970

The authors apologize for any inconvenience caused.

Since the publication of this paper, the authors have noticed an error in Table 1. The correct table is shown below.

**Table 1** Age-standardized characteristics of 4617 middle-aged men with established cardiovascular disease by quartile of dietary glycemic load (means or percentages)

Characteristic	Dietary glycemic index <sup>a</sup>					Dietary glycemic load <sup>a</sup>				
	Quartile 1	Quartile 2	Quartile 3	Quartile 4	P-value trend	Quartile 1	Quartile 2	Quartile 3	Quartile 4	P-value trend
Age (years)	67.5	67.1	67.4	67.6	0.59	67.1	67.2	67.6	67.8	0.02
Body mass index (kg/m <sup>2</sup> )	26.5	26.1	26.2	26.1	0.009	26.4	26.3	26.2	26.1	0.04
Physical activity (min per day)	60.6	60.7	57.8	54.2	<0.001	56.4	58.8	59.7	58.4	0.31
History of hypertension (%)	49.9	48.5	45.2	45.8	0.08	50.6	46.7	47.4	44.8	0.04
History of diabetes (%)	17.2	14.3	14.9	12.9	0.04	17.1	14.7	14.1	13.3	0.07
Family history of myocardial infarction at < 60 years (%)	21.5	21.8	20.0	18.9	0.26	20.5	20.4	20.9	20.5	0.99
Aspirin use (%)	65.6	67.5	64.9	63.6	0.25	62.7	67.3	65.0	66.6	0.09
Cigarette smoking (%)					0.008					0.003
Never	24.7	30.8	29.8	27.9		22.8	28.1	30.6	31.8	
Past	52.1	48.6	46.5	44.0		52.6	46.2	46.3	46.2	
Current	23.2	20.7	23.7	28.1		24.7	25.7	23.1	22.1	
Total energy (kJ per day)	10 174	10 213	10 241	9 952	0.14	10 356	10 132	10 134	9 957	0.004
Saturated fat (g per day) <sup>b</sup>	33.4	31.3	31.8	31.8	<0.001	39.0	33.5	30.4	25.4	<0.001
Monounsaturated fat (g per day) <sup>b</sup>	24.4	24.2	24.5	24.7	0.055	27.7	25.3	23.8	20.9	<0.001
Polyunsaturated fat (g per day) <sup>b</sup>	9.6	9.8	9.8	9.7	0.20	10.2	9.9	9.7	9.1	<0.001
Carbohydrate intake (g per day) <sup>b</sup>	267	282	287	294	<0.001	242	273	293	323	<0.001
Protein intake (g per day) <sup>b</sup>	95.3	89.0	85.9	82.0	<0.001	97.4	90.3	85.8	78.6	<0.001
Alcohol intake (g per day)	12.8	10.1	7.9	6.1	<0.001	13.1	10.3	7.9	5.5	<0.001
Cereal fiber intake (g per day) <sup>b</sup>	15.2	17.6	18.4	18.9	<0.001	14.3	17.2	18.5	20.1	<0.001
Dietary glycemic index <sup>a</sup>	71.9	76.6	79.4	83.4	—	74.0	77.1	78.9	81.3	<0.001
Dietary glycemic load <sup>a,b</sup>	192	216	228	245	<0.001	179	210	231	263	—

<sup>a</sup>White bread used as the reference food.

<sup>b</sup>Energy adjusted using the residuals method.