CORRIGENDUM

A dietary survey to determine if patients with coeliac disease are meeting current healthy eating guidelines and how their diet compares to that of the British general population

L Kinsey, ST Burden and E Bannerman

European Journal of Clinical Nutrition (2011) **65**, 283; doi:10.1038/ejcn.2010.256

Correction to: *European Journal of Clinical Nutrition* (2007) **62**, 1333–1342; doi:10.1038/sj.ejcn.1602856

Since the publication of this paper, the authors have noticed an error in the headings of Table 3 (*Males* and *Females*). The correct table is shown here.

Table 3 Mean intake of nutrients (s.d.) in patients with coeliac disease compared to the general population

Daily intakes	Females		Males	
	Coeliacs	NDNS	Coeliacs	NDNS
19–64 years ^a	(n = 22)	(n = 891)	(n = 6)	(n = 833)
Energy (kcal)	1754 ± 403	1632 ± 418	2039 ± 124	2313 ± 582*
Energy (kJ)	7338 ± 1687	6870 ± 1758	8530 ± 518	9720 ± 2446*
Protein (g)	83 ± 24	64 ± 17*	77 ± 10	88 ± 33*
Fat (g)	64 ± 20	61.4 ± 21.7	69 ± 11	87 ± 28.2*
CHO (g)	219 ± 65	203 ± 59	262 ± 31	275 ± 79
NSP (g)	12 ± 5	12.6 ± 5.0	16 ± 5	15.2 ± 6.0
Folate (μg)	251 ± 120	251 ± 90	258 ± 68	344 ± 127*
Vitamin B12 (μg)	6.9 ± 10.0	4.8 ± 2.7	5.1 ± 1.5	6.5 ± 4.7
Vitamin D (μg)	2.7 ± 2.07	3.7 ± 2.3*	3.1 ± 3	2.8 ± 2.1
Iron (mg)	9.9 ± 3.3	10.0 ± 3.7	11.7 ± 5.5	10.0 ± 3.7
Calcium (mg)	807 ± 283	777 ± 269	1076 ± 207	1007 ± 411
> 65 years ^b	(n = 13)	(n = 891)	(n = 6)	(n = 632)
Energy (kcal)	1918 ± 503	1909 ± 465	1457 ± 331	1374 ± 337
Energy (kJ)	8346 ± 1919	8020 ± 195	6096 ± 1385	5970 ± 1410
Protein (g)	83 ± 21	72 ± 17	60 ± 15	56 ± 13
Fat (g)	56 ± 28	75 ± 24*	49 ± 12	58 ± 18
CHO (g)	277 ± 88	232 ± 64	201 ± 60	175 ± 47
NSP (g)	13.4 ± 3.3	13.5 ± 5.8	11.8 ± 4.7	11 ± 4.7
Folate (μg)	339 ± 189	270 ± 95	214 ± 61	207 ± 75
Vitamin B12 (μg)	4.1 ± 2.1	6.1 ± 6.2*	2.7 ± 1.2	4.5 ± 4.4
Vitamin D (μg)	2.7 ± 1.7	4.07 ± 3.2*	1.9 ± 1.4	2.9 ± 2.4
Iron (mg)	11.9 ± 4.6	11.0 ± 3.6	7.5 ± 3.3	8.6 ± 2.9
Calcium (mg)	1074 ± 430	836 ± 285	798 ± 365	690 ± 246

Abbreviations: CD, coeliac disease; CHO, carbohydrate; NSP, non-starch polysaccharide.

* $P < 0.05$; one sample *t*-test. Dietary intakes of CD patients are significantly different from dietary intakes of the general UK population.

^aNDNS, 2003a.

^bNDNS, 1998.

In addition, there are five instances where Table 3 is incorrectly cited. These are all in the section 'Results compared to the DRVs' and are listed as follows:

Page 3, right-hand column, line 3

Page 4, left-hand column, line 4

Page 4, left-hand column, line 2

Page 4, right-hand column, line 10

Page 4, right-hand column, line 31

The authors would like to apologize for this error.