

CORRIGENDUM

Predictors of change in weight and waist circumference:
15-year longitudinal study in Australian adults

S Arabshahi, PH Lahmann, GM Williams and JC van der Pols

European Journal of Clinical Nutrition (2015) 69, 536; doi:10.1038/ejcn.2015.13

Correction to: *European Journal of Clinical Nutrition* 68, 309–315;
doi:10.1038/ejcn.2013.260; published online 8 January 2014

Since the publication of this article, the authors have noticed that the figures in Table 1 were incorrect due to a layout error. Below is the corrected table.

The authors apologize for any inconvenience caused.

Table 1. Weight and waist circumference by birth cohort, Nambour Study, 1992 and 2007

Age group (year)	Weight					Waist circumference				
	1992		2007		P-value ^a	1992		2007		P-value ^a
	n	Weight (kg) ^b	n	Weight (kg) ^b		n	Waist circumference (cm) ^b	n	Waist circumference (cm) ^b	
<i>Men</i>										
25–34	81	82.8 (12.2)	—	—		81	93.4 (10.7)	—	—	
35–44	105	80.8 (11.5)	18	85.5 (11.1)	0.1	104	92.7 (8.5)	18	91.2 (8.8)	0.5
45–54	133	84.6 (14.1)	72	91.8 (16.1)	0.001	131	98.1 (11.1)	72	98.3 (12.4)	0.9
55–64	102	80.0 (9.9)	83	86.8 (11.5)	0.0001	96	97.3 (9.4)	83	98.2 (9.3)	0.5
65–74	112	78.2 (12.5)	71	84.3 (15.3)	0.004	108	98.7 (10.2)	71	98.0 (12.7)	0.7
≥ 75	7	76.9 (13.6)	58	75.4 (10.8)	0.7	7	100.5 (10.8)	58	95.9 (10.4)	0.3
<i>Women</i>										
25–34	83	64.6 (12.2)	—	—		81	77.8 (10.3)	—	—	
35–44	167	67.0 (12.7)	30	68.6 (14.4)	0.5	166	79.6 (10.2)	30	78.7 (12.5)	0.7
45–54	198	69.4 (14.0)	90	75.1 (15.3)	0.003	192	83.2 (11.3)	90	84.7 (11.7)	0.3
55–64	159	68.7 (12.2)	142	74.5 (14.7)	0.003	154	85.7 (10.1)	142	87.3 (12.2)	0.2
65–74	92	67.0 (11.4)	91	70.8 (14.5)	0.05	92	88.0 (10.4)	91	86.3 (12.8)	0.3
≥ 75	9	70.1 (13.0)	57	68.6 (11.6)	0.7	9	90.2 (14.9)	57	87.7 (9.8)	0.6

^aFrom two-sample *t*-test. ^bMean (s.d.), all such values.