

ARTICLE

Copy number detection in discordant monozygotic twins of Congenital Diaphragmatic Hernia (CDH) and Esophageal Atresia (EA) cohorts

Danielle Veenma^{*,1,2}, Erwin Brosens^{1,2}, Elisabeth de Jong^{1,2}, Cees van de Ven¹, Connie Meeussen¹, Titia Cohen-Overbeek³, Marjan Boter², Hubertus Eussen², Hannie Douben², Dick Tibboel¹ and Annelies de Klein²

The occurrence of phenotypic differences between monozygotic (MZ) twins is commonly attributed to environmental factors, assuming that MZ twins have a complete identical genetic make-up. Yet, recently several lines of evidence showed that both genetic and epigenetic factors could have a role in phenotypic discordance after all. A high occurrence of copy number variation (CNV) differences was observed within MZ twin pairs discordant for Parkinson's disease, thereby stressing on the importance of post-zygotic mutations as disease-predisposing events. In this study, the prevalence of discrepant CNVs was analyzed in discordant MZ twins of the Esophageal Atresia (EA) and Congenital Diaphragmatic Hernia (CDH) cohort in the Netherlands. Blood-derived DNA from 11 pairs (7 EA and 4 CDH) was screened using high-resolution SNP arrays. Results showed an identical copy number profile in each twin pair. Mosaic chromosome gain or losses could not be detected either with a detection threshold of 20%. Some of the germ-line structural events demonstrated in five out of eleven twin pairs could function as a susceptible genetic background. For example, the 177-Kb loss of chromosome 10q26 in CDH pair-3 harbors the *TCF7L2* gene (*Tcf4* protein), which is implicated in the regulation of muscle fiber type development and maturation. In conclusion, discrepant CNVs are not a common cause of twin discordancy in these investigated congenital anomaly cohorts.

European Journal of Human Genetics (2012) 20, 298–304; doi:10.1038/ejhg.2011.194; published online 9 November 2011

Keywords: monozygotic twins; copy number variation; congenital anomaly; phenotypic discordance; congenital diaphragmatic hernia; esophageal atresia

INTRODUCTION

Monozygotic (MZ) twin comparisons have been used for many decades to specify contributions of both nature (heredity) and nurture (environment).¹ Normally, the study design is based on the presumption that MZ twins come from one fertilized egg, and therefore have complete identical genetic make-ups. Yet, recently several lines of evidence suggested that genetic and epigenetic factors could have a role in MZ phenotypic variances after all.^{2–6} Using a BAC array platform, Bruder *et al*⁶ demonstrated that discordance in their MZ Parkinson's disease (PD) twin cohort of nine individuals could be the result of copy number variation (CNV) differences. However, Baranzini *et al*⁷ could not reproduce this high intra-twin pair variability of structural variants using both array and next-generation sequencing in three twin pairs discordant for Multiple Sclerosis.

We investigated whether discrepant CNVs could cause discordance in MZ twin pairs of the Dutch Esophageal Atresia (EA (MIM 189960)) and Congenital Diaphragmatic Hernia (CDH (MIM 142340)) cohort. Blood-derived DNA from 11 (7 EA and 4 CDH) pairs of MZ twins was screened using high-resolution SNP arrays.

EA generally presents at birth with a defective formation of the esophagus with or without a fistulous tract to the trachea. Although

not lethal in most cases, long-term morbidity has a significant role in these patients. CDH is a more severe birth defect characterized by defective formation of the diaphragm, lung hypoplasia and pulmonary hypertension. Despite medical advances mortality for isolated cases is 20% and for none-isolated cases up to 60%. Both EA and CDH are presumed to have a multifactorial etiology and the identification of chromosomal aberrations and knockout animal models provide strong evidence for a genetic component.⁸ In contrast, both anomalies present with low twin concordance rates, 10.7% and 15.6% for EA and CDH, respectively, and sibling recurrence rates are low (1–2%) as well. Shaw-Smith⁹ already pointed out that the incidence of twinning in EA is 2.6 times higher than statistically expected. In all, 206 pairs are described in the literature up until now; however, information on zygosity is less thorough.^{9–16} Orford *et al*¹⁵ stated that at least 80% of reported EA twins are same-sex pairs. In total, 22 out of these 206 twin pairs are concordant for the EA phenotype. In the literature, 77 twin pairs have been described for CDH of which 53 were recognized as MZ.^{17–20} Twelve pairs were concordant for the CDH phenotype.

The rationale of this study was to investigate whether CNVs in the affected twin sibling could account for phenotypic discordance of either EA or CDH MZ twin siblings. Although results showed no such

¹Department of Paediatric Surgery, Rotterdam, The Netherlands; ²Department of Clinical Genetics, Rotterdam, The Netherlands; ³Department of Obstetrics and Gynaecology, Erasmus MC-Sophia, Rotterdam, The Netherlands

*Correspondence: Dr D Veenma, Department of Paediatric Surgery, Dr. Molewaterplein 50, Rm SK-3286, PO Box 2040, 3000 CA Rotterdam, The Netherlands.

Tel: +31 10 7043492; Fax: +31 10 7044736; E-mail: d.veenma@erasmusmc.nl

Received 30 May 2011; revised 9 September 2011; accepted 15 September 2011; published online 9 November 2011

Table 1a Clinical features EA cohort

Pair EA	GA (weeks)	Birth order		Obstetric history	EA	Fistel	Type of additional anomalies
		(Patient)	(Healthy twin)				
EA1	37.3	1	2	Breech presentation	+	+	Dysmorphic Auricular tags, Cleft uvula Abnormal dermatoglyphics Heart ASD Lung Lung hypoplasia right Neurologic/skeletal Scoliosis Fusion of vertebrae Hemivertebrae IUGR
EA2	36	2	1	Breech presentation Maternal medication: Sintrom	+	+	Heart VSD Lung Lunghypoplasia
EA3	NA	NA	NA	NA	+	+	Heart Cardiac situs Dextrocardia healthy twin
EA4	NA	NA	NA	NA	+	—	—
EA5	33.5	1	2	Breech presentation Fever durante partu	+	+	Dysmorphic Triangular face, deep set eyes, Palpebral fissures slant down Small mandible Thin fingers, hypoplastic thumbs Proximal placement of thumb Hypoplastic or absent radii Sacral hemangioma healthy twin
EA6	34.4	NA	NA	Polyhydramnion Maternal medication: Corticosteroids	+	+	Heart VSD Tricuspid incompetence
EA7	NA	NA	NA	—	+	+	—

Table 1b Clinical features CDH cohort

Pair CDH	GA (weeks)	Birth order		Obstetric history	CDH	Type of additional anomalies
		(Patient)	(Healthy twin)			
CDH1	35.3	2	1	Sectio Caesarea Breech presentation	Left	—
CDH2	33.4	1	2	Sectio Caesarea >24 h ruptured membrane	Left	—
CDH3	38.5	2	1	—	Right	Urogenital Inguinal hernia Hydrocele testis
CDH4	34.1	2	1	Sectio Caesarea	Left	Dysmorphic Small mandibula IUGR

Abbreviations: ASD, Atrial Septal defect; CDH, Congenital Diaphragmatic Hernia; EA, Esophageal Atresia; GA, gestational age; IUGR, Intra Uterine Growth Retardation; VSD, Ventricular Septal Defect. Unfortunately, for a few EA subjects detailed clinical data are unavailable.

proof, germ-line structural events were detected and these could represent a susceptible genetic background as seen in other genetic anomalies. Results are discussed in the context of earlier MZ twin reports.

MATERIALS AND METHODS

Ethics statement

Research involving human participants has been approved by the Medical Ethical Committee (METC) at Erasmus MC, which specifically approved for blood

withdrawal of both twins and their parents. Informed consent forms were obtained for the index case and his/her parents at once and for the healthy twin separately.

Patients

The 11 affected twin samples were collected from the congenital anomaly cohort in Rotterdam (Erasmus MC Sophia's Hospital, The Netherlands) in which 541 EA and 626 CDH patients are currently registered. Of these, 22 CDH patients (14 dizygotic, 5 MZ, and 3 not tested) and 35 EA patients (6 dizygotic, 9 MZ, and 20 not tested) were the result of a twin pregnancy. Included were those twin samples with a written parental informed consent, quality material of both siblings and confirmed monozygosity by STR profiling (AmpFISTR identifier PCR amplification kit, Applied Biosystems, Foster City, CA, USA). Another exclusion criterion was the identification of a genetic abnormality, most commonly an aneuploidy.

DNA isolation

Automated DNA extraction from peripheral blood (or skin fibroblasts in case of two affected CDH twins) was performed using local standard protocols. DNA quality and concentration were checked with the Quant-iT PicoGreen dsDNA Kit (Invitrogen Corporation, Carlsbad, CA, USA).

Whole-genome high-resolution SNP array

SNP analysis was carried out using the Illumina HumanCytoSNP-12 version 2.2. (Illumina, San Diego, CA, USA). This chip includes 220 000 of the most informative SNPs markers with a median physical distance of 6.2 Kb. DNA samples were processed according to the manufacturer's protocol. The call rate of this array batch was above 0.98, except for one sample.

SNP array analysis

Data for each bead chip were self-normalized in Genomestudio GT (Illumina) using information contained within the array. Copy number estimates for each individual sample were determined by comparison with a common reference set of 200 samples from the HapMAP project (www.hapmap.org/downloads/raw_data) supplied by Illumina (Manifest files) and visualized in the Nexus software program (version five, Biodiscovery, El Segundo, CA, USA) as log₂ ratios. Analysis settings were as follow: both SNP-FASST and SNP-Rank segmentation methods were executed independently with significance thresholds ranging from 1×10^{-4} to 1×10^{-6} and log-ratio thresholds of 0.18 and -0.18 for duplication and deletions, respectively. The max contiguous probe spacing was 1000 Kb and the minimum number of probes per segment was set to three, limiting CNV detection to sizes > 18.6 Kb. Subsequently, only CNVs > 50 Kb were validated. Paired analysis for deletions and duplications was performed in each affected twin versus its healthy co-twin. As described recently, high-resolution (SNP) arrays are suitable for detection of both germ-line and mosaic CNVs.^{21–25} Mosaic copy number aberrations are hallmarked by a concomitant change of log₂ intensity signal and a shift in b-allele frequency. The detection limit (sensitivity) of the Nexus SNP-FASST algorithm for mosaic CNVs is 20% using a heterozygous imbalance threshold of 0.45.²² To review functionality of each putative CNV at once, occurrence frequencies in a qualified normal pediatric cohort of 2026 individuals²⁶ (CHOP; <http://cnv.chop.edu/>) and in the DGV (<http://www.tcag.ca>) were uploaded in the Nexus program as well. Since these populations display various ethnic backgrounds, comparison with an in-house normal reference set was performed as well. Additionally, possible intra-twin pair genotype differences (with respect to all SNP markers presented on the array) were evaluated in Genomestudio GT using the paired analysis settings.

Validation using fluorescent *in-situ* hybridization and relative-quantitative PCR analysis

Confirmation of each CNV with quantitative real-time PCR and/or FISH was executed in the twin siblings and their parents according to local standard protocols with minor modifications.^{22,27} For FISH, BAC clones were selected from the UCSC genome browser (<http://genome.ucsc.edu/>), purchased at BACPAC resources center (Oakland, CA, USA) and labelled (Random Prime labelling system; Invitrogen Corporation) with Bio-16-dUTP or Dig-11-dUTP (Roche Applied Science, Indianapolis, IN, USA). After validation on control metaphases, the chromosome 22 BAC clones RP11-62K15 and RP1-66M5 were used for confirmation in EA pair-I.

Table 2 Inherited CNVs detected in MZ twins of the Rotterdam Congenital anomaly cohort

EA pair	Chromosomal location (bp)	CNV type	Length (bp)	Gene symbols	Discordant twin	Normal pediatric cohort #	Validation
1	chr22:46,163,818-46,870,578	CN loss	666 192	FLJ46257 (hypothetical)	Yes	Yes (0/2026, 0/370, overlap 25 DGV)	FISH, inherited maternally
2	chr7:7,5,420,580-75,471,147	CN loss	50 567	POR, STYX11, TMEM120A	Yes	Yes (0/2026, 0/370, overlap 2 DGV)	q-PCR, inherited (both parents heterozygous deletion)
3	chr6:162,638,827-162,840,229	CN gain	201 402	PARK2	Yes	Yes 5/2026	q-PCR, inherited maternally
	chr17:41,522,684-41,646,903	CN gain	124 219	KIAA1267	Yes	Yes: 69/2026	Copy number polymorphism
	chr18:1,895,191-1,960,898	CN loss	65 707	—	Yes	Yes: 26/2026	Copy number polymorphism
4	chr14:19,283,777-19,479,370	CN gain	195 593	OR4K1, OR4K2, OR4K5, etc	Yes	Yes: 35/2026	Copy number polymorphism
5	chr12:7,892,014-8,027,862	CN gain	135 848	SLC2A14, SLC2A3	Yes	Yes: 39/2026	Copy number polymorphism
6	chr12:31,146,084-31,303,651	CN gain	157 567	DDX11	Yes	Yes: 20/370	Copy number polymorphism
	chr2:89,728,406-89,885,025	CN gain	156 619	—	Yes	Yes: 14/2026	Copy number polymorphism
	chr8:2,322,561-2,577,455	CN gain	254 894	—	Yes	Yes: 32/2026	Copy number polymorphism
7	—	—	—	—	—	—	—
CDH pair	Chromosomal location (bp)	CNV (type)	Length (bp)	Gene symbols	Discordant twin	Normal cohort #	Validation
1	—	—	—	—	—	—	—
2	—	—	—	—	—	—	—
3	chr10:114,660,279-114,838,014	CN loss	177 735	TCF7L22	Yes	No	q-PCR, inherited maternally
4	chr4:86,595,913-87,070,050	CN loss	474 137	ARHGAP24 ^{23,24}	Yes	Yes (0/2026, 0/370, overlap 27 DGV)	q-PCR, inherited maternally
	chr4:57,743,262-57,795,280	CN gain	52 018	—	Yes	Yes (2/2026, 0/370, overlap 15 DGV)	Copy number polymorphism

Abbreviations: CN, copy number; bp, basepairs; EA, Esophageal Atresia; CDH, Congenital Diaphragmatic Hernia. # Normal Cohort; CHOP (<http://cnv.chop.edu/>) and DGV (<http://tcag.ca>). & Decipher (<http://decipher.sanger.ac.uk/application/>). Percentage of overlap with CNVs in control databases is designated if not 100% aligned.

Primer pairs for quantitative real-time PCR were designed from unique sequences within the minimal deleted or duplicated regions of each copy number change using Primer Express software v2.0 (Applied Biosystems, Carlsbad, CA, USA). The nucleotide BLAST algorithm at NCBI (<http://www.ncbi.nlm.nih.gov/BLAST/>) was used to confirm that each PCR amplification product was unique. Quantitative PCR analyses were performed using a LightCycler 1.5 instrument in combination with LightCycler FastStart DNA Master SYBR Green I kits (Roche Molecular Diagnostics, Indianapolis, IN, USA). Experiments were designed with a region of the *C14ORF145* gene serving as a control locus as previously described.²⁷

RESULTS

Clinical characterization and monozygosity screening of twin pairs

Clinical features of each twin pair are summarized in Table 1. Briefly, 7 out of 11 pairs were discordant for the phenotype of EA (Table 1a) and 4 out of 11 for CDH (Table 1b). Four out of eleven EA-affected

patients harbored (major) additional anomalies. Considering CDH, there is a variable expression of left and right CDH with all persons (as expected) featuring lung hypoplasia. We are dealing with an isolated CDH cohort since most anomalies in pairs 3 and 4 are minor. Finally, zygosity status of each twin pair was confirmed (data not shown) by STR profiling using the commercially available STR identifier kits of Applied Biosystems.

Paired CNV analysis of discordant MZ twins

Results of the paired CNV analysis of each MZ twin couple are summarized in Table 2 showing no evidence of pathogenic CNV discordance in both congenital anomaly cohorts. In order to detect mosaic (somatic) aberrations, specific attention was paid to b-allele frequency changes as well. In the EA cohort, a total of 10 germ-line CNVs were identified. Seven concerned common copy number




Figure 1 SNP and fluorescent *in-situ* hybridization results of inherited chromosome 22 CNV in EA pair-I. Nexus results (Top) of the 666-Kb deletion on chromosome 22q13.3 in both individuals of EA pair-I showing a clear drop in log₂ intensity signal validated by FISH (Bottom) on metaphase chromosomes of the affected EA twin-1. Probes are control: RP11-62K15 (green) and target: RP11-66M5 (red). Parental analysis (results not shown) demonstrated that this genomic event is inherited from the mother and therefore less likely pathogenic. In addition, no gene is allocated to this region neither are any miRNA transcripts hampering the identification of functional elements in this region as well.

polymorphisms defined by the occurrence of the CNV in at least five individuals of qualified normal pediatric cohorts in the literature. The remaining three events were present in both the twin and at least one healthy parent, and are therefore less likely to be pathogenic as well. For example, the 666-Kb sized chromosome 22q deletion in EA pair-1 (Figure 1) was found both in the healthy twin and his mother and partly overlaps with CNVs cataloged in control cohorts.

Existence of inherited CNVs was detected in the CDH cohort as well. A total of three CNVs were distinguished of which two are not prevalent in normal cohorts. All three events were present in the healthy twin as well. Figure 2 represents the 177-Kb loss of chromosome 10q26 in CDH pair-3 and harbors the *TCF7L2* gene (*Tcf4* protein), which is mainly known for its involvement in blood glucose homeostasis as a result of Wnt signalling changes.

Not ruled out in this study are the presence of balanced genomic alterations and small (<50 Kb) or very-low mosaic (<20%) chromo-

some aberrations beyond the detection level of our experimental approach.

SNP genotype analysis MZ twin cohorts

SNP genotype differences between the affected and unaffected twin siblings were evaluated for each SNP on the Illumina bead chip. After removal of less accurately called SNPs, genotyping analysis showed concordance for almost all SNPs ($n=299671$) within each MZ pair. A total of five SNPs in three EA pairs were dissimilar and three SNPs in two CDH pairs (Table 3). CDH pair-3 showed discrepancy for 99 SNPs, which could be attributed to less overall genotyping accuracy and therefore was not analyzed further. Until now, only rs2824374 (which is closely linked to the *CXADR* gene) could be associated with embryonic (mal) development, however, literature only reports on effects to the kidneys and cochlea.^{28,29} None of the other identified discordant intra-twin SNP loci are directly linked to a phenotype.




Figure 2 SNP and RT-PCR results of inherited chromosome 10-CNV in CDH pair-3. Nexus result (TOP) of the chromosome 10q26 deletion event in CDH pair-3 showing a clear drop in log₂ intensity signal which was confirmed by relative q-PCR (Bottom) in the affected proband, the unaffected twin and the mother.

Table 3 Discordant SNPs in MZ twin pairs of the Rotterdam Congenital anomaly cohort

	Discordant SNPs	Chromosomal location (dbSNP build 130) (bp)	Gene symbols
<i>Pair EA</i>			
1	rs11573502	chrX:24888693-24889193	POLA1
2	—	—	—
3	—	—	—
4	rs10125846 rs1744767 rs438895	chr9:2821641-2822141 chr20:35200380-35200880 chr1:8260692-8261192	KIAA0020 LOC140699 isoform 3
5	—	—	—
6	—	—	—
7	rs1576026	chr9:25,453,089-25,453,589	—
<i>Pair CDH</i>			
1	rs17730982 rs2824374	chr8:134851934-134852434 chr21:17879252-17879752	CXADR; possible involvement Kidney and cochlear development ^{25,26}
2	—	—	—
4	rs6571064	chr6:103400463-103400963	—
3	Not evaluated; SNP quality rate <0.95	—	—

Abbreviations: bp, basepairs; EA, Esophageal Atresia; CDH, Congenital Diaphragmatic Hernia.

DISCUSSION

A high occurrence of copy number variants that differed between siblings discordant for PD was recently suggested.⁶ However, intra-twin pair variability for germ-line CNVs could not be detected in our subset of EA and CDH MZ twins. Within the limitations of the used experimental approach, structural variants in mosaic form (above 20%) could neither be demonstrated. Application of next-generation sequencing methods will allow for an easier and more sensitive calling of the smallest mosaic aberrations in the near future and will add up to the (scarce) data generated recently on this topic by some other groups.^{30–33}

Various causes could account for the discrepancy in CNV findings between our congenital anomaly twin cohort and the Parkinson's cohort. First of all, an age factor: the rather high prevalence of mosaic CNVs in PD twins could have been generated during lifetime. This was suggested by a small study of the group of Dumanski *et al*,^{34,35} who identified mosaic aberrations in a wide range of tissues of three phenotypically normal individuals. This hypothesis would imply that age-accumulated (tissue-specific) CNV events could have a role in diseases developing symptoms later in life. Consequently, they are expected to contribute less to congenital disorders. Second, differences in CNV prevalence between our study and the Parkinson's study could be based on methodological differences such as choice of platform. Although Bruder *et al* presented confirmative evidence for a few of their CNVs using a different platform, detailed confirmation of most CNVs was lacking. On the other hand, structural DNA variation might have a minor role in EA and CDH pathophysiology, suggesting that in these congenital cohorts the focus should be widened on environmental and epigenetic factors. Two recent studies^{2,7} revealed a (significant) proportion of epigenetic variability between MZ twins in investigated tissues. However, in the Multiple Sclerosis twin cohort study these changes could not account for disease discordance. A similar study for EA, CDH, or other congenital anomalies is difficult to perform, since the target tissues cannot be obtained from the healthy co-twin for obvious reasons. Structural variations restricted to the affected esophagus and diaphragm tissue could represent another cause for twin discordance, yet was not excluded in this MZ cohort due to unavailability of the affected material.

Finally, although our results showed no proof for CNV contribution to phenotypic MZ discordance, germ-line structural events were detected in both cohorts and these events could represent a so-called susceptible genetic background. In five out of eleven twin pairs, germ-line CNVs were identified. These were rarely detected in a specific pediatric normal population²⁶ and/or our in-house control cohort and could therefore represent an increased susceptibility to congenital anomalies by means of a dosage responsive or position effect. For example, the 177-Kb loss of chromosome 10q26 in CDH pair-3 might be of functional importance. A recent report demonstrated *Tcf4* (alias *Tcf7l2*) expression in connective tissue fibroblasts during development and suggested its role in the regulation of muscle fiber type development and maturation.³⁶ Additionally, certain polymorphisms and mutations in *TCF7L2* are linked to an increased risk of type 2 diabetes.³⁷ This implies that loss of one functional *TCF7L2* allele might be associated with (super) normal glucose tolerance. Indeed, we observed evidence of increased serum glucose (a glucose of 12.2 mmol/l was identified within 24 h postnatal) in the affected individual of twin pair-3. However, also one normal glucose level (glucose 3.6 mmol/l) was determined within the same time window and since this patient was critically ill and died shortly thereafter no absolute conclusions can be drawn from these results. The healthy twin had an unremarkable medical record so far. Similarly, the haploinsufficient *ARHGAP24* gene in CDH pair-4 (encoding a vascular, cell-specific GTPase-activating protein) could confer genetic susceptibility for CDH by means of its function in modulating angiogenesis and through its interaction with filamin-A.^{38,39} Girirajan *et al*⁴⁰ recently demonstrated that a second hit may elicit a severe phenotype in offspring of healthy CNV carriers. Hypothetically, this second hit can constitute another CNV in the same or associated disease pathway as well as a pathogenic SNP. These results underline the importance of archiving all genomic events (also those with a 'benign' nature at first sight) in a freely accessible database such as initiated by the ISCA consortium (<https://www.iscaconsortium.org>). Detailed and unbiased phenotyping is crucial for the understanding of the more complex genotype–phenotype correlations as well.

In summary, we investigated whether the existence of discrepant CNVs could be causal to the phenotypic discordance in MZ twin pairs

of the EA and CDH cohort in Rotterdam and found no such proof. Prospective collection of DNA material in various MZ twin cohorts is warranted to evaluate the possibility of such genetic factors contributing to human phenotypic variability in general and to twin discordance specific. We feel that the use of high-resolution SNP arrays and sequencing-based methods are more suitable in these designs than BAC arrays. Finally, phenotypic correlations can only be made after proper analysis in normal cohorts as well.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We would like to thank all patients, specifically all twins and their parents for making this study possible. Also, we thank Tom de Vries-Lentsch for his help in preparing the figures. This study was funded by the Sophia's Children's Hospital Foundation for Medical Research, Rotterdam, The Netherlands (SSWO 493 and 551). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

- 1 Hall JG: Twinning. *Lancet* 2003; **362**: 735–743.
- 2 Kaminsky ZA, Tang T, Wang SC *et al*: DNA methylation profiles in monozygotic and dizygotic twins. *Nat Genet* 2009; **41**: 240–245.
- 3 Kimani JW, Shi M, Daack-Hirsch S *et al*: X-chromosome inactivation patterns in monozygotic twins and sib pairs discordant for nonsyndromic cleft lip and/or palate. *Am J Med Genet* 2007; **143A**: 3267–3272.
- 4 Machin GA: Some causes of genotypic and phenotypic discordance in monozygotic twin pairs. *Am J Med Genet* 1996; **61**: 216–228.
- 5 Streit A, Sommer RJ: Genetics: random expression goes binary. *Nature* 2010; **463**: 891–892.
- 6 Bruder CE, Piotrowski A, Gijbbers AA *et al*: Phenotypically concordant and discordant monozygotic twins display different DNA copy-number-variation profiles. *Am J Human Genet* 2008; **82**: 763–771.
- 7 Baranzini SE, Mudge J, van Velkinburgh JC *et al*: Genome, epigenome and RNA sequences of monozygotic twins discordant for multiple sclerosis. *Nature* 2010; **464**: 1351–1356.
- 8 Holder AM, Klaassens M, Tibboel D, de Klein A, Lee B, Scott DA: Genetic factors in congenital diaphragmatic hernia. *Am J Human Genet* 2007; **80**: 825–845.
- 9 Shaw-Smith C: Oesophageal atresia, tracheo-oesophageal fistula, and the VACTERL association: review of genetics and epidemiology. *J Med Genet* 2006; **43**: 545–554.
- 10 Sunagawa S, Kikuchi A, Yoshida S *et al*: Dichorionic twin fetuses with VACTERL association. *J Obstet Gynaecol Res* 2007; **33**: 570–573.
- 11 Depaepae A, Dolk H, Lechat MF: The epidemiology of tracheo-oesophageal fistula and oesophageal atresia in Europe. EUROCAT Working Group. *Arch Dis Child* 1993; **68**: 743–748.
- 12 Farquhar J, Carachi R, Raine PA: Twins with oesophageal atresia and the CHARGE association. *Eur J Pediatr Surg* 2002; **12**: 56–58.
- 13 Khan RA, Narashimhan KL: Discordance in twins for esophageal atresia with tracheo-oesophageal fistula. *Indian J Pediatr* 2009; **76**: 334–335.
- 14 Ohno K, Nakamura T, Azuma T *et al*: Esophageal atresia with tracheoesophageal fistula in both members of monozygotic twins. *Pediatr Surg Int* 2008; **24**: 1137–1139.
- 15 Orford J, Glasson M, Beasley S, Shi E, Myers N, Cass D: Oesophageal atresia in twins. *Pediatr Surg Int* 2000; **16**: 541–545.
- 16 Trobs RB, Knupffer M, Schutz A, Cernaianu G, Hirsch W, Regenthal R: Congenital oesophageal atresia discordant for tracheo-oesophageal fistula occurring in a set of dizygotic twins. *Eur J Pediatr Surg* 2006; **16**: 260–264.
- 17 Donahoe PK, Noonan KM, Lage K: Genetic tools and algorithms for gene discovery in major congenital anomalies. *Birth Defects Res* 2009; **85**: 6–12.
- 18 Lau ST, Kim SS, Ledbetter DJ, Healey PJ: Fraternal twins with Morgagni hernias. *J Pediatr Surg* 2005; **40**: 725–727.
- 19 Machado AP, Ramalho C, Portugal R *et al*: Concordance for bilateral congenital diaphragmatic hernia in a monozygotic dichorionic twin pair - first clinical report. *Fetal Diagn Ther* 2010; **27**: 106–109.
- 20 Pober BR, Lin A, Russell M *et al*: Infants with Bochdalek diaphragmatic hernia: sibling recurrence and monozygotic twin discordance in a hospital-based malformation surveillance program. *Am J Med Genet* 2005; **138A**: 81–88.
- 21 Conlin LK, Thiel BD, Bonnemann CG *et al*: Mechanisms of mosaicism, chimerism and uniparental disomy identified by single nucleotide polymorphism array analysis. *Hum Mol Genet* 2010; **19**: 1263–1275.
- 22 Veenma D, Beurskens N, Douben H *et al*: Comparable low-level mosaicism in affected and non affected tissue of a complex CDH patient. *PLoS One* 2010; **5**: e15348.
- 23 Shaffer LG, Kashork CD, Saleki R *et al*: Targeted genomic microarray analysis for identification of chromosome abnormalities in 1500 consecutive clinical cases. *J Pediatr* 2006; **149**: 98–102.
- 24 Ballif BC, Rorem EA, Sundin K *et al*: Detection of low-level mosaicism by array CGH in routine diagnostic specimens. *Am J Med Genet A* 2006; **140**: 2757–2767.
- 25 Xiang B, Li A, Valentin D, Nowak NJ, Zhao H, Li P: Analytical and clinical validity of whole-genome oligonucleotide array comparative genomic hybridization for pediatric patients with mental retardation and developmental delay. *Am J Med Genet A* 2008; **146A**: 1942–1954.
- 26 Shaikh TH, Gai X, Perin JC *et al*: High-resolution mapping and analysis of copy number variations in the human genome: a data resource for clinical and research applications. *Genome Res* 2009; **19**: 1682–1690.
- 27 Scott DA, Klaassens M, Holder AM *et al*: Genome-wide oligonucleotide-based array comparative genome hybridization analysis of non-isolated congenital diaphragmatic hernia. *Hum Mol Genet* 2007; **16**: 424–430.
- 28 Excoffon KJ, Avenarius MR, Hansen MR *et al*: The Coxsackievirus and Adenovirus Receptor: a new adhesion protein in cochlear development. *Hear Res* 2006; **215**: 1–9.
- 29 Raschperger E, Neve EP, Wernerson A, Hulthenby K, Pettersson RF, Majumdar A: The coxsackie and adenovirus receptor (CAR) is required for renal epithelial differentiation within the zebrafish pronephros. *Dev Biol* 2008; **313**: 455–464.
- 30 Razzaghi HR, Shahi MH, Forsberg LA *et al*: Somatic mosaicism for chromosome X and Y aneuploidies in monozygotic twins heterozygous for sickle cell disease mutation. *Am J Med Genet* 2010; **152A**: 2595–2598.
- 31 Hasler R, Begun A, Freitag-Wolf S *et al*: Genetic control of global gene expression levels in the intestinal mucosa: a human twin study. *Physiol Genomics* 2009; **38**: 73–79.
- 32 Notini AJ, Craig JM, White SJ: Copy number variation and mosaicism. *Cytogenet Genome Res* 2008; **123**: 270–277.
- 33 Yurov YB, Iourov IY, Vorsanova SG *et al*: Aneuploidy and confined chromosomal mosaicism in the developing human brain. *PLoS One* 2007; **2**: e558.
- 34 Piotrowski A, Bruder CE, Andersson R *et al*: Somatic mosaicism for copy number variation in differentiated human tissues. *Hum Mutat* 2008; **29**: 1118–1124.
- 35 De S.: Somatic mosaicism in healthy human tissues. *Trends Genet* 2011; **27**: 217–223.
- 36 Mathew SJ, Hansen JM, Merrell AJ *et al*: Connective tissue fibroblasts and Tcf4 regulate myogenesis. *Development* 2011; **138**: 371–384.
- 37 Hansson O, Zhou Y, Renström E *et al*: Molecular function of TCF7L2: Consequences of TCF7L2 splicing for molecular function and risk for type 2 diabetes. *Curr Diab Rep* 2010; **10** (6): 444–451.
- 38 Nakamura F, Heikkinen O, Pentikainen OT *et al*: Molecular basis of filamin A-FilGAP interaction and its impairment in congenital disorders associated with filamin A mutations. *PLoS One* 2009; **4**: e4928.
- 39 Su ZJ, Hahn CN, Goodall GJ *et al*: A vascular cell-restricted RhoGAP, p73RhoGAP, is a key regulator of angiogenesis. *Proc Natl Acad Sci USA* 2004; **101**: 12212–12217.
- 40 Girirajan S, Rosenfeld JA, Cooper GM *et al*: A recurrent 16p12.1 microdeletion supports a two-hit model for severe developmental delay. *Nat Genet* 2010; **42**: 203–209.