

SHORT REPORT

# Deletions in *PITX1* cause a spectrum of lower-limb malformations including mirror-image polydactyly

Eva Klopocki<sup>1,2,11</sup>, Christian Kähler<sup>3,11</sup>, Nicola Foulds<sup>4</sup>, Hitesh Shah<sup>5</sup>, Benjamin Joseph<sup>5</sup>, Hermann Vogel<sup>6</sup>, Sabine Lüttgen<sup>7</sup>, Rainer Bald<sup>8</sup>, Regina Besoke<sup>3</sup>, Karsten Held<sup>3</sup>, Stefan Mundlos<sup>1,2,9</sup> and Ingo Kurth<sup>\*,7,10</sup>

*PITX1* is a bicoid-related homeodomain transcription factor implicated in vertebrate hindlimb development. Recently, mutations in *PITX1* have been associated with autosomal-dominant clubfoot. In addition, one affected individual showed a polydactyly and right-sided tibial hemimelia. We now report on *PITX1* deletions in two fetuses with a high-degree polydactyly, that is, mirror-image polydactyly. Analysis of DNA from additional individuals with isolated lower-limb malformations and higher-degree polydactyly identified a third individual with long-bone deficiency and preaxial polydactyly harboring a heterozygous 35 bp deletion in *PITX1*. The findings demonstrate that mutations in *PITX1* can cause a broad spectrum of isolated lower-limb malformations including clubfoot, deficiency of long bones, and mirror-image polydactyly.

*European Journal of Human Genetics* (2012) 20, 705–708; doi:10.1038/ejhg.2011.264; published online 18 January 2012

**Keywords:** limb malformation; mirror-image-polydactyly; *PITX1*

## INTRODUCTION

Malformations of the limbs are common human birth defects. Genetic factors associated with these malformations often result in both upper and lower extremity anomalies.<sup>1</sup> A basic clinical classification differentiates between reduction defects and duplications, the latter most frequently causing pre- or postaxial polydactyly.<sup>2</sup> A rare variant of polydactyly is termed ‘mirror-image polydactyly’. This malformation is characterized by a high degree of polydactyly with a recognizable anterior/posterior axis of symmetry, that is, there is a hallux/thumb-like structure in the middle or an interdigital space with a flanking pair of digits that resemble a middle finger, index finger, thumb, toe or hallux.<sup>3</sup> The most lateral digits on each side typically resemble fifth fingers/toes. This malformation is often observed in combination with duplication of the ulna and/or fibula or with absent radius and/or tibia (hemimelia). The condition is observed as a leading clinical sign in Laurin–Sandrow Syndrome (MIM %135750).<sup>4,5</sup> A translocation breakpoint in 14q13 within the designated mirror-image polydactyly 1 gene (*MIPOL1*) has been described in a single case of Laurin–Sandrow syndrome,<sup>6</sup> but the genetic factors underlying mirror-image polydactyly remain largely unknown.

*PITX1* encodes a bicoid-transcription factor; mutations in which have previously been reported in a large autosomal-dominant family with a clubfoot malformation.<sup>7</sup> In this family penetrance was reduced and phenotype expression was variable, ranging from complete absence of congenital abnormalities to clubfoot, tibial hemimelia, and duplicated great toes in a single family member. This was the first indication that a broader spectrum of malformations might be due to *PITX1* dysfunction.

However, no other patient with tibial hemimelia was identified in this study. Recently, a second family with a deletion of the entire *PITX1* gene (chr5: 134 222 383–134 463 022, NCBI36/hg18) was reported in another multi-generation family with dominantly inherited clubfoot, where no additional malformations were present.<sup>8</sup>

Here, we report heterozygous deletions of 5q31 including the *PITX1* locus in two fetuses with lower-limb malformations including tibial hemimelia and mirror-image polydactyly. An intragenic deletion in *PITX1* was found in a third individual with a similar phenotype of tibial hemimelia but a milder polydactyly with some mirror elements.

## METHODS

### Subjects

Written informed consent was obtained from all the study participants after approval from the Institutional Review Boards at the participating institutions.

### Sequencing

The coding exons of *PITX1* (NM\_002653.4) and adjacent intronic sequences were amplified and sequenced. The primer sequences are given in Supplementary Table 2. Analysis was performed with SeqPilot software (JSI Medical Systems, Kippenheim, Germany). Four hundred control chromosomes were genotyped for the c.765\_799del variant using the identified mutant chromosome as positive control.

### Quantitative real-time PCR (qPCR)

Genomic DNA samples were obtained from EDTA-blood. Amplicons were located within the *PITX1* coding sequence and flanking regions (Supplementary Table 2). qPCR was performed as previously described.<sup>9</sup>

<sup>1</sup>Institut für Medizinische Genetik und Humangenetik, Charité Universitätsmedizin Berlin, Berlin, Germany; <sup>2</sup>Max Planck Institute for Molecular Genetics, Berlin, Germany; <sup>3</sup>genteQ, Hamburg, Germany; <sup>4</sup>Wessex Clinical Genetics Service, Southampton University Hospitals NHS Trust, Princess Anne Hospital, Southampton, UK; <sup>5</sup>Department of Orthopaedics, Kasturba Medical College, Manipal, Karnataka State, India; <sup>6</sup>Institut für Rechtsmedizin, University Hospital Hamburg-Eppendorf, Hamburg, Germany; <sup>7</sup>Institute of Human Genetics, University Hospital Hamburg-Eppendorf, Hamburg, Germany; <sup>8</sup>Clinical Centre Leverkusen, Leverkusen, Germany; <sup>9</sup>Berlin-Brandenburg Center for Regenerative Therapies (BCRT), Charité -Universitätsmedizin Berlin, Berlin, Germany; <sup>10</sup>Institute of Human Genetics, Jena University Hospital, Jena, Germany  
\*Correspondence: Dr I Kurth, Institute of Human Genetics, Jena University Hospital, Kollegien-gasse 10, 07743 Jena, Germany. Tel: +49 3641 934877; Fax: +49 3641 935502; E-mail: ingo.kurth@mti.uni-jena.de

<sup>11</sup>These authors contributed equally to this work.

Received 21 September 2011; revised 9 December 2011; accepted 14 December 2011; published online 18 January 2012

### Fluorescence *in situ* hybridization (FISH)

FISH analysis was performed on metaphase chromosomes according to standard procedures. The following BAC clones were used: D5S23/D5S721 (control probes), EGR1 (detection probe) (Vysis, Abbott Park, IL, USA).

### Array CGH

Total genomic DNA was purified by QIAamp DNA Blood Kit (QIAGEN, Hilden, Germany). Processing of the array (restriction digestion, fluorescent labeling, hybridization, washing, and scanning) was done in accordance with manufacturer's instructions (Agilent Oligonucleotide Array-Based CGH for Genomic DNA, Enzymatic Labelling for Blood (v6.3)). Extraction of microarray TIFF images had been done by 'Feature Extraction' and the following data analysis was done by the software 'Genomic Workbench 5.0' and 'CGH analytics' (both Agilent Technologies, Santa Clara, CA, USA). The following analysis settings were applied: algorithm ADM-2, filter 5 probes, log2ratio 0.29. The arrays used were: Human Genome CGH microarray kit 105A and 244K oligonucleotide array (Agilent Technologies).

### RESULTS

The first case we investigated was a fetus that had been diagnosed with high-degree polydactyly, hypoplasia of the corpus callosum, enlargement of the cisterna magna, and cardiomegaly. The fetus was stillborn at 34-weeks gestation after a sudden deceleration in fetal heart rate. Postmortem clinical examination revealed a small median cleft-palate, bilateral popliteal pterygia, talipes equinovarus together with mirror-image polydactyly with eight digits on each foot. The upper limbs did not show any abnormalities (Figure 1, case 1). Dysmorphic features included low-set ears, downslanting palpebral fissures, mild hypertelorism, and a flat nasal bridge. Array CGH analysis of DNA from amniotic fluid cells revealed a heterozygous 5.7 Mb deletion on chromosome 5q31.1-q31.2 (arr 5q31.1q31.2 (132 560 606–138 352 212)x1; NCBI36/hg18). This region contains 53 protein-coding genes, among them the bicoid-related homeodomain transcription factor *PITX1* (Supplementary Table S1). The deletion was verified using FISH in metaphases from cultured amniocytes with a region-specific BAC probe (Figure 1 and Supplementary Figure 2) and by quantitative PCR with amplicons located in exon 2 and 3 of *PITX1*. The deletion was not found in parental blood samples as confirmed by array CGH and FISH suggesting a *de novo* occurrence.

The second case (Figure 1, case 2) had undergone ultrasound examination at 15 weeks gestation that had revealed a single lower leg bone on the left as well as mirror-image polydactyly of the right foot together with bilateral clubfoot. The upper limbs did not show any abnormalities. In this patient, array CGH analysis of amniotic fluid cell DNA showed a heterozygous deletion of 4.9 Mb on 5q31 including *PITX1* (arr 5q31.1q31.2 (133 200 000–138 080 000)x1; NCBI36/hg18). The parents opted for termination of pregnancy.

To try to further elucidate the role of *PITX1* in complex lower-limb malformations, eight individuals diagnosed with isolated higher degree polydactyly/mirror-image polydactyly and/or tibial hemimelia,

but normally developed upper extremities underwent molecular genetic testing of *PITX1*. Direct sequencing of the coding exons of *PITX1* as well as quantitative PCR with probes in exon 2 and 3 of the gene was performed. In one individual with bilateral preaxial polydactyly, talipes equinovarus and right tibial hemimelia we identified a heterozygous deletion of 35 bp in exon 3 of *PITX1* (c.765\_799del) (Figure 1, case 3, and Supplementary Figure 1). This frameshift mutation at position 256 of the 314 amino-acid protein encoded by *PITX1* is predicted to cause a stop codon after incorporation of 303 additional irrelevant amino acids (p.Ala256ArgfsX303). The mutation was absent from lymphocyte DNA of the healthy mother and from 400 control chromosomes. DNA from the father was not available for molecular genetic testing. This frameshift mutation removes the C-terminal part of the protein and results in the loss of a 14 amino-acid motif, designated as OAR domain. This motif was suggested to trigger DNA binding and transactivation.<sup>10</sup> Thus, the mutation is highly likely to be pathogenic and thus supports a role for *PITX1* in complex lower-limb malformation.

### DISCUSSION

Evidence from several vertebrate species indicates that the transcription factor *Pitx1* is a key molecule in limb-specific morphogenesis. Targeted disruption of *Pitx1* in mice results in abnormalities of the hindlimb such as the fibula and tibia having equivalent diameters, loss of the patella and abnormalities of the calcaneus bone. This and the finding that *Pitx1* is expressed only in the hindlimb suggested a role in the determination of hindlimb morphology and identity.<sup>11–14</sup> The hypothesis was supported by studies performed in chick and mouse showing that misexpression of *Pitx1* in forelimbs results in the adoption of hindlimb characteristics.<sup>14,15</sup> In fish, pelvic loss in different natural populations of stickleback fish has occurred through regulatory mutations deleting a tissue-specific enhancer of *Pitx1* and demonstrates an evolutionary change in vertebrates via this genomic region.<sup>16</sup>



In humans, a missense mutation (p.E130K) in *PITX1* has been shown to segregate with dominantly inherited clubfoot with reduced penetrance in a large kindred, together with unilateral tibial hemimelia and preaxial polydactyly in one individual in this family.<sup>7</sup> The mutation is located in the DNA-binding domain of the protein and revealed reduced transactivation in a reporter gene assay.

A second family with dominantly inherited clubfoot was recently reported to harbor a 241 kb deletion including *PITX1*.<sup>8</sup> Interestingly, no signs of polydactyly or hemimelia were observed in this multi-generation family. The pathogenesis of *PITX1* in clubfoot was further supported by the finding that a subset of heterozygous *Pitx1* knockout mice displayed clubfoot.<sup>8</sup>




Our findings add considerable weight to the observation that *PITX1* deletions can cause a spectrum of lower-limb malformations including aplasia of long bones and high-degree polydactyly. Interestingly, the

**Figure 1** Haploinsufficiency of *PITX1* causes variable lower-limb malformations. *Case 1*: Reconstructions of postmortem CT scans, as well as clinical pictures showing the pes equinovarus configuration and mirror-image polydactyly. Whole body CT reconstruction shows normal upper limb and axial skeleton. FISH analysis indicating a deletion on 5q31 (arrow; green=control probe, red=locus specific probe). *Case 2*: Ultrasound examination in a fetus in the 15th gestational week shows polydactyly and long-bone deficiency. *Case 3*: Note long-bone deficiency, preaxial polydactyly as well as bilateral clubfoot. The right side is more severely affected (absent tibia) than the left. Molecular genetic testing of *PITX1* showed a second faster migrating amplicon of exon 3 by gel electrophoresis of the DNA from the affected individual (pat) in comparison with the maternal amplicon (co). Direct sequencing revealed a deletion of 35 bases (c.765\_799del). The electropherograms show the respective genomic sequence of exon 3 (upper panel) as well as a subcloned mutated allele (lower panel) from the patient. The breakpoint is indicated by a dashed line. *PITX1* is composed of three exons (E1-E3, NM\_002653.4, coding region in blue). Case 1 and case 2 showed larger interstitial genomic deletions on chromosome 5 including *PITX1* as determined by array CGH. The location and size of the three deletions is visualized by red bars and is given according to NCBI36/hg18.



case 1



case 2



case 3



CT-scan of case 1 also showed that the tibia and fibula become more alike and show a similar diameter, a feature that was also found in *Pitx1*<sup>-/-</sup> mice.<sup>17</sup> The findings also extend the phenotypic spectrum to include mirror-image polydactyly as variant of polydactylies. Of note, however, full mirror-image polydactyly was only present in the fetuses with larger genomic aberrations and therefore we cannot fully exclude that other genes in the deleted genomic region might cause, or at least influence this particular phenotype. As there is increasing evidence that copy number changes in non-coding genomic elements are implicated in bone disorders,<sup>18,19</sup> deletions of non-coding regulatory elements upstream or downstream of *PITX1* might also account for the clinical variability observed.

The digit number and identity along the anterior/posterior axis is controlled by signals from the zone of polarizing activity (ZPA) at the posterior margin of the limb bud. One of the key players in this scenario is the signaling molecule sonic hedgehog (SHH), which is expressed in the ZPA. In polydactylies, misregulation of SHH has been shown to be a crucial factor as demonstrated by mutations in the ZPA regulatory sequence of *SHH*, a non-coding regulatory sequence element located in 1 Mb distance of the *SHH* gene.<sup>20</sup> Copy number variations and point mutations in this regulatory region have been associated with a broad spectrum of polydactylies.<sup>19</sup> This overlapping clinical phenotype suggests an interplay between the Hedgehog pathway and *PITX1* in humans.

However, the situation might be different between men and mice. *Pitx1*<sup>-/-</sup> mice do not show a polydactyly indicating that *Pitx1* does not interfere to a significant degree with *Shh* signaling in the mouse. This hypothesis is corroborated by *Shh* expression analysis showing normal patterns of *Shh* expression in the ZPA of *Pitx1*<sup>-/-</sup> mice.<sup>12</sup> Only a combined inactivation of *Pitx1* and *Pitx2* resulted in abnormal *Shh* expression and *Pitx1* overexpression in the forelimb caused downregulation of *Shh*,<sup>14</sup> indicating some degree of interaction between the pathways. A similar discrepancy between mouse and human phenotypes has been observed for *WNT7A*. In humans, mutations in *WNT7A* result in severe limb truncations whereas inactivation in the mouse causes defects in dorsal-ventral patterning.<sup>21</sup>

In summary, our findings further define the role of *PITX1* in human limb development and illustrate that a spectrum of malformations is associated with *PITX1* haploinsufficiency.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### ACKNOWLEDGEMENTS

We thank the families for their cooperation and participation in this study. This study was supported by a grant to IK by the Fritz-Thyssen-Stiftung and to EK and SM by the Deutsche Forschungsgemeinschaft.

- Kornak U, Mundlos S: Genetic disorders of the skeleton: a developmental approach. *Am J Hum Genet* 2003; **73**: 447–474.
- Stricker S, Mundlos S: Mechanisms of digit formation: human malformation syndromes tell the story. *Dev Dyn* 2011; **240**: 990–1004.
- Stevenson RE, Judith G H: *Human Malformations and Related Anomalies*. New York: Oxford University Press, 2005.
- Marino-Enriquez A, Lapunzina P, Omenaca F, Morales C, Rodriguez JI: Laurin-Sandrow syndrome: review and redefinition. *Am J Med Genet A* 2008; **146A**: 2557–2565.
- Verghese R, Shah H, Rebello G, Joseph B: Pre-axial mirror polydactyly associated with tibial deficiency: a study of the patterns of skeletal anomalies of the foot and leg. *J Child Orthop* 2007; **1**: 49–54.
- Kondoh S, Sugawara H, Harada N *et al*: A novel gene is disrupted at a 14q13 breakpoint of t(2;14) in a patient with mirror-image polydactyly of hands and feet. *J Hum Genet* 2002; **47**: 136–139.
- Gurnett CA, Alaei F, Kruse LM *et al*: Asymmetric lower-limb malformations in individuals with homeobox *PITX1* gene mutation. *Am J Hum Genet* 2008; **83**: 616–622.
- Alvarado DM, McCall K, Aferol H *et al*: *Pitx1* haploinsufficiency causes clubfoot in humans and a clubfoot-like phenotype in mice. *Hum Mol Genet* 2011; **20**: 3943–3952.
- Klopocki E, Ott CE, Benatar N, Ullmann R, Mundlos S, Lehmann K: A microduplication of the long range SHH limb regulator (ZRS) is associated with triphalangeal thumb-polysyndactyly syndrome. *J Med Genet* 2008; **45**: 370–375.
- Semina EV, Reiter R, Leysens NJ *et al*: Cloning and characterization of a novel bicoid-related homeobox transcription factor gene, RIEG, involved in Rieger syndrome. *Nat Genet* 1999; **14**: 392–399.
- Szeto DP, Rodriguez-Esteban C, Ryan AK *et al*: Role of the Bicoid-related homeodomain factor *Pitx1* in specifying hindlimb morphogenesis and pituitary development. *Genes Dev* 1999; **13**: 484–494.
- Marcil A, Dumontier E, Chamberland M, Camper SA, Drouin J: *Pitx1* and *Pitx2* are required for development of hindlimb buds. *Development* 2003; **130**: 45–55.
- Minguillon C, Del Buono J, Logan MP: *Tbx5* and *Tbx4* are not sufficient to determine limb-specific morphologies but have common roles in initiating limb outgrowth. *Dev Cell* 2005; **8**: 75–84.
- DeLaurier A, Schweitzer R, Logan M: *Pitx1* determines the morphology of muscle, tendon, and bones of the hindlimb. *Dev Biol* 2006; **299**: 22–34.
- Logan M, Tabin CJ: Role of *Pitx1* upstream of *Tbx4* in specification of hindlimb identity. *Science* 1999; **283**: 1736–1739.
- Chan YF, Marks ME, Jones FC *et al*: Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a *Pitx1* enhancer. *Science* 2010; **327**: 302–305.
- Lancot C, Moreau A, Chamberland M, Tremblay ML, Drouin J: Hindlimb patterning and mandible development require the *Pitx1* gene. *Development* 1999; **126**: 1805–1810.
- Kurth I, Klopocki E, Stricker S *et al*: Duplications of noncoding elements 5' of *SOX9* are associated with brachydactyly-anonychia. *Nat Genet* 2009; **41**: 862–863.
- Klopocki E, Mundlos S: Copy-number variations, noncoding sequences, and human phenotypes. *Annu Rev Genomics Hum Genet* 2011; **12**: 53–72.
- Hill RE: How to make a zone of polarizing activity: insights into limb development via the abnormality preaxial polydactyly. *Dev Growth Differ* 2007; **49**: 439–448.
- Woods CG, Stricker S, Seemann P *et al*: Mutations in *WNT7A* cause a range of limb malformations, including Fuhrmann syndrome and Al-Awadi/Raas-Rothschild/Schinzler phocomelia syndrome. *Am J Hum Genet* 2006; **79**: 402–408.

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)