

SHORT REPORT

High prevalence of genetic variants previously associated with LQT syndrome in new exome data

Lena Refsgaard^{1,2}, Anders G Holst^{1,2}, Golnaz Sadjadieh^{1,2}, Stig Haunso^{1,2,3}, Jonas B Nielsen^{1,2} and Morten S Olesen^{*,1,2}

To date, hundreds of variants in 13 genes have been associated with long QT syndrome (LQTS). The prevalence of LQTS is estimated to be between 1:2000 and 1:5000. The knowledge of genetic variation in the general population has until recently been limited, but newly published data from NHLBI GO Exome Sequencing Project (ESP) has provided important knowledge on this topic. We aimed to investigate the prevalence of previously LQTS-associated variants in ESP (5400 individuals), in order to identify possible false-positive LQTS variants. With this aim, we performed a search for previously published LQTS-associated variants in ESP. In addition, a PolyPhen-2 prediction was conducted, and the four most prevalent LQTS-associated variants with significant functional effects present in ESP were genotyped in a second control population. We identified 33 missense variants previously associated with LQTS in ESP. These 33 variants affected 173 alleles and this corresponded to a LQTS prevalence of 1:31 in the ESP population. PolyPhen-2 predicted 30% of the 33 variants present in ESP to be benign compared with 13% among LQTS-associated variants not present in ESP ($P=0.019$). Genotyping of the four variants *KCNH2* P347S; *SCN5A*: S216L, V1951L; and *CAV3* T78M in the control population ($n=704$) revealed prevalences comparable to those of ESP. Thus, we identified a much higher prevalence of previously LQTS-associated variants than expected in exome data from population studies. Great caution regarding the possible disease causation of some of these variants has to be taken, especially when used for risk stratification in family members.

European Journal of Human Genetics (2012) 20, 905–908; doi:10.1038/ejhg.2012.23; published online 29 February 2012

Keywords: LQTS; false positive; exome; next-generation sequencing; NGS; arrhythmia

INTRODUCTION

Congenital long QT syndrome (LQTS) is an inherited cardiac disorder characterised by a prolonged QT interval on the electrocardiogram and sudden cardiac death secondary to cardiac arrhythmias.

The prevalence of LQTS in the general population has been estimated to be between 1:2000 and 1:5000.¹ In most cases, LQTS is considered a monogenic disorder, and to date hundreds of variants in 13 genes have been associated with this syndrome.^{1,2} However, variants previously reported to be causative in LQTS have subsequently been shown to be rare variants with only a modest disease-modifying effect or even non-pathogenic.³

Until recently, there has only been limited knowledge regarding the genetic variation in the general population, especially with regards to low-frequency variants. This was changed in June 2011, when whole exome data from the NHLBI GO Exome Sequencing Project (ESP) was published.⁴ In order to identify possible false-positive LQTS variants reported in the literature, we aimed to investigate the prevalence of previously LQTS-associated variants in new ESP exome data and compare the prevalence of these variants with the expected prevalence of LQTS in the same population.

METHODS

In ESP, next-generation sequencing of all protein-coding regions in 5400 persons from different population studies were carried out.⁴ No clinical data

were available on the ESP population, nor at request. By literature search, we found inclusion- and exclusion criteria of 9 out of 12 populations used in ESP. None of these have specifically included persons with channelopathies or other heart diseases and at least two cohorts have excluded such patients.

ESP exome data were systematically searched for previously published missense and nonsense LQTS-associated variants reported in a recent comprehensive review by Hedley *et al*¹ and in data from the The Human Gene Mutation Database (HGMD).⁵ In addition, we have included the recently published *KCNJ5* variant reported by Yang *et al*,² in order to include all 13 genes so far associated with LQTS. Due to the lack of data regarding variants positioned in introns and UTR regions in ESP, these could not be included.

The literature was searched for functional data and family co-segregation of all the previously identified LQTS-associated variants also identified in the ESP population. Co-segregation was defined as at least two family members both having the phenotype and the genotype. In addition, we conducted a PolyPhen-2 prediction on all previously reported LQTS-associated missense variants and compared protein damage predictions between LQTS-associated variants identified in ESP with those variants not identified in ESP.⁶ Nonsense variants were assumed probably damaging. Differences in proportions of PolyPhen-2 predictions between groups were assessed using a Fisher's exact test.

Furthermore, we genotyped (Sanger sequencing, as described previously⁷) variants with (1) convincing LQTS association in terms of significant functional effects on channel function and (2) a prevalence in ESP European American

¹The Danish National Research Foundation Centre for Cardiac Arrhythmia, Copenhagen, Denmark; ²Laboratory of Molecular Cardiology, Department of Cardiology, The Heart Centre, University Hospital of Copenhagen, Rigshospitalet, Denmark; ³Department of Medicine and Surgery, Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark

*Correspondence: Dr MS Olesen, Laboratory for Molecular Cardiology, Department of Cardiology, The Heart Centre, University Hospital of Copenhagen, Rigshospitalet, 9312, Juliane Mariesvej 20, Copenhagen OE 2100, Denmark. Tel: +45 35456506; Fax: +45 35456500; E-mail: morten.salling.olesen@rh.regionh.dk

Received 15 September 2011; revised 4 January 2012; accepted 17 January 2012; published online 29 February 2012

population high enough ($\geq 7:5295$) to have a reasonable chance to be detected in our own two independent healthy control populations ($n=704$) of Northern European origin with no history of arrhythmias or other cardiac diseases,^{7,8} and with available ECGs.

RESULTS

In all, 33 out of 631 missense and nonsense variants previously associated with LQTS (5.2%) were found in the ESP population. All 33 variants were missense variants and affected 173 alleles. Thirteen of

Table 1 Variants associated with LQTS present in the ESP population

Gene	Variant	Amino acid	Type	Minor allele ^a	Total alleles ^a	Variant associated with	PolyPhen-2 results	Level of LQTS evidence	
								Family co-segregation	Functional data
KCNQ1	c.532G>A	A178T	Missense	2	10 756	LQT1 ^b	Probably damaging	Equivocal	No data available
	c.613G>A	V205M	Missense	1	10 756	LQT1 ^c	Probably damaging	Yes, but incomplete penetrance	Loss of function
	c.959C>A	P320H	Missense	1	10 754	LQT1 ^c	Probably damaging	Yes	Loss of function
	c.1189C>T	R397W	Missense	2	10 758	LQT1 ^b	Probably damaging	No data available	No data available
	c.1352G>A	R451Q	Missense	1	10 754	LQT1 ^c	Possibly damaging	No data available	No data available
	c.1354C>T	R452W	Missense	2	10 754	LQT1 ^c	Probably damaging	No data available	No data available
	c.1576A>G	K526E	Missense	2	10 758	LQT1 ^b	Probably damaging	No data available	No data available
	c.1831G>A	D611N	Missense	3	10 716	LQT1 ^d	Possibly damaging	No data available	No data available
	c.1927G>A	G643S	Missense	79	10 528	aLQTS ^d	Benign	Yes, but incomplete penetrance	Loss of function
	KCNH2	c.1039G>A	P347S	Missense	7	10 758	LQT2 ^c	Benign	Equivocal
c.1912A>G		K638E	Missense	1	10 732	LQT2 ^b	Probably damaging	No data available	No data available
c.2653G>A		R885C	Missense	1	10 758	LQT2 ^c	Probably damaging	No data available	No effect
c.2660G>A		R887H	Missense	1	10 758	LQT2 ^b	Probably damaging	No data available	No data available
c.2948G>A		T983I	Missense	3	10 740	LQT2 ^b	Probably damaging	No data available	No data available
SCN5A	c.647G>A	S216L	Missense	11	10 248	LQT3 ^d	Probably damaging	No data available	Gain of function
	c.1384G>A	E462K	Missense	1	9966	LQT3 ^b	Probably damaging	No data available	No data available
	c.1715G>T	A572D	Missense	18	10 194	sLQTS ^c	Benign	No	Gain of function
	c.1844C>T	G615E	Missense	3	10 264	diLQTS ^c	Benign	No data available	Gain of function/no effect
	c.1852G>A	L618F	Missense	21	10 274	diLQTS ^b	Probably damaging	No data available	No effect
	c.1855G>A	L619F	Missense	1	10 262	LQT3 ^c	Benign	Equivocal	Gain of function
	c.2066G>A	R689H	Missense	1	10 488	LQT3 ^b	Benign	No data available	No data available
	c.2074G>T	Q692K	Missense	2	10 494	LQT3 ^c	Benign	No data available	No data available
	c.3578G>A	R1193Q	Missense	7	10 747	aLQTS ^b	Benign	No data available	Gain of function
	c.3911G>A	T1304M	Missense	5	10 384	LQT3 ^b	Probably damaging	Yes	Gain of function
	c.5336C>T	T1779M	Missense	1	10 758	LQT3 ^b	Probably damaging	No data available	No data available
	c.5360C>T	S1787N	Missense	6	10 758	LQT3 ^b	Possibly damaging	No data available	Gain of function
	c.5711C>T	S1904L	Missense	10	10 416	LQT3 ^d	Probably damaging	Equivocal	Gain of function
	c.5848C>A	V1951L	Missense	17	10 278	LQT3 ^d	Benign	No data available	Gain of function
	c.5873C>T	R1958Q	Missense	2	10 258	LQT3 ^b	Benign	No data available	No data available
c.6016G>C	P2006A	Missense	12	10 108	sLQTS ^b	Benign	Yes, but incomplete penetrance	Gain of function	
ANK2	—	—	—	—	—	—	—	—	—
KCNE1	c.253C>T	D85N	Missense	94	10 758	aLQTS ^b	Benign	No data available	Loss of function
	c.325G>A	V109I	Missense	2	10 758	LQT5 ^c	Benign	Equivocal	Loss of function
KCNE2	c.22A>G	T8A	Missense	48	10 758	diLQTS ^b	Probably damaging	No data available	Loss of function
	c.161T>C	M54T	Missense	3	10 758	LQT6 ^d	Probably damaging	No data available	Loss of function
	c.170T>C	I57T	Missense	4	10 758	LQT6 ^d	Probably damaging	Equivocal	No effect/loss of function
	c.229C>T	R77W	Missense	2	10 758	LQT6 ^c	Probably damaging	No data available	No effect
KCNJ2	—	—	—	—	—	—	—	—	—
CACNA1C	—	—	—	—	—	—	—	—	—
CAV3	c.233C>T	T78M	Missense	46	10 758	LQT9 ^c	Probably damaging	No data available	Gain of function
SCN4B	—	—	—	—	—	—	—	—	—
AKAP9	c.4709C>T	S1570L	Missense	1	10 758	LQT11 ^d	Benign	Yes	Loss of function
SNTA1	c.1169C>T	A390V	Missense	1	10 758	LQT12 ^b	Possibly damaging	No data available	Gain of function
KCNJ5 ^e	—	—	—	—	—	—	—	—	—

Abbreviations: aLQTS, acquired LQTS; diLQTS, drug-induced LQTS; sLQTS, suspected LQTS.

^aData from NHLBI Exome Sequencing Project (ESP).⁴

^bHedley *et al*.¹ and HGMD.⁵

^cHGMD.⁵

^dHedley *et al*.¹

^eYang *et al*.²

Table 2 Allele frequencies in ESP and in control population

Gene	Amino acid	rs#	MAF ESP				MAF controls	
			African-American (%)	European-American (%)	Total (%)	Northern European (%)	QTc interval (ms)	
<i>KCNH2</i>	P347S	rs138776684	0.00	0.10	0.07	0.14	414, 436	
<i>SCN5A</i>	S216L	rs41276525	0.03	0.15	0.11	0.07	400	
	V1951L	rs41315493	0.38	0.06	0.17	0.07	426	
<i>CAV3</i>	T78M	rs72546668	0.38	0.46	0.43	0.43	410, 436, 462, 435, 412, RBBB ^a	

Abbreviation: MAF, minor allele frequency.

^aRight bundle branch block, QTc estimate not possible.

the variants were identified in three or more alleles. No nonsense variants were found. The previously LQTS-associated variants identified in the ESP population are listed in Table 1. On an average, the genes investigated in this study have been screened in 5298 individuals. If we assume that no individuals in the ESP population harbour more than one of the LQTS-associated variants and that all are heterozygote carriers, it corresponds to a LQTS genotype prevalence of 1:31 (173:5298). Thirteen variants were present in three or more alleles corresponding to a LQTS genotype prevalence of 1:37 (145:5298). Six variants associated with acquired/drug-induced LQTS were also identified in ESP; these variants were not included in the present analysis. Literature search regarding functional data and family co-segregation revealed that 19 of the 33 identified LQTS-associated variants have been investigated functionally and 16 of these did have functional effects, but there was a striking lack of data regarding family co-segregation (Table 1 and Supplementary Table).

PolyPhen-2 analysis of the 33 LQTS variants predicted 19 (58%) to be probably damaging, 4 (12%) to be possibly damaging, and 10 (30%) were predicted to be benign (Table 1). Of the remaining 558 missense and 40 nonsense LQTS-associated variants, not present in ESP, 451 (75%) were predicted probably damaging, 69 (12%) possibly damaging, and 78 (13%) benign. This difference in the distribution of the three PolyPhen-2 categories among the two groups was statistically significant both for the overall comparison ($P=0.027$) and when comparing the categories benign ($P=0.016$) and probably damaging ($P=0.038$) independently. As no nonsense variants were found in ESP, a more conservative approach is to compare only the missense variants in the two groups; 558 missense LQTS-associated variants were not present in ESP and of these 411 (74%) were predicted probably damaging, 69 (12%) possibly damaging, and 78 (14%) benign. In this case, only a borderline significant difference was found ($P=0.051$) for the overall comparison.

Four variants (*KCNH2* P347S; *SCN5A* S216L, V1951L, and *CAV3* T78M) were, based on our criteria, genotyped in our control populations ($n=704$), and this revealed variant prevalences comparable to those found in ESP (Table 2). One of the variant carriers harbouring *CAV3* T78M had a QTc interval of 462 ms, but all other variant carriers had normal QTc intervals ($QTc \leq 440$ ms) including four other carriers of *CAV3* T78M (Table 2).

DISCUSSION

We identified 33 missense variants out of 631 (5.2%) variants previously associated with LQTS in the ESP data.

These 33 variants correspond to a LQTS genotype prevalence of 1:31 in the ESP. If the prevalence of LQTS is set at 1:2000 (a high estimate), then we would expect only three persons in the ESP population to have LQTS. Thus, we found a very large overrepresentation of previously LQTS-associated variants in the ESP, indicating a

very dubious LQTS association for these variants as a group. Of course, we cannot exclude the possibility that there might be an overrepresentation of LQTS patients in the ESP population, although this seems unlikely, due to the size of the overrepresentation and the fact that ESP is based on population studies mainly representing the general population. Even if we had clinical data on all individuals, we could not make a significantly different conclusion due to the possibility of reduced penetrance of the variants.

To test the possible overrepresentation of LQTS-associated variants in ESP we genotyped the four most prevalent LQTS-associated variants with significant functional effects present in ESP in two cohorts of healthy individuals with available ECGs.^{7,8} The four variants were present in our control population with prevalences consistent with those of ESP (Table 2). Thus, the present results indicate that overrepresentation of LQTS-associated variants does not seem to be a problem in ESP, but more importantly the results confirm that at least these four LQTS-associated variants with high prevalence in ESP and significant functional effects are not monogenic causes of LQTS.

In addition, we did a PolyPhen-2 prediction both for the variants found in ESP and the ones we did not find. Of the 33 variants found in ESP, 30% were predicted to be benign compared with 13% among the variants not found in ESP ($P=0.016$). This further strengthens the argument that these variants are not disease causing.

The majority of the variants have been shown to have functional effects in electrophysiological experiments performed in heterologous cell models. This suggests that they may have some pathological influence, but only a very small fraction of studies showed family co-segregation. Thirteen of the variants were present in three or more alleles in ESP, and thus it seems reasonable to assume that these variants are not monogenic causes of LQTS, as 1:37 are carriers of one of these variants.

The remaining 20 variants are present in just one or two alleles in ESP and an interpretation of the pathogenicity of these variants are therefore much less straightforward.

Taken together, the present results indicate that a large proportion of the 33 variants previously associated with LQTS but now identified in ESP are very unlikely to be monogenic causes of LQTS and are maybe just innocent bystanders. Others are likely to be variants with reduced penetrance, or in some cases disease modifiers. Thus, great caution must be taken regarding the possible disease causation of these variants when found in LQTS (Table 1).

It is noteworthy, that 94.8% of the previously LQTS-associated variants were not present in ESP and thus the pathogenicity of the majority of these variants seems to be corroborated.

Genetic screening has become an important tool in at least family examinations of LQTS. It is for this reason important that variants being associated with LQTS are truly disease causing. A patient with genuine LQTS but a false-positive variant result might have another

true disease-causing variant that is not found and this could lead to misdiagnosis of family members with consequences for treatment, clinical advice, and follow-up.

CONCLUSION

In recently released exome data, we identified a much higher prevalence of previously LQTS-associated variants than expected based on the prevalence of LQTS in the general population. Our findings suggest that some of the previously associated LQTS variants are false-positive findings and that great caution must be taken regarding the possible disease causation of these variants, especially when used for risk stratification in family members.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by The Danish National Research Foundation Centre for Cardiac Arrhythmia, The John and Birthe Meyer Foundation, The Research Foundation of the Heart Centre Rigshospitalet, and The Arvid Nilsson Foundation.

- 1 Hedley PL, Jørgensen P, Schlamowitz S *et al*: The genetic basis of long QT and short QT syndromes: a mutation update. *Hum Mutat* 2009; **30**: 1486–1511.
- 2 Yang Y, Yang Y, Liang B *et al*: Identification of a Kir3.4 mutation in congenital long QT syndrome. *Am J Hum Genet* 2010; **86**: 872–880.
- 3 Tester DJ, Valdivia C, Harris-Kerr C *et al*: Epidemiologic, molecular, and functional evidence suggest A572D-SCN5A should not be considered an independent LQT3-susceptibility mutation. *Heart Rhythm* 2010; **7**: 912–919.
- 4 Exome Variant Server, NHLBI Exome Sequencing Project (ESP), Seattle, WA (url: <http://snp.gs.washington.edu/EVS/>) (Accessed 2011 Dec 05).
- 5 The Human Gene Mutation Database at the Institute of Medical Genetics in Cardiff (url: <http://www.hgmd.org/>) (Accessed 2011 Dec 05).
- 6 Adzhubei IA, Schmidt S, Peshkin L *et al*: A method and server for predicting damaging missense mutations. *Nat Methods* 2010; **7**: 248–249.
- 7 Olesen MS, Jespersen T, Nielsen JB *et al*: Mutations in sodium channel β -subunit SCN3B are associated with early-onset lone atrial fibrillation. *Cardiovasc Res* 2011; **89**: 786–793.
- 8 Olesen MS, Jensen NF, Holst AG *et al*: A novel nonsense variant in Nav1.5 cofactor MOG1 eliminates its sodium current increasing effect and may increase the risk of arrhythmias. *Can J Cardiol* 2011; **27**: 523e17–523e23.



This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported Licence. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)