

ARTICLE

Five children with deletions of 1p34.3 encompassing *AGO1* and *AGO3*

Mari J Tokita^{1,2}, Penny M Chow^{1,2}, Ghayda Mirzaa^{1,2}, Nicola Dikow³, Bianca Maas³, Bertrand Isidor⁴, Cédric Le Caignec⁴, Lynette S Penney⁵, Giovanni Mazzotta⁶, Laura Bernardini⁷, Tiziana Filippi⁸, Agatino Battaglia⁸, Emilio Donti⁹, Dawn Earl^{1,2} and Paolo Prontera^{*,9}

Small RNAs (miRNA, siRNA, and piRNA) regulate gene expression through targeted destruction or translational repression of specific messenger RNA in a fundamental biological process called RNA interference (RNAi). The Argonaute proteins, which derive from a highly conserved family of genes found in almost all eukaryotes, are critical mediators of this process. Four *AGO* genes are present in humans, three of which (*AGO 1, 3, and 4*) reside in a cluster on chromosome 1p35p34. The effects of germline *AGO* variants or dosage alterations in humans are not known, however, prior studies have implicated dysregulation of the RNAi mechanism in the pathogenesis of several neurodevelopmental disorders. We describe five patients with hypotonia, poor feeding, and developmental delay who were found to have microdeletions of chromosomal region 1p34.3 encompassing the *AGO1* and *AGO3* genes. We postulate that haploinsufficiency of *AGO1* and *AGO3* leading to impaired RNAi may be responsible for the neurocognitive deficits present in these patients. However, additional studies with rigorous phenotypic characterization of larger cohorts of affected individuals and systematic investigation of the underlying molecular defects will be necessary to confirm this.

European Journal of Human Genetics (2015) 23, 761–765; doi:10.1038/ejhg.2014.202; published online 1 October 2014

INTRODUCTION

RNA interference (RNAi), also called posttranscriptional gene silencing (PTGS), is a process in which small noncoding RNA molecules regulate gene expression by interfering with translation of corresponding messenger RNA (mRNA).¹ MicroRNA (miRNA) are the primary RNAi effectors, and it is estimated that miRNAs regulate the translation of at least one-third of proteins encoded by the human genome.² MiRNA undergo a complex biosynthetic process starting in the nucleus where the Drosha-Pasha microprocessor complex effects the first cleavage, transforming primary transcripts, or pri-miRNA, into pre-miRNA.³ Pre-miRNA is then exported to the cytoplasm where Dicer performs the second cleavage, generating mature miRNA.³ The miRNA is then integrated into an RNA-induced silencing complex (RISC), which guides recognition of a complementary sequence of mRNA. Argonaute proteins (*AGO1-4*), which constitute the catalytic component of the RISC, cleave the mRNA, effectively blocking translation.³ *AGO* proteins are also directly involved in miRNA biogenesis⁴ and thus fulfill several essential roles in the RNA interference pathway.

Animal models carrying variants in *DGCR8* (which encodes Pasha), *DICER1*, *AGO1*, and *AGO2* display severe neurological impairment and defective development and function of the central nervous system.^{5–8} In humans, the neuropsychiatric features seen in patients with 22q11.2 microdeletion syndrome are thought to result from

haploinsufficiency of the *DGCR8* gene and associated perturbations in miRNA biogenesis.⁹

We present five patients with different but overlapping deletions involving the 1p34.3 locus, ranging in size from 1.1 to 3.1 Mb. The shared deleted region is approximately 289 kb and encompasses the *AGO1* and *AGO3* genes. The effect of *AGO* dosage alterations in humans has not been previously reported. We hypothesize that loss of *AGO1* and *AGO3* in these patients accounts for the shared neurocognitive phenotype, which includes hypotonia, poor feeding, and developmental delay.

MATERIALS AND METHODS

Contact between physicians caring for the patients described was facilitated by the Decipher Database (<https://www.decipher.sanger.ac.uk>). Variants for each patient have been submitted to this web resource and the corresponding Decipher ID specified at the end of each clinical report.

Clinical reports

Proband 1 was the first child born to non-consanguineous parents of Norwegian and German heritage. She was born at term by spontaneous vaginal delivery weighing 3000 g (10th–25th percentile) and measuring 49.5 cm in length (50th percentile) following an unremarkable pregnancy. Speech delays were noted between 18 and 24 months of age. At her initial genetics visit at age 2 years and 10 months, she was communicating primarily with poorly articulated one word utterances, gestures, and tonal inflections.

¹Division of Genetic Medicine, University of Washington, Seattle, WA, USA; ²Seattle Children's Hospital, Seattle, WA, USA; ³Institute of Human Genetics, Heidelberg University, Heidelberg, Germany; ⁴Génétique Médicale, Centre Hospitalier Universitaire de Nantes, Nantes, France; ⁵Department of Pediatrics, Dalhousie University, Halifax, Nova Scotia, Canada; ⁶School of Specialization in Childhood Neurology and Psychiatry, University of Perugia, Perugia, Italy; ⁷Mendel Laboratory, IRCSS Casa Sollievo della Sofferenza Hospital, San Giovanni Rotondo, Foggia, Italy; ⁸The Stella Maris Clinical Research Institute for Child and Adolescent Neuropsychiatry, Calambrone (Pisa), Italy; ⁹Medical Genetics Unit, Hospital Santa Maria della Misericordia, University of Perugia, Perugia, Italy

*Correspondence: Professor P Prontera, Medical Genetics Unit, Hospital Santa Maria della Misericordia, University of Perugia, via Enrico dal Pozzo, 06123 Perugia, Italy. Tel: +00 3907 5578 3555; Fax: +00 3907 5578 3945; E-mail: paolo.prontera@ospedale.perugia.it or pprontera@hotmail.com

Received 29 April 2014; revised 14 July 2014; accepted 20 August 2014; published online 1 October 2014




Figure 1 Facial phenotypes of proband 3 (a) and proband 5 (b).

At her most recent evaluation at 3 years 9 months, her weight was 13.5 kg (16th percentile), height 95.9 cm (25th percentile), and occipitofrontal circumference (OFC) 48.1 cm (10th–25th percentile). She had mildly impaired balance but was able to walk and jump without difficulty. She was using six-word phrases and approximately 50% of her speech was understandable. She was able to name body parts and count to 15. Receptive language was intact and age-appropriate. Pertinent exam findings included: high nasal bridge, upturned nasal tip, mildly down-slanting palpebral fissures, high-arched palate, elongated-appearing digits, tented upper lip, and retrognathia. She had mild hypotonia with pronated feet.

Diagnostic workup included fragile X testing which was normal, and array CGH (Roche Nimblegen, CGX-6, 135K, Madison, WI, USA) which showed a 2.7 Mb deletion at 1p34.3 (chr1.hg18:g.(36,103,080-36,130,907)_(38,861,099-38,893,034)del) (NCBI 36/hg18), encompassing 27 OMIM genes. The proband's parents have declined carrier testing to date (Decipher ID: 291841).

Proband 2 was the first child of non-consanguineous parents of French origin. She was born at term following an unremarkable pregnancy. Her birth weight was 2810 g (10th percentile), length 48 cm (25th percentile), and OFC 34.5 cm (25th–50th percentile). She had neonatal hypotonia and feeding difficulties. Development was subsequently delayed. She walked independently at 21 months, was able to string two words together at 2 years, and was speaking in short sentences at 3 years of age.

At her most recent evaluation at 10 years 6 months, her weight was 25 kg (3rd percentile), height was 137 cm (25th percentile), and OFC was 53 cm (25th percentile). She displayed ongoing learning delays and was just starting to learn how to read and write. She had velopharyngeal insufficiency that affected her speech. On exam, she had bilateral epicanthal folds, a triangular upper lip, fullness of the periorbital region, and long thin fingers.

Array CGH (Agilent Catalog, 44K) showed a 2.2 Mb deletion at 1p34.3 (chr1.hg18:g.(35,878,982-35,927,274)_(38,364,135-38,603,709)del) (NCBI 36/hg18), including 31 OMIM genes. Parental testing confirmed this deletion to be *de novo* (Decipher ID: 004679).

Proband 3 was born to healthy, non-consanguineous parents of Italian origin after an unremarkable pregnancy at 42 weeks gestational age. Birth weight was 3000 g (10th–25th percentile). She had neonatal breastfeeding difficulties and severe bilateral hip dislocation requiring surgical management. She did not walk independently until 3 years of age and also had speech delays. At age 17, she scored less than 55 on formal IQ testing consistent with moderate intellectual disability. At her most recent evaluation at 18 years of age, she weighed 46 kg (5th percentile), was 166 cm tall (50th percentile), and had an OFC of 50 cm (< 1st percentile; –3 to 4 SD). She had a friendly disposition, poor vocabulary,

and a limited attention span. Her voice had a nasal quality due to velopharyngeal insufficiency. Facial exam revealed bilateral ptosis, bitemporal narrowing, down-slanting palpebral fissures, small ears, retrognathia, a high-arched palate, maxillary hypoplasia, and a long mandible (Figure 1a). She also had moderate joint laxity, hypotonia, and flat feet.

Diagnostic workup included hematologic and biochemical testing, urinalysis, ECG, and echocardiography, all of which were normal. Brain MRI showed mild cortical atrophy in the occipital and parietal lobes. Array CGH (Cytochip, BluGnome, 180 K with 75–100 kb resolution) showed a 1.1 Mb deletion at 1p34.3 (chr1.hg18: g.(35,691,738_35,705,605)_(36,825,269_36,848,510)del) (NCBI 36/hg18) encompassing 18 OMIM genes. The result was confirmed by FISH and was found to be *de novo* (Decipher ID: 256601).

Proband 4 was the second child born to non-consanguineous healthy parents of German origin. He was born at 41 weeks gestational age after a pregnancy complicated by intrauterine growth restriction (IUGR) diagnosed at 23 weeks. His birth weight was 2660 g (< 1st percentile, –3.29 SD), length 47 cm (< 1st percentile, –2.66 SD), and OFC 34 cm (10th percentile). Following delivery, he required treatment for neonatal sepsis, showed marked muscular hypotonia with failure to thrive, and required nasogastric tube feeding for several weeks. At 7 months of age, his OFC dropped below the 3rd percentile. At 11 months of age he was 69 cm long (–3.1 SD) and had an OFC of 44 cm (–3 SD). He had a friendly disposition with prolonged eye contact and excessive babbling. He was able to turn independently from front to back and from back to front. Physical exam showed a prominent forehead, flat nasal bridge, deep-set eyes with short palpebral fissures, a small chin, and hypospadias. At his most recent follow-up at age 17 months, his muscular hypotonia had improved, however, he was not yet able to sit unsupported.

Oligo/SNP Array analysis (Affymetrix, CytoScan HD, Santa Clara, CA, USA) showed a 3.1 Mb deletion in chromosomal region 1p34.3 (chr1.hg18:g.(35,543,882_35,544,184)_(38,659,938_38,667,647)del) (GRCH36/hg18) encompassing 33 OMIM genes. Metaphase FISH analysis of the patient and his parents confirmed the deletion was a *de novo* occurrence (Decipher ID: 291728).

Proband 5 was the first child born to healthy, non-consanguineous parents of Romanian descent. He was delivered at 38 weeks gestational age for IUGR by cesarean section, which was complicated by perinatal asphyxia. Birth weight was 2400 g (< 10th percentile), birth length was 49 cm (50th percentile), and he was reportedly turricephalic. At delivery, he had poor suction but could breastfeed.

Developmentally, he started babbling at 6 months, was speaking single words at 18 months and short phrases by 2½ years. He walked independently at 16 months and was toilet-trained by 2½ years. At his initial genetics evaluation at

Table 1 Summary of the clinical and cytogenetic findings of the 1p34.3 cohort (n = 5)

	Proband 1	Proband 2	Proband 3	Proband 4	Proband 5
Array platform (hg18)	Roche Nimblegen CGX-6 135K	Agilent Catalog 44K Array	BluGnome 180K Cytochip	Affymetrix CytoScan HD	Affymetrix GeneChip 6.0
Deletion	1p34.3	1p34.3	1p34.3	1p34.3	1p34.3
Coordinates (hg18)	36 130 907–38 861 099	35 927 274–38 364 135	35 705 605–36 825 269	35 544 184–38 659 938	35 219 831–36 415 737
Size	~2.7 Mb	~2.2 Mb	~1.1 Mb	~3.1 Mb	~1.2 Mb
Inheritance	Unknown	<i>De novo</i>	<i>De novo</i>	<i>De novo</i>	<i>De novo</i>
Age at last evaluation	3 years 9 months	10 years 6 months	18 years	17 months	13 years
Development	Motor and speech delay	Motor and speech delay. Special education	Motor and speech delay. Moderate intellectual disability	Motor and speech delay	Speech delay. Moderate intellectual disability
Height (percentile)	24th	25th	50th	< 1st (-3.2 SD)	90th–97th
Weight (percentile)	15th	2nd	5th	1st (-2.4 SD)	> 97th
OFC (percentile)	3rd–10th	25th	< < 3rd (-3 to 4 SD)	< 1st (-3.3 SD)	2nd–10th
Dysmorphic features	High nasal bridge, upturned nasal tip, mildly down-slanting palpebral fissures, high-arched palate with dental crowding, retrognathia, long-appearing fingers, dysplastic toenails, pronated feet	Bilateral epicanthal folds, triangular upper lip, fullness of periorbital region, long thin fingers	Long hypotonic face, bitemporal narrowing, down-slanting palpebral fissures, ptosis, small ears, retrognathia, high-arched palate, maxillary hypoplasia, long mandible, flat feet	Prominent forehead, deep-set eyes, short palpebral fissures, small chin	Flat occiput, straight eyebrows, small mouth, bifid uvula, highly arched palate, dorsal left convex kyphoscoliosis, bilateral pes planus
Poor feeding	+	+	+	+	±
Hypotonia	+	+	+	+	+
Velopharyngeal insufficiency	+	+	+	-	-
Other medical problems	Astigmatism	Buccofacial dyspraxia	Severe bilateral hip dislocation, joint laxity, nasal voice	Hypospadias	Joint laxity
Pregnancy and delivery	Uncomplicated	Uncomplicated	Uncomplicated	IUGR at 23 weeks, neonatal sepsis	IUGR, perinatal asphyxia
Age at delivery	38 weeks	37 weeks	42 weeks	41 weeks	38 weeks
Family history	Male cousin with possible pervasive developmental delay	Not available	Healthy parents	Healthy parents	Healthy parents




Figure 3 miRNA biogenesis pathway and consequences of dysfunction observed in animal models and in humans (adapted from Yang and Lai¹⁹).

exclude the possibility that other genes in the deleted region may be causative of or contribute to the phenotype. As such, additional studies of larger numbers of affected individuals in conjunction with functional assays will be necessary to more clearly define the developmental consequences of AGO gene haploinsufficiency.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the patients and their families for their kind cooperation. This study makes use of data generated by the DECIPHER Consortium. A full list of centers who contributed to the generation of the data is available from <http://decipher.sanger.ac.uk> and via email from decipher@sanger.ac.uk. Funding for the project was provided by the Wellcome Trust. Electronic database information is available from Online Mendelian Inheritance in Man (<http://www.omim.org>); Decipher-Wellcome Trust Sanger Institute (<https://www.decipher.sanger.ac.uk>). This case report was supported in part by a grant from the Italian Ministry of Health (Ricerca Corrente 2013). MJT is funded by the University of Washington Medical Genetics Postdoctoral Fellowship (5T32 GM007454).

- 5 Kataoka Y, Takeichi M, Uemura T. Developmental roles and molecular characterization of a *Drosophila* homologue of *Arabidopsis* Argonaute1, the founder of a novel gene superfamily. *Genes Cells* 2001; **6**: 313–325.
- 6 Giraldez AJ, Cinalli RM, Glasner ME *et al*: MicroRNAs regulate brain morphogenesis in zebrafish. *Science* 2005; **308**: 833–838.
- 7 Liu J, Carmell MA, Rivas FV *et al*: Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 2004; **305**: 1437–1441.
- 8 Stark KL, Xu B, Bagchi A *et al*: Altered brain microRNA biogenesis contributes to phenotypic deficits in a 22q11-deletion mouse model. *Nat Genet* 2008; **40**: 751–760.
- 9 Forstner AJ, Degenhardt F, Schrott G, Nothen MM. MicroRNAs as the cause of schizophrenia in 22q11.2 deletion carriers, and possible implications for idiopathic disease: a mini-review. *Front Mol Neurosci* 2013; **6**: 47.
- 10 Martinez JE, Tuck-Muller CM, Gasparrini W, Li S, Wertelecki W. 1p microdeletion in sibs with minimal phenotypic manifestations. *Am J Med Genet* 1999; **82**: 107–109.
- 11 Biswas S, Munier FL, Yardley J *et al*: Missense mutations in COL8A2, the gene encoding the alpha2 chain of type VIII collagen, cause two forms of corneal endothelial dystrophy. *Hum Mol Genet* 2001; **10**: 2415–2423.
- 12 Gottsch JD, Sundin OH, Liu SH *et al*: Inheritance of a novel COL8A2 mutation defines a distinct early-onset subtype of Fuchs corneal dystrophy. *Invest Ophthalmol Vis Sci* 2005; **46**: 1934–1939.
- 13 Mok JW, Kim HS, Joo CK. Q455V mutation in COL8A2 is associated with Fuchs corneal dystrophy in Korean patients. *Eye* 2009; **23**: 895–903.
- 14 Meng H, Matthaehi M, Ramanan N *et al*: L450W and Q455K Col8a2 knock-in mouse models of Fuchs endothelial corneal dystrophy show distinct phenotypes and evidence for altered autophagy. *Invest Ophthalmol Vis Sci* 2013; **54**: 1887–1897.
- 15 Schrott G. MicroRNAs at the synapse. *Nat Rev Neurosci* 2009; **10**: 842–849.
- 16 Rio Frio T, Bahubeshi A, Kanellopoulou C *et al*: DICER1 mutations in familial multinodular goiter with and without ovarian Sertoli-Leydig cell tumors. *JAMA* 2011; **305**: 68–77.
- 17 Vorstman JA, Staal WG, van Daalen E, van Engeland H, Hochstenbach PF, Franke L. Identification of novel autism candidate regions through analysis of reported cytogenetic abnormalities associated with autism. *Mol Psychiatry* 2006; **1**: 18–28.
- 18 Zhang F, Gu W, Hurles ME, Lupski JR. Copy number variation in human health, disease, and evolution. *Annu Rev Genomics Hum Genet* 2009; **10**: 451–481.
- 19 Yang JS, Lai EC. Alternative miRNA biogenesis pathways and the interpretation of core miRNA pathway mutants. *Mol Cell* 2011; **43**: 892–903.

- 1 Valencia-Sanchez MA, Liu J, Hannon GJ, Parker R. Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev* 2006; **20**: 515–524.
- 2 Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenines, indicates that thousands of human genes are microRNA targets. *Cell* 2005; **120**: 15–20.
- 3 Meister G. Argonaute proteins: functional insights and emerging roles. *Nat Rev Genet* 2013; **14**: 447–459.
- 4 Yang JS, Lai EC. Dicer-independent, Ago2-mediated microRNA biogenesis in vertebrates. *Cell Cycle* 2010; **9**: 4455–4460.