ARTICLE

EIF3G is associated with narcolepsy across ethnicities

Anja Holm^{1,2}, Ling Lin³, Juliette Faraco³, Sara Mostafavi⁴, Alexis Battle⁴, Xiaowei Zhu⁵, Douglas F Levinson⁵, Fang Han⁶, Steen Gammeltoft¹, Poul Jennum², Emmanuel Mignot³ and Birgitte R Kornum^{*,1}

Type 1 narcolepsy, an autoimmune disease affecting hypocretin (orexin) neurons, is strongly associated with HLA-DQB1*06:02. Among polymorphisms associated with the disease is single-nucleotide polymorphism rs2305795 (c.*638G>A) located within the *P2RY11* gene. *P2RY11* is in a region of syntenic conservation in mammals and zebrafish containing *PPAN*, *EIF3G* and *DNMT1* (DNA methyltransferase 1). As mutations in *DNMT1* cause a rare dominant form of narcolepsy in association with deafness, cerebellar ataxia and dementia, we questioned whether the association with *P2RY11* in sporadic narcolepsy could be secondary to linkage disequilibrium with *DNMT1*. Based on genome-wide association data from two cohorts of European and Chinese ancestry, we found that the narcolepsy association signal drops sharply between *P2RY11/EIF3G* and *DNMT1*, suggesting that the association with narcolepsy does not extend into the *DNMT1* gene region. Interestingly, using transethnic mapping, we identified a novel single-nucleotide polymorphism rs3826784 (c.596–260A>G) in the *EIF3G* gene also associated with narcolepsy. The disease-associated allele increases *EIF3G* mRNA expression. *EIF3G* is located in the narcolepsy risk locus and *EIF3G* expression correlates with *PPAN* and *P2RY11* expression. This suggests shared regulatory mechanisms that might be affected by the polymorphism and are of relevance to narcolepsy.

European Journal of Human Genetics (2015) 23, 1573–1580; doi:10.1038/ejhg.2015.4; published online 11 February 2015

INTRODUCTION

Narcolepsy with cataplexy (Type 1 narcolepsy) affects 1 in 3000 individuals and is caused by the loss of around 70 000 hypocretin (hcrt, also known as orexin) producing neurons in the hypothalamus. The disease is strongly associated with HLA-DQB1*06:02 and DQA1*01:02,¹ the T-cell receptor alpha locus² and polymorphisms in other immune-related genes.³ The loss of hcrt-producing cells is hypothesized to have an autoimmune basis.⁴

One of the genetic associations in narcolepsy is rs2305795, a polymorphism located in the purinergic receptor *P2RY11*.^{5,6} The disease-associated allele of rs2305795 decreases *P2RY11* expression and increases the sensitivity of CD8⁺ T cells to cell death induced by ATP. Interestingly, we also recently found that missense mutation in exon 21 of the DNA methyltransferase 1 (*DNMT1*) gene causes a rare hereditary form of narcolepsy-associated deafness, cerebellar ataxia and eventually dementia (ADCA-DN).⁷ As the *P2RY11* and *DNMT1* genes are located in close proximity (within 18 kb) on chromosome 19, we hypothesized that the two signals could be related at the pathophysiological level. In support of this hypothesis, Kornum *et al.*⁵ observed a correlation between *P2RY11* and *DNMT1* expression in peripheral blood mononuclear cells (PBMCs) of both patients and healthy controls. These findings made us question whether the association with *P2RY11* in spontaneous narcolepsy could be secondary to linkage disequilibrium (LD) with *DNMT1*. In the original study, Kornum *et al.*⁵ fine-mapped the locus using seven single-nucleotide polymorphisms (SNPs) and reported an association between *DNMT1* and narcolepsy in individuals of European ancestry but not Asians. In Asians, LD pattern pointed towards an association with *P2RY11*.⁵

One of the biggest challenges of association studies is to identify which genetic variant is causal within a given locus. To address this challenge, increased sample size, functional studies and transethnic mapping may be beneficial. In transethnic mapping, different LD patterns in the disease-associated locus can be explored to narrow down the associated genetic region. This approach has previously been shown to be very powerful in narcolepsy and other diseases.^{1,8,9} In this study, we reinvestigated SNPs in the *P2RY11-DNMT1* region in narcolepsy across three different ethnic backgrounds. Based on genome-wide association study (GWAS) data available in Chinese and Caucasians, we observed that the narcolepsy association signal drops sharply between *P2RY11* and *DNMT1* in both cohorts. Based on LD patterns, we next selected four SNPs for genotyping in a cohort of African Americans. Following genotyping, we identified a novel SNP in *EIF3G*, rs3826784, best associated with narcolepsy across all three cohorts. *EIF3G* is located between *P2RY11* and *DNMT1*. To further explore association between these different genes and narcolepsy, we finally also examined gene expression patterns in the entire region.

MATERIALS AND METHODS

Subjects

Narcolepsy with cataplexy cases were selected based on documented hypocretin deficiency or clearcut cataplexy and HLA-DQB1*06:02.^{2,6} The initial European ancestry GWA data set was comprised of 807 cases and 1074 DQB1*06:02-positive controls; 415 cases and 753 controls were recruited from the United States and Canada; 392 cases and 321 controls were recruited from European centers. The Chinese GWA data set was comprised of 1078 cases and 1903 controls. Analysis of the GWA data (549 596 SNPs in Europeans and 603 382 SNPs in Chinese) was performed as described.^{2,6} The African-American sample

¹Molecular Sleep Laboratory, Department of Diagnostics, Glostrup University Hospital, Glostrup, Denmark; ²Danish Center for Sleep Medicine, Department of Neurophysiology, Glostrup Hospital, University of Copenhagen, Glostrup, Denmark; ³Center for Sleep Sciences in Medicine and Department of Psychiatry, Stanford University School of Medicine, Palo Alto, CA, USA; ⁴Department of Computer Science, Stanford University, Palo Alto, CA, USA; ⁵Department of Psychiatry and Behavioral Sciences, Stanford University, Palo Alto, CA, USA; ⁶Department of Pulmonary, Critical Care Medicine, Peking University People's Hospital, Beijing, China

*Correspondence: Dr BR Kornum, Molecular Sleep Laboratory, Department of Diagnostics, Glostrup University Hospital, Nordre Ringvej 69, Glostrup 2600, Denmark. Tel: +45 38634541; Fax: +45 38633958; E-mail: birgitte.kornum@regionh.dk

Received 26 February 2014; revised 28 October 2014; accepted 9 December 2014; published online 11 February 2015

contained 1297 individuals (249 cases, 1048 controls). The controls were recruited from the National Institute of Mental Health (NIMH).

For the expression quantitative trait locus (eQTL) analysis, we used a large case-control study of major depression, for which RNA-sequencing data from whole blood together with genotype data were available for 922 individuals of European ancestry.

Imputation of Chinese and European cohorts

Imputation for the *PPAN-DNMT1* region (from rs7250025 to rs2043305) was performed in the European and Chinese cohorts separately, using Beagle v.3.2.1¹⁰ against reference genotypes from the 1000 genomes integrated data set. Chinese genotypes, originally obtained through the Affymetrix Axiom CHP array, were imputed against the CHB reference population. European ancestry genotypes from the Affymetrix 6.0 array were imputed using four European populations as a reference (CEU, TSI, GBR, IBS). In the paper by Kornum *et al.*,⁵ SNPs rs2305795 and rs12460842 were fine typed in many of the Caucasian samples. Further, rs2305795 was also retyped in the Chinese cohort⁶ and all the results were consistent with imputed results. The imputation quality score for rs3826784 was consistently above 0.8 and the overall association on pooled imputed genotypes is considered valid. Therefore, SNP markers with poor imputation quality score in either Chinese or Europeans ($r^2 < 0.8$) were excluded from all further analysis.

The data of this study is submitted to the ClinVar Database (<http://www.ncbi.nlm.nih.gov/clinvar/>; accession numbers: SCV000196702–SCV000196705).

Fine mapping of the chromosome 19 locus, and replication of published SNPs

We used 1297 African-American individuals (249 cases, 1048 controls) for genotyping in the region of association. Based on LD and r^2 data from the International HapMap Project (Hapmap.org), we observed that the LD pattern between the SNPs found from the GWAS was different in African Americans.

The selected SNPs were rs1551570: NM_020230.5:c.190–151C>T; rs12460842: NM_002566.4:c.–203A>G; rs2305795: NM_002566.4:c.*638G>A; and rs3826784: NM_003755.3:c.596–260A>G. For genotyping, we used predesigned Taqman SNP genotyping assays (Applied Biosystems, Carlsbad, CA, USA) to validate the observed LD pattern from the International HapMap project in a cohort of African-American narcolepsy patients and controls. Genotyping was performed at Stanford University. SNPs rs1551750, rs12460842 and rs3826784 are all located within introns of *P2RY11* and *EIF3G*, respectively. SNP rs2305795 is located in the 3'-UTR of *P2RY11*. Genotype results were analyzed (allelic associations, LD calculations and analysis) and standard r^2 plots were generated using Haploview 4.2.¹¹

RNA sequencing and genotyping of 922 individuals of European ancestry

The complete procedure for whole blood processing, RNA sequencing and genotyping is described in Battle *et al.*¹² Briefly, whole blood was collected in PAXGene tubes for RNA and in acid-citrate-dextrose tubes for DNA. Tubes were stored at –80 °C. Total RNA was extracted using the GLOBINclear Kit (Invitrogen, Carlsbad, CA, USA). Libraries for RNA sequencing was performed using Illumina TruSeq protocol. DNA was genotyped on the Illumina HumanOmni1-Quad BeadChip (Illumina, San Diego, CA, USA).

Statistical analysis

Genome-wide association analysis of the European and Chinese ancestry has been described previously.^{2,6} Statistical analysis of genotyped and imputed genotype data were performed using the PLINK software package (v.1.07) (<http://pngu.mgh.harvard.edu/purcell/plink/>),¹³ and in Haploview 4.2.¹¹

Associations within ethnic groups were performed with basic allelic χ^2 statistic tests. For associations combining multiple ethnic groups, the Mantel–Haenszel test was used together with the Breslow Day test of homogeneity of the odds ratio (both implemented in Plink).

The details of the *cis*-eQTL analysis are described in Battle *et al.*¹² Briefly, expression data was first normalized to remove the effect of technical factors,

population structure and 'hidden' factors (which have been shown to reduce the statistical power to identify *cis*-eQTLs¹⁴). Such normalization approaches can significantly increase power for detecting *cis*-eQTLs; however, in doing so, such approaches also remove broader expression patterns that may be driven by *trans*-regulation (as a *trans*-acting factor may affect the expression levels of multiple target genes) or reflect coregulation. As a result, previous work has shown that separate normalization of gene expression data may be appropriate for detecting *cis* versus *trans* associations.^{12,14} Therefore, we quantified co-expression levels between genes using the normalization approach optimized for detecting *trans*-eQTLs. For each SNP, a *P*-value for association with gene expression level of each nearby gene (1 Mb from the transcription start site) was computed based on the Spearman's correlation coefficient. Significant associations were reported using 0.05 FDR (false discovery rate).

PPAN: NM_020230.4; *P2RY11*: NM_002566.4; *EIF3G*: NM_003755.3; *DNMT1*: NM_001130823.1.

Rs1551570: NM_020230.5:c.190–151C>T; rs12460842: NM_002566.4:c.–203A>G; rs2305795: NM_002566.4:c.*638G>A; rs3826784: NM_003755.3:c.596–260A>G.

SNPs reported in Table 1: rs1551570: NM_020230.5:c.190–151C>T; rs12460842: NM_002566.4:c.–203A>G; rs2010353: NM_002566.4:c.19+142G>T; rs3826784: NM_003755.3:c.596–260T>C; rs2305795: NM_002566.4:c.*638G>A; rs12462506: NM_002566.4:c.20–178T>G; rs3826785: NM_003755.3:c.595+327G>A; rs73011220: NM_002566.4:c.19+880G>A; rs112647895: NM_002566.4:c.19+446G>A; rs3745601: NM_002566.4:c.259G>A; rs2305789: NM_020230.5:c.291+50A>G; rs6511570: NM_001130823.1:c.2895–855C>T; rs2288349: NM_001130823.1:c.2721–45C>T; rs35693490: NM_001130823.1:c.2721–959G>A; rs8101626: NM_001130823.1:c.4773+383C>T; rs4804122: NC_000019.10:g.10131268C>T; 9 rs2114724: NM_001130823.1:c.1832+14A>G; rs2290684: NM_001130823.1:c.3394+34T>C; rs8112801: NM_001130823.1:c.3117–203A>C; rs11880553: NM_001130823.1:c.3117–660G>A; rs11880388: NM_001130823.1:c.3117–677C>T; rs11085587: NC_000019.10:g.10124423G>C; rs35374357: NC_000019.10:g.10129267T>C; rs7710: NM_003755.3:c.846T>C; rs870612: NC_000019.10:g.10120814A>C; rs4555265: NC_000019.10:g.10121508G>A; rs11667630: NM_003755.3:c.300+47G>T; rs2290687: NM_003755.3:c.240+23G>A; rs55752217: NM_020230.5:c.292–46C>T; rs2305792: NM_020230.5:c.902–10C>G; rs1037686: NM_002566.4:c.19+419T>A; rs3745600: NM_002566.4:c.237C>T; rs34484805: NM_002566.4:c.20–882A>G; rs7401: NM_002566.4:c.*363A>G; rs11666402: NM_020230.5:c.513+266T>C; rs10414661: NC_000019.10:g.10131561T>A; rs12462004: NC_000019.10:g.10127588T>G; rs10404209: NC_000019.10:g.10126751T>C; rs9305012: NM_001130823.1:c.2266–225A>G; rs10407514: NM_001130823.1:c.2895–449G>C; rs10418707: NM_001130823.1:c.2894+810C>T; rs12611113: NM_001130823.1:c.2721–212A>G; rs60565702: NM_001130823.1:c.2720+999C>T; rs72620548: NM_001130823.1:c.4490–306C>T; rs10854076: NM_001130823.1:c.4293+238C>G; rs4804489: NM_001130823.1:c.2265+418T>C; rs57366074: NM_001130823.1:c.2586+99G>T; rs10414537: NC_000019.10:g.10126048C>T.

RESULTS

Association signal within the risk locus

Using two GWA data sets from narcoleptic patients and controls of European and Chinese ancestry, we investigated polymorphisms in the *P2RY11-DNMT1* region. The cohorts included 1881 individuals of European ancestry (807 cases and 1074 controls) and 2981 Chinese (1078 cases and 1903 controls). SNPs in the region of interest were imputed in each GWA data set, and ranked by statistical significance in the Chinese cohort (Table 1). The most significant SNPs in both GWA data sets are in high LD with rs2305795 and located in the *PPAN-P2RY11-EIF3G* haplotype block, and not in *DNMT1* (Table 1). R^2 plots clearly illustrate that SNPs in high LD with rs2305795 are restricted to the *PPAN-P2RY11-EIF3G* region, but not *DNMT1* (Figure 1), as was also observed by Han *et al.*⁶ Based on this analysis, five SNP markers (rs1551570 (*PPAN*), rs12460842 (*P2RY11*), rs2010353 (*P2RY11*), rs2305795 (*P2RY11*) and rs3826784 (*EIF3G*))

Table 1 SNP markers located in the PPAN-P2RY11-EIF3G-DNMT1 locus and their association with Type 1 narcolepsy

Gene	SNP	Position (bp)	European						Chinese									
			Major allele	Minor allele	Freq cases	Freq cont.	P-value	OR (95% CI)	Imp. quality	r ²	Major allele	Minor allele	Freq cases	Freq cont.	P-value	OR (95% CI)	Imp. quality	r ²
PPAN	Rs1551570	10 218 030	T	C	0.42	0.48	4.16E-04	0.79 (0.69-0.90)	0.84	0.98	T	C	0.25	0.30	1.20E-05	0.77 (0.68-0.86)	1.00	0.94
	Rs12460842	10 222 195	G	A	0.42	0.48	3.45E-04	0.79 (0.69-0.90)	0.86	0.99	G	A	0.25	0.30	1.47E-05	0.77 (0.68-0.87)	0.98	0.94
P2RY11	Rs2010353	10 222 558	T	G	0.42	0.48	3.45E-04	0.79 (0.69-0.90)	0.86	0.99	T	G	0.25	0.30	1.47E-05	0.77 (0.68-0.87)	0.97	0.94
EIF3G	Rs3826784	10 227 010	G	A	0.38	0.44	1.28E-04	0.77 (0.68-0.88)	0.84	0.87	G	A	0.25	0.30	1.47E-05	0.77 (0.68-0.87)	0.93	0.94
P2RY11	Rs2305795	10 226 052	A	G	0.42	0.48	2.31E-04	0.78 (0.69-0.89)	0.89	1.00	A	G	0.26	0.31	5.99E-05	0.79 (0.70-0.88)	0.93	1.00
P2RY11	Rs12462506	10 224 131	T	G	0.14	0.11	4.72E-03	1.33 (1.09-1.61)	0.87	0.11	T	G	0.46	0.42	2.23E-03	1.18 (1.06-1.31)	0.95	0.28
EIF3G	Rs3826785	10 227 149	C	T	0.14	0.11	3.55E-03	1.33 (1.10-1.62)	0.99	0.11	C	T	0.44	0.40	4.06E-03	1.17 (1.05-1.30)	0.94	0.31
P2RY11	Rs73011220	10 222 862	G	A	0.14	0.11	4.72E-03	1.33 (1.10-1.63)	0.97	0.11	G	A	0.44	0.40	4.68E-03	1.17 (1.05-1.30)	0.97	0.31
P2RY11	Rs112647895	10 223 296	G	A	0.14	0.11	3.63E-03	1.34 (1.09-1.61)	0.94	0.11	G	A	0.44	0.40	4.68E-03	1.17 (1.05-1.30)	0.97	0.31
P2RY11	Rs3745601	10 224 548	G	A	0.14	0.11	4.72E-03	1.33 (1.09-1.61)	0.96	0.11	G	A	0.44	0.40	4.68E-03	1.17 (1.05-1.30)	0.95	0.31
PPAN	Rs2305789	10 218 332	A	G	0.14	0.11	3.63E-03	1.34 (1.10-1.63)	0.84	0.11	A	G	0.44	0.41	7.40E-03	1.16 (1.04-1.29)	1.00	0.31
DNMT1	Rs6511570	10 246 029	A	G	0.39	0.45	2.35E-04	0.78 (0.68-0.89)	0.98	0.62	A	G	0.24	0.27	2.29E-02	0.87 (0.77-0.98)	0.97	0.51
DNMT1	Rs2288349	10 255 518	G	A	0.39	0.45	1.88E-04	0.78 (0.69-0.89)	0.98	0.61	G	A	0.24	0.27	2.29E-02	0.87 (0.77-0.98)	0.97	0.51
DNMT1	Rs35693490	10 257 245	G	A	0.39	0.45	1.88E-04	0.78 (0.68-0.89)	0.95	0.61	G	A	0.24	0.27	2.29E-02	0.87 (0.77-0.98)	0.97	0.51
DNMT1	Rs8101626	10 258 159	C	T	0.39	0.45	1.88E-04	0.78 (0.68-0.89)	0.97	0.61	C	T	0.24	0.27	2.29E-02	0.87 (0.77-0.98)	0.92	0.51
Intergenic	Rs4804122	10 241 944	T	C	0.40	0.46	4.78E-05	0.76 (0.67-0.87)	0.97	0.65	T	C	0.26	0.28	2.78E-02	0.87 (0.78-0.99)	0.96	0.55
DNMT1	Rs2114724	10 265 248	T	C	0.46	0.50	9.66E-03	0.84 (0.74-0.96)	0.99	0.50	C	T	0.30	0.29	1.79E-01	1.08 (0.96-1.22)	1.00	0.10
DNMT1	Rs2290684	10 251 747	G	A	0.46	0.51	1.02E-02	0.84 (0.74-0.96)	0.98	0.50	A	G	0.30	0.29	1.84E-01	1.08 (0.96-1.21)	0.99	0.10
DNMT1	Rs8112801	10 253 099	G	T	0.46	0.51	1.02E-02	0.84 (0.74-0.96)	0.98	0.50	T	G	0.30	0.29	1.84E-01	1.08 (0.96-1.21)	0.99	0.10
DNMT1	Rs11880553	10 253 556	T	C	0.46	0.51	1.02E-02	0.84 (0.74-0.96)	0.98	0.50	C	T	0.30	0.29	1.84E-01	1.08 (0.96-1.21)	0.99	0.10
DNMT1	Rs11880388	10 253 573	A	G	0.46	0.51	1.02E-02	0.84 (0.74-0.96)	0.97	0.50	G	A	0.30	0.29	1.84E-01	1.08 (0.96-1.21)	0.99	0.10
Intergenic	Rs11085587	10 235 099	C	G	0.46	0.51	8.90E-03	0.84 (0.74-0.96)	0.98	0.50	G	C	0.31	0.29	1.96E-01	1.08 (0.96-1.21)	0.91	0.10
Intergenic	Rs35374357	10 239 943	T	C	0.43	0.41	1.69E-01	1.10 (0.96-1.25)	0.91	0.35	T	C	0.30	0.29	2.08E-01	1.08 (0.96-1.21)	0.94	0.10
EIF3G	Rs7710	10 226 256	A	G	0.44	0.41	6.65E-02	1.13 (0.99-1.29)	0.88	0.63	A	G	0.31	0.29	2.24E-01	1.07 (0.96-1.21)	0.98	0.13
Intergenic	Rs870612	10 229 297	C	A	0.45	0.42	5.15E-02	1.14 (1.00-1.30)	0.90	0.62	C	A	0.29	0.28	3.09E-01	1.06 (0.95-1.19)	1.00	0.16
Intergenic	Rs4555265	10 229 521	C	T	0.45	0.42	5.51E-02	1.14 (1.00-1.30)	0.88	0.62	C	T	0.29	0.28	3.09E-01	1.06 (0.95-1.19)	1.00	0.16
EIF3G	Rs11667630	10 231 490	A	C	0.45	0.42	5.15E-02	1.14 (1.00-1.30)	0.90	0.62	A	C	0.29	0.28	3.09E-01	1.06 (0.95-1.19)	1.00	0.16
EIF3G	Rs2290687	10 232 184	G	A	0.45	0.42	5.15E-02	1.14 (1.00-1.29)	0.90	0.62	G	A	0.29	0.28	3.09E-01	1.06 (0.95-1.19)	1.00	0.16
PPAN	Rs55752217	10 218 434	C	T	0.44	0.41	1.02E-01	1.12 (0.98-1.27)	0.88	0.62	C	T	0.29	0.28	3.24E-01	1.06 (0.94-1.19)	0.99	0.16
PPAN	Rs2305792	10 221 151	C	G	0.44	0.41	9.99E-02	1.12 (0.98-1.27)	0.87	0.62	C	G	0.29	0.28	3.33E-01	1.06 (0.94-1.19)	0.99	0.16
P2RY11	Rs1037686	10 222 835	T	A	0.44	0.42	9.25E-02	1.12 (0.98-1.27)	0.87	0.62	T	A	0.29	0.28	3.33E-01	1.06 (0.94-1.19)	0.99	0.16
P2RY11	Rs3745600	10 223 427	A	G	0.44	0.41	4.67E-02	1.14 (0.99-1.28)	0.88	0.62	A	G	0.29	0.28	3.33E-01	1.06 (0.94-1.19)	0.98	0.16
P2RY11	Rs344484805	10 224 526	C	T	0.44	0.41	7.70E-02	1.13 (1.00-1.30)	0.88	0.62	C	T	0.29	0.28	3.33E-01	1.06 (0.94-1.19)	0.98	0.16
P2RY11	Rs7401	10 225 777	A	G	0.44	0.41	4.27E-02	1.14 (1.00-1.30)	0.87	0.62	A	G	0.29	0.28	3.33E-01	1.06 (0.94-1.19)	0.98	0.16
PPAN	Rs11666402	10 219 076	T	C	0.44	0.42	9.43E-02	1.12 (0.98-1.27)	0.90	0.62	T	C	0.29	0.28	3.52E-01	1.06 (0.94-1.19)	0.91	0.16
Intergenic	Rs10414661	10 237 427	T	C	0.07	0.06	3.21E-02	1.32 (0.99-1.66)	0.98	0.01	T	C	0.44	0.43	4.11E-01	1.05 (0.94-1.16)	0.97	0.15
Intergenic	Rs12462004	10 238 264	T	G	0.08	0.06	3.73E-02	1.31 (1.02-1.70)	0.94	0.01	T	G	0.44	0.43	4.11E-01	1.05 (0.94-1.16)	0.94	0.15
Intergenic	Rs10404209	10 242 237	T	A	0.08	0.06	6.20E-02	1.28 (1.02-1.71)	0.95	0.01	T	A	0.44	0.43	4.11E-01	1.05 (0.94-1.16)	0.94	0.15
DNMT1	Rs9305012	10 255 112	C	G	0.07	0.06	2.17E-02	1.36 (1.03-1.73)	0.93	0.02	C	G	0.46	0.45	4.35E-01	1.04 (0.94-1.16)	0.97	0.12
DNMT1	Rs10407514	10 260 869	T	C	0.07	0.06	2.99E-02	1.33 (1.05-1.76)	1.00	0.02	T	C	0.46	0.45	4.35E-01	1.04 (0.94-1.16)	0.97	0.12
DNMT1	Rs10418707	10 247 269	G	A	0.07	0.06	2.17E-02	1.36 (1.05-1.76)	1.00	0.02	G	A	0.44	0.43	4.48E-01	1.04 (0.94-1.16)	0.98	0.15

Table 1 (Continued)

Gene	SNP	Position (bp)	European						Chinese									
			Major allele	Minor allele	Freq cases	Freq cont.	P-value	OR (95% CI)	Imp. quality	r ²	Major allele	Minor allele	Freq cases	Freq cont.	P-value	OR (95% CI)	Imp. quality	r ²
<i>DNMT1</i>	Rs12611113	10 248 270	G	C	0.07	0.06	2.17E-02	1.36 (1.05–1.76)	1.00	0.02	G	C	0.44	0.43	4.48E-01	1.04 (0.94–1.16)	0.98	0.15
<i>DNMT1</i>	Rs60565702	10 256 217	G	A	0.07	0.06	2.17E-02	1.36 (1.05–1.76)	0.94	0.02	G	A	0.44	0.43	4.48E-01	1.04 (0.94–1.16)	0.98	0.15
<i>DNMT1</i>	Rs72620548	10 257 412	T	C	0.07	0.06	2.17E-02	1.36 (1.05–1.76)	0.99	0.02	T	C	0.44	0.43	4.48E-01	1.04 (0.94–1.16)	0.98	0.15
<i>DNMT1</i>	Rs10854076	10 258 561	G	A	0.07	0.06	2.17E-02	1.36 (1.05–1.76)	0.97	0.02	G	A	0.44	0.43	4.48E-01	1.04 (0.94–1.16)	0.98	0.15
<i>DNMT1</i>	Rs4804489	10 260 030	C	A	0.07	0.06	2.55E-02	1.35 (1.05–1.76)	0.98	0.02	C	A	0.44	0.43	4.48E-01	1.04 (0.94–1.16)	0.97	0.15
<i>DNMT1</i>	Rs57366074	10 261 656	A	G	0.07	0.06	2.17E-02	1.36 (1.04–1.75)	0.99	0.02	A	G	0.44	0.43	4.48E-01	1.04 (0.94–1.16)	0.92	0.15
<i>Intergenic</i>	Rs10414537	10 236 724	C	T	0.07	0.06	6.20E-02	1.28 (0.99–1.66)	0.95	0.01	C	T	0.42	0.41	4.74E-01	1.04 (0.93–1.16)	0.87	0.18

Abbreviations: bp, base pair; CI, confidence interval; freq., frequency; imp., imputation; OD, odds ratio; r², pairwise r² with rs2305795; SNP, single-nucleotide polymorphism. P-values and imputation quality were calculated with Beagle. Pairwise LD (r²) with rs2305795 was calculated with Haploview. Bold letters indicate SNPs of interest. Only SNP marker with an imputation quality score of r² > 0.8 were included. Intergenic: SNP markers located between *EIF3G* and *DNMT1*, which is ranked by statistical significance in the Chinese cohort. PPA: NM_020230.4; P2RY11: NM_002566.4; *EIF3G*: NM_003755.3; *DNMT1*: NM_001130823.1.

were significantly associated with narcolepsy in both cohorts, and were selected for further analysis. Owing to a failure of the genotyping assay, however, rs2010353 was excluded, leaving four SNPs for all subsequent analyses.

Based on data from the International HapMap Project, we next explored LD patterns for the four markers across ethnic groups. Interestingly, two SNPs rs12460842 (*P2RY11*) and rs3826784 (*EIF3G*) had a lower LD with rs1551570 (*PPAN*) and rs2305795 (*P2RY11*) in an African-American cohort (Figure 2), suggesting these to be candidates for transethnic mapping. Genotyping of these four markers in 1297 African Americans (249 cases and 1048 controls) showed that surprisingly rs3826784, and not the previously reported rs2305795, showed the strongest narcolepsy association based on odds ratio (OR) values, although none of the four SNPs reached statistical significance in this data set. A meta-analysis was performed across all ethnic groups and here we replicated the finding by Kornum *et al.*⁵ of an increased OR with the major allele rs2305795A (Table 2 shows data on minor allele). The meta-analysis further revealed that rs3826784 showed the strongest association with narcolepsy across all ethnic groups (Table 2) with a slightly higher OR for the rs3826784G risk allele and a lower P-value. There were no significant differences between the ORs from the four SNPs.

Expression quantitative trait loci analysis

To determine if rs3826784 regulates gene expression, we next performed an eQTL analysis of the genes in the risk locus. For this analysis, we used a large case-control study of major depression, for which RNA-sequencing data from whole blood together with genotype data were available for 922 individuals of European ancestry.¹² After appropriate normalization, whether or not cases or controls were used it did not influence the eQTL analysis;¹² thus, we performed all analysis in the combined sample.

Using this data set, we found that the expression of *EIF3G* increased with the risk allele, *DNMT1* expression decreased and the expression of *PPAN* and *P2RY11* did not change with the risk allele (rs3826784G) (Figure 3). Furthermore, the expression of *EIF3G* and *DNMT1*, but not *PPAN* and *P2RY11*, correlated with rs3826784, the strongest association being with *EIF3G* (Table 3). The other four SNPs in our analysis showed equally strong association with *EIF3G* consistent with the high LD between these four SNPs in Caucasians (Figure 2). The expression of *DNMT1* was most strongly correlated with eQTL SNP for *DNMT1* rs2290684 (highlighted in Table 1). The effect of rs3826784 on *DNMT1* expression is therefore likely explained by LD between rs3826784 and rs2290684 (r² = 0.35). In line with this, conditional analysis showed that rs3826784 does not independently affect *DNMT1* expression when the effect of rs2290684 is controlled for (P > 0.1).

The effect of rs2305795 on *P2RY11* was also re-examined. Using our data, we found that rs2305795 was not an eQTL for *P2RY11* (see Supplementary Table S1). In addition, as rs2305795 is located in the 3'-UTR of *P2RY11*, it was possible to analyze allele-specific association for this SNP. Using a test of association between rs2305795 and the allelic ratio of *P2RY11* (as an eQTL), we did not observe a significant association between rs2305795 and the allelic expression of *P2RY11* in the human whole blood data set in contrast to earlier reports in PBMCs.⁵

Gene expression correlation

We finally investigated the possibility of coregulation and/or regulatory relationships between the four genes in our locus by analyzing correlations between expression of *PPAN*, *P2RY11*, *EIF3G* and

Figure 1 Risk locus on Chr. 19q13.2. Gene organization and LD in the region of interest (10 209 865–10 276 167) compared between European (CEU) and Chinese ancestry (CHB). Top: R^2 -based LD plots using data from GWA data set of European ancestry (CEU). Bottom: R^2 -based LD plots using data from GWA data set of Chinese ancestry (CHB). Darker colors illustrate higher levels of LD. Dashed and dotted line indicates the position of rs2305795 and rs3826784, respectively.

Figure 2 R^2 -based LD plots for individuals of CEU, CHB and ASW. Utah residents with Northern and Western European ancestry from the CEPH collection (CEU), Han Chinese in Beijing, China (CHB) and African ancestry in Southwest USA (ASW). Curly braces indicate LD values between SNP candidates: rs1551570, rs12360842, rs2305795 and rs3826784. Darker color illustrates a higher level of LD. Chromosome 19q13.2 (10 209 865–10 230 599). See Figure 1 for entire locus.

DNMT1. *PPAN* and *P2RY11* were found to be expressed at very low levels overall in whole blood cells, which correlate with the expression in the brain (Supplementary Tables S2+S3). Expression of *EIF3G* correlated with the expression of *PPAN* in normalized data optimized

for detecting *trans*-eQTLs but not in data optimized for detecting *cis*-eQTLs (for more information see Materials and methods). Finally, the expression level of *EIF3G* was not correlated with *DNMT1* (Table 4). We also did not observe a significant correlation between *P2RY11* and

Table 2 Association of SNP markers with Type 1 narcolepsy

SNP	Position	Major allele > minor allele	Freq. cases (807)	Freq. controls (1074)	P-value (MH)	OR (95% CI)	Pairwise r^2 with rs2305795	
<i>European ancestry (1881)</i>								
Rs1551570	10 218 030	T > C	0.42	0.48	4.00E-04	0.79 (0.69–0.90)	0.98	
Rs12460842	10 222 195	G > A	0.42	0.48	3.00E-04	0.79 (0.69–0.90)	0.99	
Rs2305795	10 226 052	A > G	0.42	0.48	2.00E-04	0.78 (0.69–0.90)	1	
Rs3826784	10 227 010	G > A	0.38	0.44	1.00E-04	0.77 (0.68–0.88)	0.87	
SNP	Position	Major allele > minor allele	Freq. cases (1078)	Freq. controls (1903)	P-value (MH)	OR (95% CI)	Pairwise r^2 with rs2305795	
<i>Chinese ancestry (2981)</i>								
Rs1551570	10 218 030	T > C	0.25	0.30	1.20E-05	0.77 (0.68–0.86)	0.94	
Rs12460842	10 222 195	G > A	0.25	0.30	1.47E-05	0.77 (0.68–0.87)	0.94	
Rs2305795	10 226 052	A > G	0.26	0.31	5.99E-05	0.79 (0.70–0.88)	1	
Rs3826784	10 227 010	G > A	0.25	0.30	1.47E-05	0.77 (0.68–0.87)	0.94	
SNP	Position	Major allele > minor allele	Freq. cases (249)	Freq. controls (1048)	P-value (MH)	OR (95% CI)	Pairwise r^2 with rs2305795	
<i>African Americans (1297)</i>								
Rs1551750	10 218 030	T > C	0.37	0.37	8.90E-01	0.99 (0.80–1.21)	0.65	
Rs12460842	10 222 195	G > A	0.52	0.54	6.20E-01	0.95 (0.86–1.28)	0.47	
Rs2305795	10 226 052	A > G	0.39	0.40	7.20E-01	0.96 (0.79–1.18)	1	
Rs3826784	10 227 010	G > A	0.43	0.46	1.90E-01	0.87 (0.71–1.07)	0.44	
SNP	Position	Major allele > minor allele	Freq. cases (2134)	Freq. controls (4025)	P-value (MH)	OR (95% CI)	CHISQ_BD	P_BD
<i>All samples (6159)</i>								
Rs1551750	10 218 030	T > C	0.33	0.37	1.97E-07	0.81 (0.74–0.88)	4.472	0.1069
Rs12460842	10 222 195	G > A	0.34	0.41	1.14E-07	0.80 (0.74–0.87)	3.397	0.1829
Rs2305795	10 226 052	A > G	0.34	0.38	2.94E-07	0.81 (0.75–0.88)	3.33	0.1892
Rs3826784	10 227 010	G > A	0.32	0.38	6.35E-09	0.79 (0.73–0.85)	1.246	0.536

Abbreviations: BD, Breslow day; CI, confidence interval; Freq., frequency; MH, Mantel-Haenszel; OR, odds ratio; SNP, single-nucleotide polymorphism. P-values were calculated with Beagle. Pairwise LD (r^2) with rs2305795 was calculated with Haploview. Results for minor allele (for European and Chinese ancestry) are presented for each cohort. The selected SNPs were: rs1551570:T>G), rs12460842:A>G), rs2305795:G>A) and rs3826784:A>G). The most significant SNP across all samples is indicated in bold. PPAN: NM_020230.4; P2RY11: NM_002566.4; EIF3G: NM_003755.3; DNMT1: NM_001130823.1.

DNMT1 in the normalized data for detecting *cis*-eQTL or *trans*-eQTL in contrast to what has earlier been reported for PBMCs.⁵

DISCUSSION

In this study, we found rs3826784, a SNP located within an intron of the *EIF3G* gene, to be the most significantly associated SNP of the region. Transethnic mapping further revealed that rs3826784 could be a better marker for the genetic association with narcolepsy than the previously identified rs2305795. However, the magnitude of the association is very similar to that of the previously reported SNP rs2305795.⁵ The new SNP, rs3826784, is in high LD with SNPs located in *PPAN* and *P2RY11*, but not in *DNMT1*. Genetic LD between *P2RY11* and *DNMT1* is weak in narcolepsy and control cohorts, confirming the observation by Han *et al.*⁶ Importantly, however, elements located within the *P2RY11* gene may still regulate *DNMT1* expression. We also found that the expression of *EIF3G* increased with the risk allele, *DNMT1* expression decreased and there was no effect of rs3826784G on *PPAN* or *P2RY11* mRNA levels in whole blood. The results suggest that the narcolepsy-associated SNP in this locus affects *EIF3G* expression rather than *P2RY11*.

To gain insight into gene regulation in the entire risk locus, we investigated gene expression correlations. In whole blood, expression

of *EIF3G* strongly correlated with the expression of *PPAN* and *P2RY11*, and less with *DNMT1*. These results were consistent with strong LD found with SNPs located in the *EIF3G*, *P2RY11* and *PPAN* region, an effect not extending to *DNMT1*. Surprisingly, however, expression of *P2RY11* and *DNMT1* did not correlate as earlier discovered in PBMCs.⁵ *PPAN* and *P2RY11* were found to be expressed at very low levels in whole blood cells. The discrepancy in *P2RY11* expression is most likely caused by the different input material in the two studies. Kornum *et al.*⁵ showed that *P2RY11* expression was predominantly seen in CD8⁺ cells. These account for ~20% of PBMCs. However, whole blood also contains granulocytes, which accounts for 65% of all cells, and may thus mask the expression of *P2RY11* in the CD8⁺ cells. In agreement with our results, it has previously been reported that *EIF3G* is highly expressed in blood cells.¹⁵ These findings illustrate the complexity of interpreting eQTL studies in whole blood, considering heterogeneity of the effect of regulatory regions on gene expression in various white blood cell subsets. Indeed, cell-specific regulatory mechanisms in the *PPAN*-*P2RY11*-*EIF3G* region are likely present. Future studies will need to address the regulation of the *PPAN*-*P2RY11*-*EIF3G* narcolepsy risk locus within distinct sub-populations of white blood cells. That cell-specific effects are present is not surprising considering the importance

Figure 3 Gene expression in whole blood with various rs3826784 genotypes. (a and b) The expression of *PPAN* and *P2RY11* did not change with the risk allele (rs3826784G). (c) The expression of *EIF3G* increased with the risk allele (rs3826784GG). (d) The expression of *DNMT1* decreased with the risk allele (rs3826784GG). The data are reported as normalized effect size, which equals the correlation coefficient (Spearman's ρ correlation coefficient is used as a measure of change in expression level on the y axis).

Table 3 Association between SNPs and eQTLs in the Type 1 narcolepsy risk locus (negative log₁₀ *P*-values for association between genotype and expression levels)

Gene/SNP name	<i>Rs12460842</i> (genotyped)	<i>Rs2305795</i> (genotyped)	<i>Rs3826784</i> (imputed)	<i>Rs1551570</i> (genotyped)
<i>PPAN</i>	0.94	1.01	1.10	1.00
<i>P2RY11</i>	0.14	0.22	0.27	0.18
<i>EIF3G</i>	42.94	43.95	38.83	43.25
<i>DNMT1</i>	10.48	10.86	9.10	10.12

PPAN: NM_020230.4; *P2RY11*: NM_002566.4; *EIF3G*: NM_003755.3; *DNMT1*: NM_001130823.1.

Table 4 Correlation coefficient between gene expression levels of each gene pair was measured using the Spearman's correlation coefficient

Correlation coefficient					Negative log10 P-value				
	PPAN	P2RY11	EIF3G	DNMT1		PPAN	P2RY11	EIF3G	DNMT1
PPAN		0.276	0.213	0.064	PPAN		16.75	10.10	1.28
P2RY11	0.276		0.155	0.008	P2RY11	16.75		5.64	0.09
EIF3G	0.212	0.155		0.050	EIF3G	10.10	5.64		0.88
DNMT1	0.064	0.008	0.050		DNMT1	1.28	0.09	0.88	

PPAN: NM_020230.4; *P2RY11*: NM_002566.4; *EIF3G*: NM_003755.3; *DNMT1*: NM_001130823.1.

of the regulation of apoptosis in various immune cells types in reaction to changes in the environmental milieu.

How could *EIF3G* be involved in narcolepsy? Eukaryotic initiation factors (EIF) have an important role in the initiation phase of eukaryotic translation by forming a complex with the 40S ribosomal subunit and recruiting a Met-tRNA_i called the 43S to the preinitiation complex. This complex recognizes and binds the 5' cap structure of mRNA, promoting ribosomal scanning of mRNA and regulating recognition of the AUG initiation codon.^{16,17} In mammals, EIF3 is the largest scaffolding initiation factor (750 kDa), comprising 13 subunits (A–M). These subunits cluster together in subcomplexes, in which EIF3G connects to EIF3B and EIF3I. EIF3B:G contains RNA recognition motif subunits that are involved in RNA binding. Interestingly, EIF3B:G:I also interacts directly with certain viral mRNAs to promote the translation of viral proteins.¹⁸ EIF4G, which interacts with EIF3,¹⁹ contains a binding domain for the non-structural influenza protein, NS1,²⁰ and is recruited to the viral mRNA to initiate translation of influenza virus. This is particularly interesting as recent findings have implicated a role of the H1N1 pandemic virus of 2009 as a trigger for narcolepsy development.⁶ It has been suggested that the disruption of EIF pathways in childhood H1N1/09 influenza increases disease severity,²¹ so polymorphisms in this locus could indirectly change immune responses to influenza and narcolepsy risk.

Another possibility could involve regulation of cell death, a function that could be shared with *P2RY11* and *DNMT1*, with coordinated regulation of these loci explaining the conservation of synteny. The N terminus of EIF3G indeed interacts with the C-terminal region of apoptosis-inducing factor (AIF). AIF is a ubiquitous FAD-binding flavoprotein that has an important role in caspase-independent apoptosis. Mature AIF inhibits newly translated protein synthesis via its interactions with EIF3G. Additionally, mature AIF overexpression specifically activates caspase-7, thereby amplifying the inhibition of protein synthesis including EIF3G cleavage.^{22,23} Additionally, *EIF3G* is expressed in the brain and the lateral hypothalamus (Allen Brain Atlas: www.brain-map.org).¹⁵ Higher EIF3G expression with the narcolepsy risk allele could cause a dysregulated cell death, which again could have an important role in white blood cells in the context of autoimmunity, or in the brain in the context of hypocretin cell loss.

In conclusion, using transethnic mapping, we found a novel genetic association of Type 1 narcolepsy with rs3826784, an SNP located in *EIF3G*. The new SNP is in high LD with the previously reported *P2YR11* rs2305795 SNP in individuals of European and Chinese ancestry, but displays lower LD in African Americans. As the association signal from rs3826784 is nominally better compared with rs2305795 in all three cohorts, rs3826784 may be a better marker for the genetic association with narcolepsy, although this conclusion is tentative considering sample size of the African-American Cohort. Further confirmation is needed through extended GWA studies that combine multiple ethnic groups, and local resequencing. A complex correlation between the expression of three genes in the region suggests that a shared regulatory mechanism exists and might be affected by the polymorphism.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We are indebted to all the narcoleptic patients of the study. Without their contributions this study would not have been possible. This work was supported by the Lundbeck Foundation and NIH Grant P50 NS23724.

- Matsuki K, Grumet FC, Lin X *et al*: DQ (rather than DR) gene marks susceptibility to narcolepsy. *Lancet* 1992; **339**:1052.
- Hallmayer J, Faraco J, Lin L *et al*: Narcolepsy is strongly associated with the T-cell receptor alpha locus. *Nat Genet* 2009; **41**: 708–711.
- Faraco J, Lin L, Kornum BR *et al*: ImmunoChip study implicates antigen presentation to T cells in narcolepsy. *PLoS Genet* 2013; **9**: e1003270.
- Partinen M, Kornum BR, Piazzi G *et al*: Narcolepsy as an autoimmune disease: the role of H1N1 infection and vaccination. *Lancet Neurol* 2014; **13**: 600–613.
- Kornum BR, Kawashima M, Faraco J *et al*: Common variants in P2RY11 are associated with narcolepsy. *Nat Genet* 2011; **43**: 66–71.
- Han F, Faraco J, Dong XS *et al*: Genome wide analysis of narcolepsy in China implicates novel immune loci and reveals changes in association prior to versus after the 2009 H1N1 influenza pandemic. *PLoS Genet* 2013; **9**: e1003880.
- Winkelmann J, Lin L, Schormair B *et al*: Mutations in DNMT1 cause autosomal dominant cerebellar ataxia, deafness and narcolepsy. *Hum Mol Genet* 2012; **21**: 2205–2210.
- Mignot E, Lin L, Rogers W *et al*: Complex HLA-DR and -DQ interactions confer risk of narcolepsy-cataplexy in three ethnic groups. *Am J Hum Genet* 2001; **68**: 686–699.
- Wu Y, Waite LL, Jackson AU *et al*: Trans-ethnic fine-mapping of lipid loci identifies population-specific signals and allelic heterogeneity that increases the trait variance explained. *PLoS Genet* 2013; **9**: e1003379.
- Browning SR, Browning BL: Rapid and accurate haplotype phasing and missing data inference for whole genome association studies by use of localized haplotype clustering. *Am J Hum Genet* 2007; **81**: 1084–1097.
- Barrett JC, Fry B, Maller J *et al*: Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 2005; **21**: 263–265.
- Battle A, Mostafavi S, Zhu X *et al*: Characterizing the genetic basis of transcriptome diversity through RNA-sequencing of 922 individuals. *Genome Res* 2014; **24**: 14–24.
- Purcell S, Neale B, Todd-Brown K *et al*: PLINK: a toolset for whole-genome association and population-based linkage analysis. *Am J Hum Genet* 2007; **81**: 559–575.
- Stegle O, Parts L, Durbin R, Winn J: A Bayesian framework to account for complex non-genetic factors in gene expression levels greatly increases power in eQTL studies. *PLoS Comput Biol* 2010; **6**: e1000770.
- Wu C, Orozco C, Boyer J *et al*: BioGPS: an extensible and customizable portal for querying and organizing gene annotation resources. *Genome Biol* 2009; **10**: R130.
- Zhou M, Sandercock AM, Fraser CS *et al*: Mass spectrometry reveals modularity and a complete subunit interaction map of the eukaryotic translation factor eIF3. *Proc Natl Acad Sci USA* 2008; **105**: 18139–18144.
- Hinnebusch AG: eIF3: a versatile scaffold for translation initiation complexes. *Trends Biochem Sci* 2006; **31**: 553–562.
- Park HS, Himmelbach A, Browning KS, Hohn T, Ryabova LA: A plant viral 'reinitiation' factor interacts with the host translational machinery. *Cell* 2001; **106**: 723–733.
- Spahn CM, Kieft JS, Grassucci RA *et al*: Hepatitis C virus IRES RNA-induced changes in the conformation of the 40S ribosomal subunit. *Science* 2001; **291**: 1959–1962.
- Aragón T, de la Luna S, Novoa I, Carrasco L, Ortín J, Nieto A: Eukaryotic translation initiation factor 4G1 is a cellular target for NS1 protein, a translational activator of influenza virus. *Mol Cell Biol* 2000; **20**: 6259–6268.
- Herberg JA, Kafrou M, Gormley S *et al*: Transcriptomic profiling in childhood H1N1/09 influenza reveals reduced expression of protein synthesis genes. *J Infect Dis* 2013; **208**: 1664–1668.
- Kim JT, Kim KD, Song EY *et al*: Apoptosis-inducing factor (AIF) inhibits protein synthesis by interacting with the eukaryotic translation initiation factor 3 subunit p44 (eIF3g). *FEBS Lett* 2006; **580**: 6375–6383.
- Kim JT, Lee SJ, Kim BY *et al*: Caspase-mediated cleavage and DNase activity of the translation initiation factor 3, subunit G (eIF3g). *FEBS Lett* 2013; **587**: 3668–3674.

Supplementary Information accompanies this paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)