ARTICLE

Dominant variants in the splicing factor *PUF60* cause a recognizable syndrome with intellectual disability, heart defects and short stature

Salima El Chehadeh^{*,1,2}, Wilhelmina S Kerstjens-Frederikse³, Julien Thevenon^{1,4}, Paul Kuentz^{1,4,5}, Ange-Line Bruel⁴, Christel Thauvin-Robinet^{1,4}, Candace Bensignor⁶, Hélène Dollfus^{2,7}, Vincent Laugel^{7,8}, Jean-Baptiste Rivière^{1,4,5}, Yannis Duffourd¹, Caroline Bonnet⁹, Matthieu P Robert^{10,11}, Rodica Isaiko¹², Morgane Straub¹², Catherine Creuzot-Garcher¹², Patrick Calvas¹³, Nicolas Chassaing¹³, Bart Loeys¹⁴, Edwin Reyniers¹⁴, Geert Vandeweyer¹⁴, Frank Kooy¹⁴, Miroslava Hančárová¹⁵, Marketa Havlovicová¹⁵, Darina Prchalová¹⁵, Zdenek Sedláček¹⁵, Christian Gilissen¹⁶, Rolph Pfundt¹⁶, Jolien S Klein Wassink-Ruiter³ and Laurence Faivre^{1,4}

Verheij syndrome, also called 8q24.3 microdeletion syndrome, is a rare condition characterized by ante- and postnatal growth retardation, microcephaly, vertebral anomalies, joint laxity/dislocation, developmental delay (DD), cardiac and renal defects and dysmorphic features. Recently, *PUF60* (*Poly-U Binding Splicing Factor 60 kDa*), which encodes a component of the spliceosome, has been discussed as the best candidate gene for the Verheij syndrome phenotype, regarding the cardiac and short stature phenotype. To date, only one patient has been reported with a *de novo* variant in *PUF60* that probably affects function (c.505C>T leading to p.(His169Tyr)) associated with DD, microcephaly, craniofacial and cardiac defects. Additional patients were required to confirm the pathogenesis of this association and further delineate the clinical spectrum. Here we report five patients with *de novo* heterozygous variants in *PUF60* identified using whole exome sequencing. Variants included a splice-site variant (c.24+1G>C), a frameshift variant (p.(Ile136Thrfs*31)), two nonsense variants (p.(Arg448*) and p.(Lys301*)) and a missense change (p.(Val483Ala)). All six patients with a *PUF60* variant (the five patients of the present study and the unique reported patient) have the same core facial gestalt as 8q24.3 microdeletions patients, associated with DD. Other findings include feeding difficulties (3/6), cardiac defects (5/6), short stature (5/6), joint laxity and/or dislocation (5/6), vertebral anomalies (3/6), bilateral micropthalmia and irido-retinal coloboma (1/6), bilateral optic nerve hypoplasia (2/6), renal anomalies (2/6) and branchial arch defects (2/6). These results confirm that *PUF60* is a major driver for the developmental, craniofacial, skeletal and cardiac phenotypes associated with the 8q24.3 microdeletion.

European Journal of Human Genetics (2017) 25, 43–51; doi:10.1038/ejhg.2016.133; published online 2 November 2016

INTRODUCTION

The increasing use of next-generation sequencing technologies, including whole exome sequencing and whole genome sequencing, has led to the massive identification of novel genes with *de novo* heterozygous variants as a major cause of intellectual disability (ID).^{1–3} The *PUF60* (*Poly-U Binding Splicing Factor 60 kDa*) gene encodes a general splicing factor that facilitates 3' splice-site recognition at the early stages of spliceosome assembly.^{4,5} *PUF60* has been partly associated with the rare 8q24.3 microdeletion syndrome phenotype,

in particular craniofacial and heart manifestations, in six patients and one fetal case. This rare condition, also named Verheij syndrome, is characterized by low birth weight and growth retardation, microcephaly, vertebral anomalies, joint laxity, and/or dislocation, developmental delay (DD), cardiac defects, renal defects and dysmorphic features.^{6,7} It can be considered, at least for now, as a contiguous gene syndrome characterized by the existence of apparently unrelated clinical features in the same patient, resulting from a chromosomal rearrangement involving contiguous dosage-sensitive genes. Indeed, in

¹FHU TRANSLAD, Centre de Référence Maladies Rares «Anomalies du développement et syndromes malformatifs» de l'Est, Centre de Génétique, CHU de Dijon, Dijon, France; ²Service de génétique médicale, Institut de génétique médicale d'Alsace (IGMA), Centre de Référence Maladies Rares «Anomalies du développement et syndromes malformatifs» de l'Est, Hôpitaux Universitaires de Strasbourg, Hôpital de Hautepierre, Strasbourg, France; ³Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; ⁴EA 4271, Génétique et Anomalies du Développement (GAD), Université de Bourgogne, Dijon, France; ⁵Laboratoire de Biologie Moléculaire, Plateau Technique de Biologie, CHU de Dijon, Dijon, France; ⁶Service de Pédiatrie, Hôpital d'enfants, CHU de Dijon, Dijon, France; ⁷U1112 Laboratoire de Génétique Médicale, Faculté de médecine, Université de Strasbourg, Strasbourg, France; ⁸Service de Neuropédiatrie, Hôpitaux Universitaires de Strasbourg, Hôpital de Hautepierre, Strasbourg, France; ⁹Service de Cardiopédiatrie, Hôpital d'enfants, CHU de Dijon, Dijon, France; ¹⁰Service d'Ophtalmologie, APHP, Hôpital Universitaire Necker-Enfants Malades, Paris, France; ¹¹COGNAC-G, UMR 8257: CNRS-IRBA-Université Paris Descartes, Sorbonne Paris Cité, Paris, France; ¹²Service d'Ophtalmologie, Hôpital Civil, CHU de Dijon, Dijon, France; ¹³Service de Génétique Médicale, & UMR1056 INSERM-UPS, Hôpital de Purpan, CHU de Toulouse, Toulouse, France; ¹⁴Centre of Medical Genetics, University Hospital of Antwerp, Antwerpen-Edegem, Belgium; ¹⁵Department of Biology and Medical Genetics, Charles University 2nd Faculty of Medicine and University Hospital Motol, Prague, Czech Republic; ¹⁶Department of Human Genetics, Radboud Institute for Molecular Life Sciences and Donders Centre for Neuroscience, Radboud University Medical Center, Nijmegen, The Netherlands

*Correspondence: Dr S El Chehadeh, Service de génétique médicale, Hôpital de Hautepierre, Avenue Molière, 67098 Strasbourg Cedex, France. Tel: +33 3 88 12 81 20; Fax: +33 3 88 12 81 25; E-mail: salima.elchehadeh@chru-strasbourg.fr

Received 13 March 2016; revised 28 July 2016; accepted 23 August 2016; published online 2 November 2016

2013, Dauber *et al*⁸ reported on five individuals carrying overlapping 8q24.3 microdeletions that included *SCRIB*, *NRBP2* and *PUF60*, and demonstrated by using functional experiments on zebrafish that the patients' phenotype resulted from a combination of endophenotypes induced by the haploinsufficiency of *PUF60* (short stature, craniofacial and cardiac anomalies) and *SCRIB* (coloboma and renal anomalies). They also identified a patient with microcephaly, short stature, DD and heart defects with a *de novo* c.505C>T variant leading to a p.His169Tyr change in *PUF60*.⁸ This is the only patient reported to date with a *de novo* function-affecting variant in *PUF60*.

Here, we report five additional patients with similar clinical features and *de novo* heterozygous variants in *PUF60* identified by whole exome sequencing (WES) and confirm the major role of *PUF60* in the phenotype of patients with the 8q24.3 microdeletion syndrome.

SUBJECTS AND METHODS

Patients

WES was performed in five unrelated individuals with developmental delay, congenital malformations and distinct facial dysmorphism in order to identify the molecular cause of their conditions, after negative standard genetic investigations and malformation screening. WES was used in a diagnostic approach in patient 1 and international data sharing was initiated to find other patients with very rare *de novo* *PUF60* variants. This allowed us to identify patients 2–5. Written informed consent was obtained from all of the subjects' parents for WES and for the use of photographs for patients 1, 3, 4 and 5.

Whole exome sequencing

WES was performed for patient 1 at Integragen (Evry, France). Targeted regions were captured using the SureSelect Human All Exon V5 kit (Agilent, Santa Clara, CA, USA) according to the manufacturer's recommendations. Sequencing of paired-end 76-bp reads was carried out on a HiSeq 2000 (Illumina, San Diego, CA, USA). Reads were aligned to the human genome reference sequence (GRCh37/hg19 build of UCSC Genome Browser; <http://genome.ucsc.edu>) with the Burrows–Wheeler Aligner (BWA, v.0.6.2). Duplicate paired-end reads were marked with Picard v.1.77 (<http://picard.sourceforge.net/>). The Genome Analysis Toolkit (GATK) v.2.6–4 was used for base quality score recalibration, indel realignment and variant discovery. Variants were annotated with SeattleSeq SNP Annotation 138 (<http://snp.gs.washington.edu/SeattleSeqAnnotation138/>). Candidate *de novo* events were systematically identified by focusing on protein-altering and splice-site variants present at a frequency of $\leq 1\%$ in the database of single-nucleotide polymorphisms (dbSNP) (dbSNP 146; <http://www.ncbi.nlm.nih.gov/SNP/>), absent from 100 local exomes of unaffected individuals and supported by at least four reads in the index subject and absent in both parents (less than two variant reads) at base-pair positions covered by at least four reads in the entire trio. Candidate *de novo* events were then manually inspected using the Integrative Genomics Viewer (IGV, Broad Institute; <http://www.broadinstitute.org/igv/>).

WES was performed in patients 2 and 3 in a *trio* diagnostic approach (patient and both parents) at the Radboud University Medical Center in Nijmegen, The Netherlands, essentially as described before.⁹ Exome capture was performed with the Agilent SureSelect Human All Exon v4 enrichment kit (Agilent Technologies). WES was performed on the Illumina HiSeq platform (BGI, Copenhagen, Denmark). Data were analyzed with BWA (read alignment) and GATK (variant calling) software packages. Variants were annotated using an in-house-developed pipeline. Variants were prioritized using an in-house-designed 'variant interface' and manual curation. *De novo* variants were considered the main candidates. Potentially causative variants were confirmed by Sanger sequencing.

WES was performed in patient 4 in a *trio* research approach at the Charles University and University Hospital Motol in Prague, Czech Republic. Whole exome capture was performed with SeqCap EZ Human Exome Library v3.0 (Roche NimbleGen, Madison, WI, USA). WES was performed on the Illumina HiSeq 1500 platform (Illumina) according to the manufacturer's recommendations for paired-end 101 bp reads. Sequences aligned using Novoalign (Novocraft, Petaling Jaya, Selangor, Malaysia) were processed according to GATK best

practices workflow. Variants were annotated using the GEMINI framework and subjected to several levels of filtering. Common variants (allele frequency >0.01) from public human genetic variation databases (dbSNP, 1000 Genomes, Exome Variant Server (EVS) and the Exome Aggregation Consortium (ExAC)) or from the in-house database were removed.

WES was performed in patient 5 in a *trio* diagnostic approach (patient and both parents) at the Center of Medical Genetics of the University and University Hospital of Antwerp, Belgium, essentially as described before.¹⁰ In brief, exome capture was performed with the SeqCap EZ Human Exome Library enrichment kit (Nimblegen). WES was performed on the Illumina HiSeq platform. Data were analyzed using BWA (read alignment) and GATK (variant calling) software packages. Variants were annotated using an in-house-developed pipeline. Variants were prioritized using our program VariantDB.¹¹ Potentially causative variants were confirmed by Sanger sequencing.

We have submitted data for the variants to the gene variant database at www.LOVD.nl/PUF60 (patient IDs 73735–73739, 78890).

RESULTS

Clinical assessment

Patient 1. This 3-year-old French girl (Figure 1) was the second child of healthy non-consanguineous French parents. There was no family history. The pregnancy was marked by isolated IUGR. She was born at 39 weeks of gestation (WG), she weighed 2675 g (7th centile), had a length of 43 cm (≤ 3 rd centile) and a head circumference of 34 cm (50th centile). At birth, bilateral microphthalmia associated with iris coloboma, a *naevus flammeus* on the forehead and eyelids, bilateral branchial cleft cysts, bilateral clinodactyly of the fifth digits and valgus deformity of one foot were noted. Several facial features, including asymmetric and posteriorly rotated ears, retrognathism, long philtrum with thin upper lip and anteverted nares, were also noted (Figure 1). She also had a ventricular septal defect and unilateral hip dislocation. Fundus and brain MRI showed bilateral microphthalmia and retinal coloboma involving the optic discs, bilateral optic nerve hypoplasia (Figure 2) and an ectopic posterior pituitary without interruption of the pituitary stalk. Endocrinological tests, abdominal/renal ultrasonography and skeleton X-rays were normal. Standard karyotyping (46, XX) and array-CGH were normal. Molecular analysis of several genes, including *RAX*, *VSX2*, *OTX2*, *SOX2* and *FOXE3*, known to be responsible for microphthalmia, was performed (Sanger sequencing). The results were negative, as was the molecular analysis of a targeted panel of 187 known or candidate genes involved in ocular malformations (Toulouse, France). This panel comprised genes that were already associated with ocular developmental defects (*ALDH1A3*, *ATOH7*, *BCOR*, *BMP4*, *BMP7*, *c12orf57*, *CHD7*, *COX7B*, *FNBP4*, *FOXC1*, *FOXE3*, *GDF6*, *GJA8*, *HCCS*, *MFRP*, *OTX2*, *PAX2*, *PAX6*, *PITX2*, *PITX3*, *RARB*, *RAX*, *SIX6*, *SMOC1*, *SOX2*, *STRA6*, *TENM3*, *VAX1* and *VSX2*) and other candidate genes, including candidates from the retinoic acid or SHH pathways or identified through our previous exome sequencing experiments. Libraries were pooled and sequenced on an Illumina HiSeq 2000 as paired-end 75 bp reads. The bioinformatics analysis of sequencing data was based on the Illumina pipeline (CASAVA1.8.2). Mean coverage of the targeted exons was $250\times$ and $\geq 99\%$ of the targeted exons were covered over $25\times$. The proband had developmental delay but was able to sit without support at 11 months. On the last examination at age 3 years, she had just acquired independent walking. Her hearing appeared to be normal. She had poor eye contact because of visual impairment (refraction +6), reacted to light and could follow large moving targets. She had behavioral anomalies with intolerance to frustration and echolalia. She had severe feeding difficulties and could only eat blended food, with frequent vomiting. Her measurements were 83 cm for height (< -3 SD), 10 kg for weight (< -3 SD) and 47.5 cm for OFC (-1.5 SD)

Figure 1 Facial characteristics of the individuals with *PUF60* variants. (a–c) Frontal and lateral views of patient 1 at birth (a) at 3 years and 3 months (b) and at 1 year (c) demonstrating facial dysmorphism including a square face, full cheeks, prominent forehead with bitemporal narrowing, low-set eyebrows, asymmetric and posterior rotated ears, bilateral colobomatous microphthalmia with horizontal nystagmus and convergent strabismus, bilateral epicanthal folds, wide nasal bridge with broad nasal tip and anteverted nares, thin upper lip with a long philtrum, micrognathism and broad short neck. (d) Lateral view of the neck of patient 1 showing a branchial cleft cyst (arrow). (e) Full-body view of patient 1 showing the contrast between the normal head circumference and the low body height and weight with relatively short arms, and a pectus excavatum. (f, g) Frontal and lateral views of patient 3 at birth (f) and at 3 years and 10 months (g) showing an elongated face, bilateral epicanthal folds, short up-slanted palpebral fissures, long eyelashes, anteverted nares, long prominent philtrum, thin upper lip and a short neck. (h–k) Frontal and lateral views of patient 4 at the age of 8 months (h), 2 years (i) and 17 years (k). (j) Lateral view of the neck of patient 4 showing a branchial cleft cyst (arrow). (l, m) Frontal and lateral view of patient 5 at the age of 11 years (l) showing deep set eyes, broad nasal bridge with protruding ears. (m) Flat feet with short halluces and sandal-gap deformity.

Figure 2 Fundus and brain MRI results. (a) Fundus view showing the retinal coloboma (full-line arrow) of left eye involving the optic disk (dotted-line arrow) – type IV of the Gopal classification. (b–d) Brain and optic nerve MRI (2 months) showing bilateral microphthalmia (b) associated with irido coloboma (black arrow) and retinal coloboma (white arrow), and the bilaterally thin (hypoplastic) optic nerve hypoplasia predominantly in the intra-orbital portion (c, d) (white arrows).

with micromelia. Her facial features are summarized in Table 1. Bilateral clinodactyly of the fifth fingers, pectus excavatum and two branchial cleft cysts were also noted (Figure 1).

Patient 2. The second patient, a 14-year-old boy, was the fourth child of healthy, non-consanguineous Dutch parents. He was born after an uneventful pregnancy. His growth was normal. Developmental delay was noted from the age of 1 year. At the age of 6 years, his IQ was 46 and an autism spectrum disorder was diagnosed. The family history was negative for ID or congenital malformations. Distinct dysmorphic features were noted: exotropia, a square face with a prominent forehead, low-set eyebrows, a wide nasal bridge with broad nasal tip and anteverted nares, a long philtrum and bilateral clinodactyly of the fifth digits (Figure 1 and Table 1). The parents did not permit a brain MRI. *FMRI* testing and metabolic screening were negative. Array-CGH revealed two numerical variants inherited from the parents: hg19 chr11:g.(133072607_133078519)_(133691682_133693512)dup; chr13:g.(48465866_48466014)_(48540617_48540958)del (ISCN 2016 nomenclature: arr[GRCh37] 11q25(133078519_133691683)x3 pat, 13q14.2(48466014_48540618)x1 mat). The maternal deletion at 13q14.2 (75 kb) only includes the *SUCLA2* gene that is associated with autosomal recessive mitochondrial depletion syndrome type 5. *SUCLA2* gene was therefore sequenced in the child, but no variants were detected on the other allele. The paternal duplication at 11q25 (600 Kb) includes *OPCML* gene that has uniquely been involved in ovarian and colorectal cancer to date. Hence, the absence of potential causative gene, notably syndromic ID gene, in both rearrangements, combined to the fact that they were inherited from unaffected parents, led us to consider those as not likely to contribute to the patient's phenotype. In addition, no affected patients have been reported in the literature or in the DECIPHER (<https://decipher.sanger.ac.uk/>) database with such chromosomal aberrations.

Patient 3. This 7-year-old Dutch boy was the first child of healthy, non-consanguineous Dutch parents. The family history was unremarkable as was the pregnancy. He was born at 38 WG, weighed 2570 g (−2.1 SD) and his height was 49.5 cm (−0.7 SD). He was microcephalic (head circumference 32 cm (−3.2 SD) at age 3 weeks). Soon after birth, he was diagnosed with a heart defect, including a truncus arteriosus and a ventricular septal defect, for which he had multiple surgical interventions. He developed an inspiratory stridor

and was diagnosed with a cricoid stenosis at the age of 2 years, probably because of multiple intubations. He also had difficulties swallowing and was fed by a percutaneous gastric tube from age 2 years. His facial features included an elongated face, short up-slanted palpebral fissures, bilateral epicanthal folds, long eyelashes, anteverted nares, long prominent philtrum, thin upper lip, short neck and micromelia (Figure 1 and Table 1). Ocular coloboma were not detected. A cervical spine CT-scan showed fusion of the left facet-joints of C5–C6. The brain MRI was normal. There were no renal malformations.

On follow-up, he showed DD and growth retardation (height −2 SD, weight −2.5 SD and head circumference < −2.5 SD). At the age of 4 years and 5 months, his verbal IQ was normal (100) but his performance IQ was much lower (67). His motor skills showed a delay of 1 year. At the last evaluation (6 years and 10 months), he attended a special school, because of his developmental delay and behavioral problems.

A 105K oligo-array (Agilent) was normal and molecular analysis of the *NIPBL* and *SMC1A* genes (Cornelia de Lange syndrome) was negative.

Patient 4. Patient 4 is a 17-year-old Czech boy, the first child of healthy non-consanguineous parents. The delivery was induced at 36 WG. His birth weight was 2470 g, length was 48 cm and head circumference was 33 cm. He needed intensive care because of respiratory distress syndrome during the first 3 weeks of life. Abdominal and hip ultrasound showed hypoplastic kidneys with left pyelectasis and hip dysplasia, respectively. His psychomotor development was delayed from the beginning (sitting without support at the age of 9 months, crawling at the age of 11 months and first steps from the age of 18 months). Eye examinations performed when he was a toddler were normal and no further examinations were done. Speech development was also delayed, with babbling until 4 years of age. At the last examination, speech was slow with signs of dyslalia. At the age of 4 years, moderate aortic insufficiency was revealed, associated with asymmetric coarctation and a dysplastic aortic valve. At 17 years of age, the height was 164.5 cm (−2.1 SD), weight was 59.6 kg (−0.9 SD) and the head circumference was 54 cm (< −1.6 SD). He had bifrontal narrowing, fused, thick and arched eyebrows, large dysplastic ears, a wide nasal bridge and long nose with a broad nasal tip and anteverted

Table 1 Phenotype of the patients with a *PUF60* variant and the patients with a 8q24.3 deletion comprising *PUF60*

Patients with a PUF60 variant													Patients with a 8q24.3 deletion				
	P1	P2	P3	P4	P5	Dauber et al, ⁸ P6	Verheij et al, ⁶ P1	Dauber et al, ⁸ P1	Dauber et al, ⁸ P2	Dauber et al, ⁸ P3	Dauber et al, ⁸ P4	Dauber et al, ⁸ P5	Wells et al, ⁷ P1				
Gender	F	M	M	M	F	F	M	F	M	F	F	F	F				
Age	3 y	14 y	7 y	17 y	12 y 2 m	21 y	4 m	10 y	6 y	17 y	9 y	11 y	28+6 WG				
Facial features																	
Square face	+	+	–	+	+	+	+	+	+	+	+	+	+				
Full cheeks	+	–	+	+	+	–	+	+	+	+	+	+	–				
Bitemporal narrowing	+	–	+	+	–	ND	+	ND	ND	+	+	+	+				
Prominent forehead	+	+	+	+	+	+	+	+	+	–	+	+	–				
Low set eyebrows	+	+	+	+	+	+	+	–	+	–	+	+	+				
Wide nasal bridge	+	+	–	+	+	–	+	ND	+	–	+	+	+				
Broad nasal tip	+	+	+	+	+	+	+	+	+	+	+	+	+				
Anteverted nares	+	+	+	+	–	+	+	+	+	+	+	+	+				
Long philtrum	+	+	+	+	+	+	+	+	+	+	+	+	+				
Thin upper lip	+	–	+	+	+	+	+	+	+	+	+	+	+				
Micro- or retrognathism	+	–	–	+	–	+	+	+	–	–	–	+	+				
Short neck	+	–	–	+	+	+	+	–	+	+	+	+	+				
Branchial arch anomalies	Branchial cleft cysts	–	–	Branchial cleft cysts, uvula dimple	–	–	Branchial arch remnant, fistula	Bilateral preauricular pits	–	–	Bilateral preauricular pits	–	–				
Eye anomalies																	
Coloboma	+ (IRC)	–	–	–	–	–	+ (IRC)	+ (IC)	+ (IRC)	–	–	–	–				
Other	Bilateral micro- phthalmia and optic nerve hypoplasia	Strabismus	–	– ^a	Bilateral optic nerve hypoplasia	–	–	–	–	–	–	Hyperopia	–				
Cardiac defects																	
VSD	+	–	+	–	–	+	+	–	–	+	–	+	–				
Other	–	–	Truncus arteriosus	Aortic insufficiency	AVSD	Bicuspid aortic valve	–	–	–	Truncus arteriosus	–	Aortic coarctation	AVSD, hypoplastic aortic arch				
Laryngo-tracheo-bronchomalacia																	
Growth	–	–	+	–	–	–	+	–	–	–	–	+	–				
IUGR	+	–	+	–	–	+	+	+	–	+	+	–	+				
Short stature	+	–	+	+	+	+	+	+	+	+	+	+	NA				
Micromelia	+	–	+	–	–	+ (asymmetric)	ND	ND	ND	ND	ND	ND	+				
Microcephaly	–	–	+	–	–	+	+	+	+	–	–	+	–				
Skeletal anomalies																	
Joint laxity/dislocation	+ ^b	–	+	+	+	+ ^b	+ ^b	–	+	+ ^c	+	+ ^b	–				
Distal limb anomalies	+ a, b	+ a	–	+ b, f, i	j, k	+ c	+ d, e	–	–	–	+	+ a, h	+ a, f, g				
Vertebral anomalies	–	–	Fusion of the left facet joints of C5–C6	Unfused arch of C7, scoliosis	–	Hemivertebrae, scoliosis	–	+	–	Fusion of L5-S1	–	Hemivertebrae, scoliosis	Mild scoliosis, abnormal third sacral vertebra				

Table 1 (Continued)

Patients with a PUF60 variant														Patients with a 8q24.3 deletion													
	P1	P2	P3	P4	P5	Dauber <i>et al</i> , ⁸ P6	Verheij <i>et al</i> , ⁶ P1	Dauber <i>et al</i> , ⁸ P1	Dauber <i>et al</i> , ⁸ P2	Dauber <i>et al</i> , ⁸ P3	Dauber <i>et al</i> , ⁸ P4	Dauber <i>et al</i> , ⁸ P5	Wells <i>et al</i> , ⁷ P1														
Feeding difficulties	+	-	+	-	+	-	+	-	-	+	+	+	NA														
Neurology																											
Developmental delay	+	+	+	+	+	+	+	-	+	+	+	+	NA														
Seizures	-	-	-	- ^d	-	+	+	-	-	-	-	+	-														
Brain malformation	Posterior pituitary ectopia	ND	-	ND	CC hypoplasia	Enlarged 3rd and lateral ventricles, thin CC	Hypoplastic CC, cerebral atrophy, enlarged ventricles	-	Cerebral atrophy	-	Ischemic encephalomalacia	Volume loss in the left temporal lobe	-														
Renal anomalies	-	-	-	Hypoplastic kidneys	Pelvic left kidney with unilateral VUR	-	-	Left renal agenesis	-	Left polycystic kidney	Ectopic fused kidneys	Right kidney hypoplasia with renal cyst	-														
Other	Naevus flammeus on the forehead and eyelids, pectus excavatum	Autistic features	c.1342C>T p.(Arg448*)	Autistic features, pectus excavatum, scoliosis	Trilobar left lung	Cleft palate, BPD, pes planus, hemifacial microsomia		Coarse hair, CHL	Recurrent infections, hiatus hernia	-	Forehead capillary hemangioma	Cyclic vomiting	Small medial alveolar cleft														
<i>De novo</i> PUF60 variants (NG_033879.1; NM_078480.2)	c.24+1G>C chr8:g.144911449C>G	c.1342C>T p.(Arg448*)	c.1448T>C p.(Val483Ala)	c.407_410delTCTA p.(Ile136Thrfs*31)	c.901A>T p.(Lys301*)	c.505C>T p.(His169Tyr)		Deletion	Deletion	Deletion	Deletion	Deletion	Deletion														

Abbreviations: AVSD, atrioventricular septal defect; BPD, bronchopulmonary dysplasia; CC, corpus callosum; CHL, conductive hearing loss due to middle ear malformation; IC, iris coloboma; IRC, irido-retinal coloboma; IUGR, intrauterine growth retardation; NA, not applicable; ND, not determined; VSD, ventricular septal defect; VUR, vesicoureteral reflux; WG, weeks of gestation.
 Distal limb anomalies: a: clinodactyly of the fifth digit; b: valgus deformity of the feet; c: absent right thumb, hypoplastic metacarpal and scaphoid bones; d: bilateral post-axial polydactyly; e: oligodactyly and fusion of the third and fourth metatarsal of the left foot; f: brachymesophalangy of the fifth digit; g: rocker-bottom feet; h: Bilateral transverse palmar creases; i: overlying proximal thumb; j: brachydactyly, k: flat feet with short halluces and sandal gap deformity.
^aNo focused ophthalmological examinations have been performed since the age of 3 years.
^bHip dislocation.
^cElbow subluxation with malformation of the great sigmoid cavity.
^dEEG showed epileptiform discharges with absence of seizures.

nares, a long smooth philtrum, thin upper lip and short neck. A branchial cleft cyst was noted on the neck as well as a fistula above the uvula. Skeletal features included slight shortening of the trunk, significant flattening of the chest, pectus excavatum, scoliosis, valgus deformity of the feet, long fingers and toes, overlying proximal thumbs, short fifth fingers with clinodactyly on the right digit, bilateral cervical ribs and an unfused arch of vertebra C7. He has moderate ID and autism, hyperactivity and transitional tic disorder. The EEG showed epileptiform discharges with the absence of seizures. SNP array analysis (Illumina HumanCytoSNP-12 BeadChips) was performed and was normal. Brain MRI has not yet been performed.

Patient 5. This female proband was born at 41 WG as the first child of non-consanguineous Belgian parents. Birth weight was 3560 g for a length of 50 cm. At birth, she was diagnosed with atrioventricular septal defect that was surgically corrected at age 1 month. Lung imaging revealed a trilobar left lung. She was also diagnosed with a left pelvic kidney associated with unilateral vesicoureteral reflux. Over time, left kidney function decreased significantly. During the first months of life, she had severe feeding problems and was fed by gastric tube. She developed postnatal growth retardation. At the age of 11 years and 2 months, she had a skeletal age of 9 years and 2 months. At the age of 3 years, she underwent tonsillectomy because of extreme snoring and obstructive apnea. She attended a regular school until the age of 7 years but is now in special education. MRI of the brain showed hypoplasia of the corpus callosum. The ophthalmological evaluation revealed bilateral optic hypoplasia.

At the clinical examination at age 12 years and 2 months, occipitofrontal circumference was 53.2 cm (0 SD), length 131 cm (−2.5 SD) and weight 33.3 kg (−1 SD). Craniofacial characteristics included deep set eyes, a broad nasal bridge and protruding ears. Both hands showed brachydactyly. She had flat feet with short halluces and sandal-gap deformity. Skin was normal.

Exome sequencing

Exome sequencing from a blood-derived DNA sample from patient 1 generated over 2.96 Gigabases of mappable sequence with a median sequencing depth provided for RefSeq coding exons and splice junctions of 85.8×. Depth of coverage of RefSeq coding bases and splice junctions was 94.2× for at least 10 reads and 91× for at least 20 reads calculated with the Genome Analysis Toolkit Depth of Coverage tool. After filtering, among the 16 622 non-synonymous splice-site and indel variants, 368 rare variants remained. Seventeen candidate variations of OMIM genes were found. Among these, after studying their expression and the disorders they were involved in, and

comparisons with our patient's phenotype, only three OMIM gene variants were retained. These candidate variants were validated and tested in parental DNA samples. A *PAX6* missense change (NM_000280.4:c.518C>T, p.(Thr173Met)) and a *PACSI* missense change (NM_018026.3:c.2197C>T, p.(Arg733Trp)) were found to be inherited from an asymptomatic parent, and a *de novo* *PUF60* NM_078480.2:c.24+1G>C (chr8:g.144911449C>G) variant was found. Taking into account the familial segregation and reverse phenotyping in favor of Verheij syndrome,¹² the *PUF60* variant was considered the one affecting gene function and thus the likely cause of the phenotype (Table 2). Indeed, exon 1 skipping is expected as the wild-type donor site is predicted to be broken (Human Splicing Finder score of −31.86: most probably affecting splicing; <http://www.umd.be/HSF3/HSF.html>).

Trio WES and data analysis of patients 2 and 3 and their parents did not reveal any function-affecting variants of genes known to be associated with ID. Exome-wide analysis revealed a *de novo* *PUF60* nonsense variant (NM_078480.2:c.1342C>T; p.(Arg448*)) in patient 2, and a *de novo* missense variant in *PUF60* (NM_078480.2; c.1448T>C; p.(Val483Ala)) in patient 3 (Table 1 and Figure 3) that were considered of interest given the patients' phenotypes. The nonsense variant p. Arg448* in exon 11 leads to protein truncation at position 448 with loss of RRM3 domain (462–549) (Uniprot). These variant were confirmed by Sanger sequencing. Both *PUF60* variants identified were absent from all of the public genetic variant databases used (dbSNP142; Exome Variant Server, NHLBI GO Exome Sequencing Project (ESP); ExAC, accessed in February 2016). Genomic positions were conserved (GERP 5.270) and bioinformatics tools supported the deleterious effect of the *de novo* missense change (PolyPhen-2 0.989, PhastCons 0.998, Grantham 64, CADD 24.1, Mutation Taster 0.99).

Trio WES and data analysis of patient 4 revealed a private heterozygous *de novo* deletion of four nucleotides in the *PUF60* gene (NM_078480.2:c.407_410delTCTA, p.I136Tfs*31). Thanks to reverse phenotyping, the patient's phenotype was deemed to correspond to the Verheij syndrome. This variant was also absent from all of the public genetic variant databases used (dbSNP142; Exome Variant Server, NHLBI GO ESP; ExAC, accessed in June 2016). The genomic position was conserved (GERP 4.28).

Trio WES and data analysis of patient 5 and her parents revealed a single *de novo* variant in known ID genes from a predefined panel, namely a nonsense mutation in the *PUF60* gene: NM_07848.2: c.901C>T leading to p.Lys301*. Taking into account the familial segregation, the nature of the identified variant and reverse phenotyping in favor of Verheij syndrome, the *PUF60* variant was considered to

Table 2 Pathogenicity of each variant evaluated by different databases

	Variant (hg19)	Amino-acid change	dbSNP	Frequency						
				in ExAC	Grantham	Polyphen	GERP	CADD	MutationTaster	SIFT
Patient 1	chr8:g.144911449C>G NG_033879.1; NM_078480.2:c.24+1G>C	NA	NA	0	NA	NA	4.740	21.400	1	NA
Patient 2	chr8:g.144899118G>A NG_033879.1; NM_078480.2:c.1342C>T	p.(Arg448*)	NA	0	NA	NA	5.280	36.000	1	NA
Patient 3	chr8:g.144898922A>G NG_033879.1; NM_078480.2:c.1448T>C	p.(Val483Ala)	NA	0	64	0.989	5.270	24.100	0.999	0.001
Patient 4	chr8:g.144900643_144900646delTAGA NG_033879.1; NM_078480.2:c.407_410delTCTA	p.(Ile136Tfrs*31)	NA	0	NA	NA	5.180	NA	1	NA
Patient 5	chr8:g.144899869T>A NG_033879.1; NM_07848.2:c.901A>T	p.(Lys301*)	NA	0	NA	NA	3.930	33.000	1	NA

Interestingly, one of the patients of the present study with a *PUF60* *de novo* variant had bilateral microphthalmia and irido-retinal coloboma (Table 1 and Figure 2). The presence of coloboma, as well as renal defects, observed in patients with the 8q24.3 deletion has been attributed to the *SCRIB* deletion, as 53% and 32% of the zebrafish embryos injected with a splice-blocking morpholino against *scrib* showed, respectively, coloboma and gross edema with abnormal diffusion of dextran, reflecting the loss of ability to maintain fluid homeostasis.⁸ In addition, *SCRIB* plays a key role in planar cell polarity and is involved in the formation of the pronephros and eye.¹³ *SCRIB* seems to remain a good candidate gene for kidney anomalies, as only two of the six patients with a *PUF60* variant had a renal phenotype. However, the presence of a coloboma in one patient from this report prompted us to reconsider the conclusion that the coloboma observed in (3/7) patients with Verheij syndrome was induced by *SCRIB* haploinsufficiency alone (Table 1).

We observed several distal limb anomalies in both groups of variant or deletion carriers, including clinodactyly of the fifth digit, valgus deformity of feet, absent right thumb, hypoplastic metacarpal and scaphoid bones, bilateral post-axial polydactyly, oligodactyly and fusion of the third and fourth metatarsal of the left foot, brachymesophalangy of the fifth digit and rocker bottom feet. None of these were recurrent except for clinodactyly of the fifth digit, which was present in 4/11 patients, although this feature is not very specific. It is well known that *PUF60* encodes a splicing factor that interacts directly with SF3B4, a component of the U2 pre-mRNA spliceosomal complex involved in Nager syndrome, suggesting that *PUF60* and *SF3B4* haploinsufficiency-associated phenotypes are partially due to a defect of spliceosome function.^{8,14} Interestingly, the patient carrying a *PUF60* variant reported by Dauber *et al*⁸ (patient 6) harbors clinical signs belonging to the Nager syndrome spectrum, including hemifacial microsomia with micro-retrognathism and preaxial upper-limb anomalies with an absent right thumb and hypoplastic metacarpal and scaphoid bones (Table 1).⁸

In conclusion, these results highlight the power of WES as an unbiased NGS-based strategy for identifying new genes causing very rare multisystemic conditions,¹⁵ and the importance of data sharing to identify their underlying clinical expression that is otherwise hampered by the paucity of reported cases. This strategy allowed us to report autosomal dominant *PUF60* variants in five unreported patients with overlapping features of 8q24.3 microdeletion syndrome and, thereby, to confirm the involvement of *PUF60* in the Verheij syndrome phenotype. Further screening of *PUF60* in other patients will provide important information about the frequency of *PUF60* variants and may help to further define the phenotype associated with these variants.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the patients and their families involved in the study, the University of Burgundy Centre de Calcul (CCuB) for technical support and management of the informatics platform. The Regional Council of Burgundy and the Centre

Hospitolo-Universitaire de Dijon supported this work (PARI 2014). Grant supports were NF-CZ11-PDP-3-003-2014, NT/14200 and 00064203. We thank Philip Bastable from the 'Pôle de Recherche' of Dijon University Hospital for a helpful review of this article.

- 1 Rauch A, Wieczorek D, Graf E *et al*: Range of genetic mutations associated with severe non-syndromic sporadic intellectual disability: an exome sequencing study. *Lancet* 2012; **380**: 1674–1682.
- 2 de Ligt J, Willemsen MH, van Bon BW *et al*: Diagnostic exome sequencing in persons with severe intellectual disability. *N Engl J Med* 2012; **367**: 1921–1929.
- 3 Gilissen C, Hehir-Kwa JY, Thung DT *et al*: Genome sequencing identifies major causes of severe intellectual disability. *Nature* 2014; **511**: 344–347.
- 4 Kielkopf CL, Lücke S, Green MR.: U2AF homology motifs: protein recognition in the RRM world. *Genes Dev* 2004; **18**: 1513–1526.
- 5 Hastings ML, Allemand E, Duelli DM, Myers MP, Krainer AR.: Control of pre-mRNA splicing by the general splicing factors PUF60 and U2AF(65). *PLoS One* 2007; **2**: e538.
- 6 Verheij JB, de Munick SA, Dijkhuizen T *et al*: An 8.35 Mb overlapping interstitial deletion of 8q24 in two patients with coloboma, congenital heart defect, limb abnormalities, psychomotor retardation and convulsions. *Eur J Med Genet* 2009; **52**: 353–357.
- 7 Wells C, Spaggiari E, Malan V *et al*: First fetal case of the 8q24.3 contiguous genes syndrome. *Am J Med Genet* 2016; **170**: 239–242.
- 8 Dauber A, Golzio C, Guenot C *et al*: SCRIB and PUF60 are primary drivers of the multisystemic phenotypes of the 8q24.3 copy-number variant. *Am J Hum Genet* 2013; **93**: 798–811.
- 9 Neveling K, Feenstra I, Gilissen C *et al*: A post-hoc comparison of the utility of sanger sequencing and exome sequencing for the diagnosis of heterogeneous diseases. *Hum Mutat* 2013; **34**: 1721–1726.
- 10 Helsmoortel C, Vandeweyer G, Ordoukhanian P, Van Nieuwerburgh F, Van der Aa N, Kooy RF.: Challenges and opportunities in the investigation of unexplained intellectual disability using family-based whole-exome sequencing. *Clin Genet* 2015; **88**: 140–148.
- 11 Vandeweyer G, Van Laer L, Loeys B, Van den Bulcke T, Kooy RF.: VariantDB: a flexible annotation and filtering portal for next generation sequencing data. *Genome Med* 2014; **6**: 74.
- 12 Hennekam RC, Biesecker LG.: Next-generation sequencing demands next-generation phenotyping. *Hum Mutat* 2012; **33**: 884–886.
- 13 Skouloudaki K, Puetz M, Simons M *et al*: Scribble participates in Hippo signaling and is required for normal zebrafish pronephros development. *Proc Natl Acad Sci USA* 2009; **106**: 8579–8584.
- 14 Bernier FP, Caluseriu O, Ng S *et al*: Haploinsufficiency of SF3B4, a component of the pre-mRNA spliceosomal complex, causes Nager syndrome. *Am J Hum Genet* 2012; **90**: 925–933.
- 15 Boycott KM, Dymant DA, Sawyer SL, Vanstone MR, Beaulieu CL: Identification of genes for childhood heritable diseases. *Annu Rev Med* 2014; **65**: 19–31.