Cholera toxin mediated regulation of the expression of Gq α and G11 α GTP binding proteins

Bukhtiar H. Shah

Department of Physiology and Pharmacology, The Aga Khan University, Karachi-74800, Pakistan; Present address; Endocrinology and Reproductive research branch, National Institutes of Health, Bethesda, MD 20892-4510, U.S.A.

Accepted 17 May 1999

Abbreviations: CT, cholera toxin; cAMP, cyclic adenosine monophosphate; db-cAMP, dibutyryl cAMP; FSK, forskolin; PKA, protein kinase A; IP₃, inositol 1,4,5-triphosphate; DAG, diacylglycerol

Abstract

Previously it has been shown that persistent activation of the stimulatory adenylyl cyclase pathway with cholera toxin (CT) downregulates the Gs α polypeptide (80%) in a cAMP-independent manner in C6 glioma cells (Shah, 1997). This study was conducted to examine the short and long term effects of CT on the regulation of pertussis toxin-sensitive and -insensitive G proteins and their transcripts in C6 glioma cells. Treatment of C6 cells with CT (100 ng/ml) up to 16 h had no effect on either Gi or Gq/11 α proteins. However, prolonged exposure (24-48 h) caused increased expression of Gi (20-30%) and Gq/11 α proteins (40%). Urea gradient gels, which can separate Gq α and G11 α proteins, revealed that prolonged CT treatment increased the expression of both of these G proteins. The CT-mediated enhanced expression of Gq α and G11 α proteins was accompanied by increased mRNA levels of these proteins as determined by RT/PCR. Cyclic-AMP elevating agents like forskolin (10 μ M) and db-cAMP (1 mM) mimicked the effect of CT on Gi but not Gq/11 α proteins. These studies show long term cAMP-dependent regulation of Gi and cAMP-independent expression of Gq/11 α proteins in C6 glioma cells.

Keywords: G-proteins, cholera toxin, C6 glioma cells, forskolin, dibutyryl cAMP, protein kinase A.

Introduction

The heterotrimeric guanine nucleotide binding regulatory proteins (G-proteins) mediate a variety of cellular processes by coupling transmembrane receptors to different effector molecules, including adenylyl cyclase

and inositol phospholipid-specific phospholipase C (PLC) (Exton, 1997; Eyster, 1998). Activation of Gs/adenylyl cyclase results in the production of cyclic AMP (cAMP) and activation of cAMP-dependent protein kinase (PKA). Activation of Gi protein leads to inhibition of adenylyl cyclase activity and thus decrease in cAMP levels (Sunahara *et al.*, 1996). The pertussis toxin-insensitive G-proteins include a family of Gq proteins (Gq, G11, G15 and G16) which mediate their effects by activating PLC and thus generating second messengers, inositol-1,4,5-triphosphate (IP₃) and diacylglycerol (DAG), thereby leading to activation of protein kinase C (PKC) and the mobilization of intracellular calcium, respectively (Berridge, 1993; Smrcka *et al.*, 1991). The two members of the Gq family, Gq and G11, are widely and quite often co-expressed in the body tissues (Milligan, 1993a).

Our previous studies have shown that rat C6 glioma cells express all these G-proteins (Gs, Gi, Gq and G11) (Shah and Milligan, 1995). C6 glioma cells are a bipotential cell line with oligodendrocytic and astrocytic properties. C6 cells can differentiate into type 2 astrocytes by addition of agents that increase intracellular cAMP (Messens and Slegers, 1992). These cells endogenously express receptors for a number of neurotransmitters including the adrenergic system and have proved to be a valuable model for studying the G-protein coupled signal transduction cascade (Hosoda *et al.*, 1994; Kato *et al.*, 1996). Cyclic AMP has been shown to exert multiple effects in C6 cells. These include differentiation of these cells towards more astrocyte phenotype (Segovia *et al.*, 1994), regulation of adrenergic receptors (Hosoda *et al.*, 1994), induction of interleukines (Slegers and Joniau, 1996) heat shock proteins (Kato *et al.*, 1996) and prostaglandin synthesis (Peterson *et al.*, 1996). It has been shown that PLC activation and phosphoinositide breakdown can be modulated by the adenylyl cyclase pathway and this effect is mediated by PKA (Haddock *et al.*, 1990; Liu and Simon, 1996). It has been shown that persistent CT-treatment of C6 glioma cells causes marked down-regulation of stimulatory G protein, Gs α (Shah, 1997), however, little is known about the regulation of Gi and Gq/11 α proteins. The present study was carried out to examine the effects of short and long term exposure of C6 cells to CT and agents that increase intracellular cAMP (forskolin and dibutyryl cAMP) on the expression of Gi and Gq/G11 proteins and their mRNA levels.

Materials and Methods

Materials

Taq polymerase and PCR buffer were from Promega. The cDNA synthesis kit was from Pharmacia LKB. Cholera toxin and all other chemicals were reagent grade and purchased from Sigma Chemical Co. All materials for tissue culture were from Gibco/BRL. Antibodies to various G-proteins were kindly provided by Dr. Graeme Milligan (University of Glasgow, UK).

Cell culture

Cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum, penicillin (100 units/ml) and streptomycin (100 µg/ml) in 5% CO₂ at 37°C. Cells were grown in 75 cm³ tissue-culture flasks and were harvested just before confluency. For these experiments, cells up to 25 passages were used. The treatment of cells with cholera toxin (100 ng/ml), dB-cAMP (1 mM) or forskolin (10 µM) for different times was carried out as described in the figure legends.

Preparation of membranes

Membranes were prepared from the cells by homogenization with a Teflon-on-glass homogenizer and differential centrifugation as previously described (Shah and Milligan, 1994). Frozen cell pellets were suspended in 5 ml of 10 mM Tris/HCl, 0.1 mM EDTA, pH 7.5 (buffer A) and the cells ruptured with 25 strokes of homogenizer. The resulting homogenate was centrifuged at 500 *g* for 10 min in a Beckman L5-50B centrifuge with a Ti 50 rotor, to remove unbroken cells and nuclei. The supernatant was further centrifuged at 48,000 *g* for 10 min. The pellet from the second centrifugation was washed with buffer A and recentrifuged at 48,000 *g* for 10 min. Finally the pellet was resuspended in buffer A to yield a protein concentration of 1-3 mg/ml and stored at -80°C until required. Protein concentration was measured according to the method of Lowry et al (1951).

Western blots

The generation and specificities of the various antisera used in this study are already described (Mitchell *et al.*, 1991). Membrane samples were resolved by SDS/PAGE in 10% (w/v) acrylamide gels overnight at 60 V. For resolving Gqα and G11α, urea linear gradient gels were prepared and run overnight at 100 V. Proteins were transferred to nitrocellulose (Schleicher and Schuell) and blocked for 3 h in 5% (w/v) gelatin in phosphate buffered saline (PBS), pH 7.5. Primary antisera were added in 1% gelatin in PBS containing 0.2% Nonidet P-40 (NP-40) and incubated overnight. The primary antiserum was removed and blots washed extensively with PBS containing 0.2% NP-40. Secondary antiserum (donkey anti-rabbit IgG coupled to horseradish peroxidase; HRP) in 1% gelatin/PBS/0.2% NP-40 was added and left for 3h. After removal of the second antiserum, blots were washed extensively as above and developed with

o-dianisidine hydrochloride as substrate for HRP. The developed immunoblots were scanned and quantified as described previously (Shah *et al.*, 1995).

Reverse transcription/polymerase chain reaction (RT/PCR)

Total RNA was extracted according to the acid phenol/guanidinium thiocyanate method of Chomcynski and Sacchi (1987) using RNAzol B. Purity and quantification of RNA were assessed by the ratio of absorbance at 260 and 280 nm. The RT/PCR procedure was carried out as described previously (Shah *et al.*, 1995b). Briefly, samples of 5 mg RNA (20 ml) were denatured by incubation at 65°C for 10 min followed by chilling on ice and reverse transcribed in 33 ml of reaction mixture using the first strand cDNA synthesis kit (Pharmacia LKB Biotechnology) as detailed by the manufacturer. Incubation was carried out at 37°C for 1 h and the reaction was stopped by heating samples at 95°C for 5 min followed by chilling on ice. PCR reactions were carried out using the following 24-mer primers for the α-subunit of the G-proteins.

Gsα-sense, 5' CCACCTGAATTCGAGCATGCC 3'; Gsα-antisense, 5' GCGTGGGTCTCT CCGGGCTCGGG 3'; Gqα-sense, 5'ATGACTTGACCGTGTAGCCGACC 3' G11α-sense, 5' ACGTGGACCGCATCGCCACAGTAG 3'; Gq/11α-antisense, 5' CCATGC GGTTCCTATTGTCTGACT 3', HPRT-sense, 5' CCTGCTGGATTACATTAAAGCACT 3' and HPRT-antisense, 5' CCTGAAGTACTCATTATAGTCAAG 3'.

Amplifications were performed in 100 µl of buffer containing 20-40 pmol of primers, 2.5 units of Taq polymerase in a HYBAID Omnigene temperature cycler. Cycles were as follows: 95°C/5 min, 60°C/30 s, 72°C/1 min (1 cycle); 95°C/30 s, 60°C/30 s, 72°C/1 min (25-30 cycles); 95°C/30 s, 60°C/30 s, 72°C/5 min (1 cycle). These conditions ensured a linear relationship between the amount of template mRNA and the amount of amplified product (Shah *et al.*, 1995). Either the constitutively expressed housekeeping *hprt* gene (Steel and Buckley, 1993) or β-actin (Mullaney *et al.*, 1995) was selected as a reference template for this purpose. Reaction products were separated by 1.5-1.75% agarose gel electrophoresis. Data analysis was performed using the Kaleidagraph (version 2.1) curve fitting package driven by an Apple Macintosh computer.

Results and Discussion

The membranes prepared from C6 cells exposed to CT (100 ng/ml) and other agents for different time periods were subjected to immunoblotting with antipeptide antiserum which was generated against the α-subunit of Gi and Gq/11 proteins. CT-treatment (100 ng/ml) of C6 cells for 24 or 48 h caused little increase (25 ± 6%; mean

Figure 1. Effect of cholera toxin (CT; 100 ng/ml), forskolin (FSK; 10 μ M) and dibutyryl cAMP (db-cAMP; 1 mM) on immunoreactivity of membrane Gi2 α levels in C6 cells at various time periods. Immunoblots of a representative experiment are shown here (n=4).

Figure 2. (A). Immunological detection of a combination of Gq/G11 α proteins at various time points of CT treatment (100 ng/ml). Membranes prepared from C6 cells were resolved by SDS-PAGE (10% w/v, acrylamide) and immunoblotted using antiserum CQ2 (B). Quantitation of data from Figure 4A (Mean $\pm$ SD, n=4).

and SEM) in the membrane Gi2 α levels (Figure 1). However, CT (100 ng/ml) treatment for 24 and 48 h increased levels of Gq/11 α (40%) as detected by immunoblotting using antiserum which was generated against a region that is completely conserved between the two proteins, Gq α and G11 α (Mitchell *et al.*, 1991) as shown in Figure 2.

Since CT-mediated ribosylation of Gs α results in marked production of cAMP within the cells (Chang and Bourn, 1989), it was tested if the increased intracellular cAMP levels were responsible for effects of CT on G proteins. For that, cells were treated with agents that increase intracellular cAMP like forskolin (FSK) or a cell permeable cAMP analogue (db-cAMP). Treatment of C6 cells with either FSK (10 μ M) or db-cAMP (1 mM) for 48

h caused little increase ($25 \pm 5\%$) in the level of the Gi2 α (Figure 1). The effects of CT on Gq/11 α were not mimicked by treatment with both FSK and db-cAMP (data not shown).

To examine the effect of CT on individual polypeptides (Gq α and G11 α), urea containing SDS-PAGE was used to resolve Gq α and G11 α as described previously (Shah and Milligan, 1994). Quantitation of the CT-induced changes in Gq α and G11 α demonstrated that a similar percentage of each G protein polypeptide was regulated at different time periods (Figure 3). To see whether the changes produced in G proteins are due to alterations in their transcripts, the mRNA levels were measured by RT/PCR (Shah *et al.*, 1995). Results show that CT enhanced the expression of G11 α and

Figure 3. Time-course effect of CT (100 ng/ml) on Gq α and G11 α proteins. Membranes were resolved by SDS-PAGE (12.5% acrylamide/0.0625% bisacrylamide) with a linear gradient of 4-8 M urea and were immunoblotted with antiserum which recognizes both G11 α and Gq α . Under these conditions G11 α migrates more rapidly than Gq α (Shah *et al.*, 1994).

Figure 4. (A): RT/PCR analysis of HPRT (as control), Gs α , Gq α and G11 α mRNA levels in C6 cells treated with CT (100 ng/ml) for various time periods. (B): The RT/PCR data is quantitated by normalization to HPRT mRNA as an internal standard (Mean $\pm$ SD, n=3).

Gq α (60-75%) transcripts without any appreciable change in Gs α mRNA (Figure 4).

CT-induced increase in cAMP occurs as a result of ADP-ribosylation of the α -subunit of stimulatory G-protein (Gs) and thus constitutive activation of adenylyl cyclase (Chang and Bourn, 1989). While CT markedly reduces the membrane Gs α levels, it increases Gi2 α and Gq/11 α levels at 24-48 h of treatment in C6 cells. The downregulation of Gs α was not produced by increased cAMP levels as the effect could not be reproduced by forskolin or cAMP analogue, db-cAMP (Shah, 1997). It has been suggested that ADP-ribosylation of Gs α increases the protein's susceptibility to proteolytic degradation

(Levis and Bourn, 1992; Wedegaertner and Bourn, 1994), mainly through depalmitoylation which causes loosening of Gs α attachment from the cell membrane (Levis and Bourn, 1992, see Mumby, 1997). Such a reduction in the level of G-protein is, in fact, an adaptive mechanism to protect cells from the excessive stimulation of the signalling cascade (Milligan, 1993b). Similar responses are also observed with other G-proteins like Gq α and G11 α upon prolonged receptor stimulation with pharmacological doses of the receptor agonist (Shah and Milligan, 1994; Shah *et al.*, 1995).

Measurement of mRNA levels by RT/PCR showed that the ability of CT to produce a reduction in levels of membrane Gs α was not related to alterations in levels of Gs α mRNA. This is consistent with the observations of Carr *et al.* (1990) but in contrast to findings of Lin *et al.* (1993) in GH3 lactotrophs where the later authors showed a biphasic response to CT; an initial increase and then decline in both Gs α mRNA and protein. Our previous studies in NCB20 cells transfected with β 2-adrenoceptor showed that isoproterenol mediated decrease in Gs α protein was independent of any changes in its transcript (Mullaney *et al.*, 1995). It seems as reduction in membrane Gs α requires either occupancy of receptors with the agonists or needs a direct action (e.g., ADP-ribosylation) on the Gs α protein (Levis and Bourn, 1992).

In C6 cells, the upregulation of Gq/11 α proteins in response to CT (100 ng/ml) was observed only after prolonged exposure. It has been shown that PLC activation and phosphoinositide breakdown can be modulated by the adenylyl cyclase pathway and this effect is mediated by PKA (Haddock *et al.*, 1990; Liu and Simon, 1996). Moreover, cAMP-mediated transcriptional regulation of G-proteins (Hosoda *et al.*, 1994; McKenzie and Milligan, 1990) and the linked receptors (Collins *et al.*, 1989; Thomas *et al.*, 1992) is well documented. Recent studies have shown that cAMP-mediated regulation of adrenergic receptors occur at the level of gene transcription, not mRNA stability (Hosoda *et al.*, 1994). Further

studies in this direction in C6 glioma cells are required.

cAMP exerts multiple effects on various enzymes and thus alters cellular activities. Such effects are tissue-dependent. For example cAMP causes induction of interleukine-6 in C6 cells which corresponds well with the differentiation of these cells (Slegers and Joniau, 1996). Cyclic AMP causes differentiation of C6 cells towards more astrocyte phenotype as reflected by a decrease in neuronal marker glutamate decarboxylase and increased expression of glial fibrillary acidic protein (Messens and Slegers, 1992). Present studies show that upregulation of Gq α and G11 α proteins by CT occurs only after prolonged exposure (24 and 48 h) and this effect seems to be specific to C6 cells as it was not seen in other cells like pituitary gonadotrophs (unpublished results). Similarly induction of heat shock proteins by cAMP occurs only in C6 glioma cells but not in other neuronal cell lines (Kato *et al.*, 1996). Thus, the possibility that regulation of PLC-coupled G proteins (Gq/11 α) through long term stimulation of the adenylyl cyclase pathway is a result of differentiation of C6 cells can not be overlooked.

While exposure to CT downregulated Gs α , it led to upregulation of Gq/11 α and Gi2 α proteins with no significant changes in Go or G-protein β -subunits. Earlier studies in C6 cells (Carr *et al.*, 1990) and NG-108 cells (McKenzie and Milligan, 1990) show no change in Gia protein in response to increase in intracellular cAMP. Contrary to that forskolin and isoproterenol increase the expression of Gi2 α protein in S49 cells (Haddock *et al.*, 1990). Mullaney *et al.* (1988) reported that prolonged treatment for 8 days of neuroblastoma x glioma hybrid cells with dibutyryl cAMP increased the expression of Goa but decreased Gia proteins. Taken together our previous and present studies, it is evident that CT exhibits diverse effects on G-proteins (Gs α , Gi2 α and Gq/11 α) and shows long-term cross regulation between stimulatory Gs/adenylyl cyclase cascade and phosphoinositidase C coupled Gq/11 α proteins.

Acknowledgements

Author acknowledges the support and guidance of Professor Graeme Milligan in conducting this study. This work was started in Professor Milligan's lab at the University of Glasgow, Scotland, UK. Author is also thankful to the Aga Khan University for financial assistance.

References

Berridge, M. J. (1993) Inositol triphosphate and calcium signalling. *Nature* 361: 315-325

Carr, C., Loney, C., Unson, C., Knowler, J. and Milligan, G. (1990) Chronic exposure of rat glioma C6 cells to cholera toxin induces loss of the α -subunit of the stimulatory guanine-nucleotide binding protein (Gs). *Eur. J. Pharmacol.* 188: 203-209

Chang, F.-H., and Bourne, H. R. (1989) Cholera toxin induces cAMP-independent degradation of Gs. *J. Biol. Chem.* 264: 5352-5357

Chomcynski, P. and Sacchi, N. (1987) Single-step method of RNA iso-lation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162: 156-159

Collins, S., Bouvier, M., Bolanowski, M. A., Caron, M. G. and Lefkowitz, R. J. (1989) cAMP stimulates transcription of the β 2-adrenergic receptor gene in response to short term agonist exposure. *Proc. Natl. Acad. Sci. USA* 86: 4853-4857

Exton, J. E. (1997) Cell signalling through guanine-nucleotide-binding regulatory proteins (G proteins) and phospholipases. *Eur. J. Biochem.* 243: 10-20

Eyster, K. M. (1998) Introduction to signal transduction. *Biochem. Pharmacol.* 55: 1927-1938

Haddock, J. R., Ros, M., Watkins, D. C. and Malbon, C. C. (1990) Cross regulation between G-protein mediated pathways. Stimulation of adenylyl cyclase increases expression of the inhibitory G-protein Gi α 2. *J. Biol. Chem.* 265: 14784-14790

Hosoda, K., Feussner, G. K., Rydelek-Fitzgerald, L., Fishman, P. H. and Duman, R. S. (1994) Agonist and cyclic AMP-mediated regulation of β 1-adrenergic receptor mRNA and gene transcription in rat C6 glioma cells. *J. Neurochem.* 63: 1635-1645

Kato, K., Ito, H., Hasegawa, K., Inaguma, Y., Kozawa, O. and Asano, T. (1996) Modulation of the stress-induced synthesis of hsp27 and α B-crystallin by cyclic AMP in rat C6 glioma cells. *J. Neurochem.* 66: 946-950

Levis, M. J. and Bourne, H. R. (1992) Activation of the α -subunit of Gs in intact cells alters its abundance, rate of degradation, and membrane avidity. *J. Cell Biol.* 119: 1297-1307

Lin, J. H., Wang, H. Y., Fong, J. C., Pan, J. T. and Wang, F. F. (1993) Correlation between prolactin secretion and Gs protein expression during sustained cholera-toxin stimulation. *Biochem. J.* 296: 335-340

Liu, M. and Simon, M. I. (1996) Regulation by cAMP-dependent protein kinase of a G-protein-mediated phospholipase C. *Nature* 382: 83-87

Lowery, O. H., Rosenbrough, N. J., Farr, A. L. and Randall, R. J. (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 265-275

McKenzie, F. R. and Milligan, G. (1990) Prostaglandin E1-mediated, cyclic AMP-independent, down-regulation of Gs in neuroblastoma x glioma hybrid cells. *J. Biol. Chem.* 265: 17084-17093

Messens, J. and Slegers, H. (1992) Synthesis of glial fibrillary acidic protein in rat C6 glioma in chemically defined medium: cyclic AMP-dependent transcriptional and translational regulation. *J. Neurochem.* 58: 2071-2080

Milligan, G. (1993a) Regional distribution and quantitative measurement of the phosphoinositidase C-linked guanine nucleotide binding proteins G11 α and Gq α in rat brain. *J. Neurochem.* 61: 845-851

Milligan, G. (1993b) Mechanisms of multifunctional signalling by G-protein linked receptors. *Trends Pharmacol. Sci.* 14: 239-244

Mitchell, F. M., Mullaney, I., Godfrey, P. P., Arkinstall, S. J., Wakelam, M. J. O. and Milligan, G. (1991) Widespread distribution of Gq α /G11 α detected immunologically by an antipeptide antiserum directed against predicted C-terminal decapeptides. *FEBS Lett.* 287: 171-174

Mullaney, I., Magee, A. I., Unson, C. G. and Milligan, G. (1988) Differential regulation of amounts of the guanine-nucleotide-binding proteins Gi and Go in neuroblastoma x glioma hybrid cells in response to dibutyryl cyclic AMP. *Biochem. J.* 256: 649-656

Mullaney, I., Shah, B. H., Wise, A. and Milligan, G. (1995) Expression of the human β 2-adrenoceptor in NCB20 cells results in agonist activation of adenylyl cyclase and agonist-mediated selective down-regulation of Gs α . *J. Neurochemistry* 65: 545-553

Mumby, S. M. (1997) Reversible palmitoylation of signaling proteins. *Curr. Opin Cell Biol.* 9: 148-154

Peterson, J. W., Saini, S. S., Dickey, W. D., Klimpel, G. R., Bomalaski, J. S., Clark, M.

- A., Xu, X.-J. and Chopra, A. K. (1996) Cholera toxin induces synthesis of phospholipase A₂-activating protein. *Infection and Immunity* 64: 2137-2143
- Segovia, J., Lawless, G. M., Tillakaratne, N. J. K., Brenner, M. and Tobin, A. J. (1994) Cyclic AMP decreases the expression of a neuronal marker (GAD67) and increases the expression of an astroglial marker (GFAP) in C6 cells. *J. Neurochem.* 63: 1218-1225
- Shah, B. H. and Milligan, G. (1995) Cyclic AMP differentially regulates the expression of the alpha subunit of Gs, Gq and G11 G proteins and their mRNA levels in rat C6 glioma cells. *Biochem. Soc. Trans.* 22:10S
- Shah, B. H. and Milligan, G. (1994) The gonadotrophin releasing hormone receptor of α T3-1 cells interacts with and regulates phosphoinositidase C-linked G-proteins, Gq and G11 equally. *Mol. Pharmacol.* 46: 1-7
- Shah, B. H., McEwan, D. J. and Milligan, G. (1995) Gonadotrophic releasing hormone receptor agonist-mediated down-regulation of Gq α /G11 α G-proteins in α T3-1 gonadotroph cells reflects increased G-protein turnover but not alterations in mRNA levels. *Proc. Natl. Acad. Sci. USA* 92: 1886-1890
- Slegers, H. and Joniau, M. (1996) Lipopolysaccharide-enhanced expression of interleukine-6 in dibutyl cyclic AMP-differentiated rat C6 glioma. *J. Neurochemistry* 66: 466-473
- Smrcka, A. V., Hepler, J. R., Brown, K. O. and Sternweis, P. C. (1991) Regulation of polyphosphoinositide-specific phospholipase C activity by purified Gq. *Science* 251: 804-807
- Steel, M. C. and Buckley, N. J. (1993) Differential regulation of muscarinic receptor mRNA levels in neuroblastoma cells by chronic agonist exposure: a comparative polymerase chain reaction study. *Mol. Pharmacology* 43: 694-701
- Sunahara, R. K., Dessauer, C. W. and Gilman, A. G. (1996) Complexity and diversity of mammalian adenylyl cyclases. *Ann. Rev. Pharmacol. Toxicol.* 36: 461-480
- Thomas, R. F., Holt, B. D., Schwinn, D. A. and Liggett, S.B. (1992) Long-term agonist exposure induces upregulation of β 3-adrenergic receptor expression via multiple cAMP response elements. *Proc. Natl. Acad. Sci. USA* 89: 4490-4494
- Wedegaertner, P. B. and Bourne, H. R. (1994) Activation and depalmitoylation of Gs α . *Cell* 77: 1063-1070