

α -Tocopheryl succinate potentiates the paclitaxel-induced apoptosis through enforced caspase 8 activation in human H460 lung cancer cells

Soo-Jeong Lim^{1,4*}, Moon Kyung Choi^{2*},
Min Jung Kim¹ and Joo Kyoung Kim³

¹Department of Bioscience and Biotechnology
Sejong University
Seoul 143-747, Korea

²Research Institute
National Cancer Center
Goyang 410-769, Korea

³Seoul Medical Science Institute
Seoul 140-809, Korea

⁴Corresponding author: Tel, 82-2-3408-3767;
Fax, 82-2-3408-3334; E-mail, sjlim@sejong.ac.kr

*These authors contributed equally to this work.
DOI 10.3858/emmm.2009.41.10.080

Accepted 8 June 2009

Abbreviations: AP-1, activator protein-1; Cox, cyclooxygenase; NSCLC, nonsmall cell lung cancer; PARP, poly(ADP-ribose) polymerase; ROS, reactive oxygen species; TOS, α -tocopheryl succinate

Abstract

Paclitaxel is one of the chemotherapeutic drugs widely used for the treatment of nonsmall cell lung cancer (NSCLC) patients. Here, we tested the ability of α -tocopheryl succinate (TOS), another promising anticancer agent, to enhance the paclitaxel response in NSCLC cells. We found that sub-apoptotic doses of TOS greatly enhanced paclitaxel-induced growth suppression and apoptosis in the human H460 NSCLC cell lines. Our data revealed that this was accounted for primarily by an augmented cleavage of poly(ADP-ribose) polymerase (PARP) and enhanced activation of caspase-8. Pretreatment with z-VAD-FMK (a pan-caspase inhibitor) or z-IETD-FMK (a caspase-8 inhibitor) blocked TOS/paclitaxel cotreatment-induced PARP cleavage and apoptosis, suggesting that TOS potentiates the paclitaxel-induced apoptosis through enforced caspase 8 activation in H460 cells. Furthermore, the growth suppression effect of TOS/paclitaxel combination on human H460, A549 and H358 NSCLC cell lines were synergistic. Our observations indicate that combination of paclitaxel and TOS may offer a novel therapeutic strategy for improving paclitaxel drug effi-

cacy in NSCLC patient therapy as well as for potentially lowering the toxic side effects of paclitaxel through reduced drug dosage.

Keywords: apoptosis; carcinoma, non-small-cell lung; caspase 8; paclitaxel; α -tocopherol

Introduction

α -Tocopheryl succinate (TOS), which is obtained by the esterification of α -tocopherol is a more stable powder form of α -tocopherol, since the succinate group protects the hydroxyl group of the chromanol ring from oxidation. Despite their similar chemical structures, TOS differs in activity from α -tocopherol. For example, TOS is redox-insensitive prior to being cleaved to α -tocopherol by intracellular esterases. In contrast to α -tocopherol, TOS induces cytotoxic responses in a wide range of cancer cells while sparing normal cells (Weber *et al.*, 2002; Kang *et al.*, 2004). The antitumor effect of TOS appears to be mediated primarily through the induction of apoptosis. TOS has been shown to induce apoptosis in tumor cell lines via multiple pathways. For example, TOS was observed to induce apoptosis in cancer cells by transcriptional activation of activator protein-1 (AP-1)-controlled genes. In other cell lines, the apoptotic activity of TOS was linked to its blockade of cell cycle progression by inhibiting the transcriptional activity of E2F, or by enhancing the expression of p21^{waf1/cip1}, a cell cycle checkpoint protein (Turley *et al.*, 1997). TOS has also been shown to induce apoptosis by activating Fas signaling pathways (Israel *et al.*, 2000). Recently, the proapoptotic activity of TOS has been linked to its ability to induce the accumulation of reactive oxygen species (Kang *et al.*, 2004; Dong *et al.*, 2008), leading to mitochondrial destabilization (Neuzil *et al.*, 2007). TOS-induced apoptosis appears to be mediated by caspase activation in majority of cancer cell lines, although the contribution of caspase 9 or caspase 8 differs depending on the cellular context (Swettenham *et al.*, 2005).

Recently, considerable attention have been given to combination treatment with reduced doses of two or more different anticancer agents, since combination treatment may enhance the therapeutic efficacy of each agent alone, and/or enable

the use of lower doses of each agent. With this regards, in addition to induction of apoptosis at higher doses, lower doses of TOS acts synergistically with anticancer agents, including selenium, exisulind and SC-560 a selective Cox-1 inhibitor (Lee *et al.*, 2006b; Lim *et al.*, 2007; Reagan-Shaw *et al.*, 2008) to induce apoptosis. The mechanisms supporting the synergistic effects observed in combination treatments remain largely undefined, but involve enhanced apoptosis through enforced caspase activation. These earlier studies suggest a possibility that TOS may be a clinically useful adjuvant in cancer treatment since antitumor effects of chemotherapeutic drugs are mediated at least partly through caspase-mediated apoptosis induction.

Lung cancer is one of leading cause of cancer-related death worldwide. Majority of lung cancer patients are diagnosed with already metastasized or advanced diseases, thus being refractory to current treatment modalities, warranting the needs for developing new preventive or therapeutic drugs. In particular, nonsmall cell lung cancer (NSCLC) is a cancer with poor prognosis despite recent progress made in early diagnosis and treatment. Chemotherapy is an established treatment for advanced NSCLC but still show poor response rates in NSCLC patients (Ihde and Minna, 1991).

Paclitaxel is one of the chemotherapeutic drugs widely used for the treatment of NSCLC. This drug binds and stabilize microtubules, and interferes

with the dynamic changes that occur during the formation of the mitotic spindles, thereby arresting the cells at the G₂/M phase (Honore *et al.*, 2005). Paclitaxel also has the ability to induce apoptosis and its cell killing effects are mediated, at least partly, by inducing apoptosis (Wang *et al.*, 2000). Although there are numerous reports about the mechanism underlying apoptosis induced by paclitaxel, how this microtubule polymerizing agent triggers apoptosis remains incompletely understood. Apoptotic cell death caused by paclitaxel often involves various caspase activation (von Haefen *et al.*, 2003; Park *et al.*, 2004). In NSCLC cells *in vitro*, paclitaxel treatment caused caspase-8- and caspase-3-dependent apoptosis and the apoptosis induction was correlated with the antitumor effect of paclitaxel (Ferreira *et al.*, 2000).

Despite of its potent anticancer activity, the effectiveness of paclitaxel chemotherapy is often limited because of dose-limiting side effects (Zanetta *et al.*, 2000). A major challenge to improving their chemotherapeutic index is by combined treatments that cooperatively or synergistically enhance apoptosis. We herein examined whether TOS and paclitaxel could act synergistically to induce enhanced growth suppression and apoptosis in human H460 NSCLC cells. Our data herein demonstrate that paclitaxell cotreatment with a subapoptotic dose of TOS greatly enhances paclitaxel-induced apoptosis in H460 cells, at least partially through enforced caspase-8 activation.




Figure 1. TOS augmented paclitaxel (TAX)-induced growth suppression in H460 cells. (A) Cells were seeded at a density of 2.5×10^3 /well in 96-well plates and, starting 24 h later, were treated with increasing doses of paclitaxel in the presence or absence of TOS (25 μ M). At 48 h post-incubation, the growth and viability of cells were determined by using MTT. Results are expressed as percentage growth (mean \pm S.D. of triplicate wells) relative to DMSO-treated cells. (B) Effect of combination treatment on H460 colony formation. Cells were plated at 500 cells per dish. The medium covering the cells was replaced every three days at the Taxol concentrations indicated on the figure with or without 2.5 μ M TOS. Colony numbers were counted after 10 days treatment following crystal violet staining. The number of colonies in the dish treated with vehicle alone was used as an index for a 100% survival rate (control) and this value was used to obtain the percentage survival rates for other dishes. Data are mean \pm SD of three independent experiments. Significant differences are indicated by asterisks: * $P < 0.05$ and ** $P < 0.005$, compared with cells treated with paclitaxel alone.

Results

TOS augmented paclitaxel effect on the growth and colony formation of H460 cells

To examine the effect of TOS on paclitaxel-induced growth inhibition, we treated human H460 lung cancer cells with increasing doses of paclitaxel in the presence or absence of TOS. In the absence of TOS, 10 and 100 nM paclitaxel inhibited the growth of H460 cells by 45.1 and 63.3%, whereas same doses of paclitaxel inhibited the growth of cells by 60.5 and 89.6%. (Figure 1A). Incubation with 25 μ M TOS alone showed only a slight inhibitory effect (7.5%, data not shown) on the H460 cell growth.

We performed colony formation assay to evaluate the combination effect by TOS and paclitaxel after long-term treatment. TOS combination increased the paclitaxel-induced inhibition of colony formation in H460 cells by 5.7- and 1.3 fold

compared with paclitaxel alone (Figure 1B). These data collectively indicate that TOS enhanced paclitaxel effect on the growth and colony formation of H460 cells.

TOS did not affect paclitaxel-induced tubulin polymerization and G₂/M arrest

Because the anticancer effect of paclitaxel is known to be initiated by binding to and stabilizing microtubules, we sought to investigate whether TOS combination directly affect paclitaxel-induced tubulin polymerization. The soluble, nonpolymerized tubulin was separated from insoluble, polymerized tubulin and analyzed by immunoblotting using anti-tubulin antibodies. Paclitaxel at 100 nM dose induced polymerization whereas TOS had no effect. The combination of paclitaxel with TOS did not enhance the paclitaxel effect on tubulin polymerization (Figure 2A).

Stabilization of microtubules interferes with the dynamic changes that occur during the formation of the mitotic spindles, thereby arresting the cells at the mitotic phase. When we analyzed the percentage of cells accumulated at G₂/M phase after 36 h paclitaxel treatment, we found that 10 and 100 nM paclitaxel increased the percentage of H460 cells accumulated in the G₂/M phase in a dose-dependent manner. TOS alone slightly increased the percentage of cells in the G₂/M phase. However, TOS combination did not change paclitaxel-induced G₂/M accumulation of cells (Figure 2B). These observations demonstrate that TOS do not affect the paclitaxel effect at the level of tubulin polymerization and mitotic arrest to induce growth suppression.

TOS augmented paclitaxel-induced apoptosis

We then investigated whether the observed augmenting effect of TOS on paclitaxel-induced growth suppression was associated with potentiation of paclitaxel-induced apoptosis. ELISA-based measurements of histone-associated DNA fragments revealed that 100 nM paclitaxel alone increased DNA fragmentation by 4.1-fold whereas combination treatment with TOS did by 10.2-fold (Figure 3A). The increase in the apoptosis induced by 100 nM paclitaxel plus TOS combination was even much higher compared with that by 300 nM paclitaxel alone (10.2 vs 4.8-fold). In addition, we observed an increase in apoptosis-associated morphological changes, including cell shrinkage and chromatin condensation, in cells treated with both TOS and paclitaxel versus cells treated with paclitaxel alone (data not shown). These results




Figure 2. Augmentation by TOS of paclitaxel (TAX)-induced growth inhibition is independent of tubulin polymerization and cell accumulation at G₂/M phase. (A) H460 cells were treated with 100 nM paclitaxel in the presence or absence of 25 μ M TOS for 36 h. Polymerized (cytoskeleton) and non-polymerized (cytosol) tubulin were isolated as described under experimental procedures and resolved on a gel and blotted against anti- β -tubulin and anti- β -actin. (B) Accumulation of cells at G₂/M phase induced by taxol with or without 25 μ M TOS co-treatment. Values represent percentages of accumulated cells in the G₂/M phase and are from three independent experiments.




Figure 3. Augmentation by TOS of paclitaxel (TAX)-induced apoptosis. (A) Exponentially growing H460 cells were treated with the indicated doses of paclitaxel in the absence or presence of TOS (25 μ M) for 42 h. Apoptosis was quantified by an ELISA that specifically detected histone-associated DNA. Bar represents the net absorbances at the suggested wavelengths ($A_{405\text{ nm}}$ minus $A_{490\text{ nm}}$) (mean \pm SD of four experiments performed in triplicate). Significant differences are indicated by asterisks: ** $P < 0.005$ compared with cells treated with paclitaxel alone. (B) Combination treatment induced the cleavage of PARP and decreased the expression of BCL-X_L anti-apoptotic protein without affecting that of BCL-2. The expression of proteins was assessed by immunoblotting, in cells harvested after 30 h treatment. Immunoblotting with an antibody to β -actin was used to control for equal loading of proteins per lane.

strongly suggest that TOS augments paclitaxel-induced apoptosis in H460 cancer cells.

Apoptosis was also evaluated by assaying for cleaved PARP, a general marker of apoptosis. Immunoblotting analysis demonstrated the dramatic increase in the levels of cleaved fragments of PARP by TOS combination with paclitaxel: in the absence of TOS, fragmented PARP was detected after treatment with 100 nM paclitaxel, but not with 10 nM paclitaxel. In contrast, in the presence of TOS, PARP fragment was detected already by 10 nM paclitaxel treatment (Figure 3B). TOS alone did not cause PARP fragmentation. Furthermore, when the expression of Bcl-2 family of apoptosis regulators was assayed by immunoblotting analysis, we found that the expression of Bcl-XL antiapoptotic protein, but not that of Bcl-2 antiapoptotic protein, was decreased by TOS combination (Figure 3B). These data suggest that TOS combination affected the expression profiles of some apoptosis-related proteins.

TOS augmented paclitaxel-induced apoptosis through enforced caspase 8 activation


Apoptosis is generally associated with caspase activation (Lee *et al.*, 2006a), and paclitaxel-induced apoptosis has been shown to involve the caspase activation cascade. To determine whether paclitaxel/TOS cotreatment-induced apoptosis was mediated through enforced caspase activation, we assayed the activities of caspase-8 and -9, two major initiator caspases. The decrease in the

expression of procaspase-8 protein was detected by paclitaxel alone from 100 nM doses but from paclitaxel doses as low as 10 nM by TOS combination (Figure 4A). On the other hand, the expression of procaspase-9 protein was not decreased by paclitaxel treatment regardless of TOS combination (Figure 4A). We next investigated whether paclitaxel/TOS cotreatment induced activation of caspase-3, a major effector caspase. Without TOS combination, the decrease in the expression of procaspase-3 protein was detected by paclitaxel at 100 nM dose, whereas it was detected by 10 nM paclitaxel plus TOS combination. Concomitantly, the catalytically active form of caspase-3 (p17/p19) was evident in H460 cells when they were treated with paclitaxel alone at doses as high as 100 nM, whereas it was detected in cells treated with paclitaxel at doses as low as 10 nM in combination with TOS (Figure 4A).

Figure 4B shows that the activity of caspase-8 was increased by paclitaxel and further increased by paclitaxel and TOS combination, in consistent with immunoblotting analysis data. The activity of caspase-9 was not affected by paclitaxel treatment regardless of TOS combination.

We then examined whether pretreatment of cells with z-LEHD-FMK (a caspase-9 inhibitor), z-IETD-FMK (a caspase-8 inhibitor) or z-VAD-FMK (a pan-caspase inhibitor) could inhibit the apoptotic cell death induced by paclitaxel/TOS cotreatment. Our results revealed that pan-caspase inhibition and specific inhibition of caspase-8 greatly blocked cotreatment-induced apoptosis, as shown by signi-

A



B




Figure 4. TOS combination potentiated paclitaxel (TAX)-induced caspase 8 activation. (A) Combination treatment induced activation of caspase-8 and -3, but not -9, in H460 cells. Expression of active caspase-3 fragment, procaspase-3, -8 and -9 were assayed by immunoblotting of cell lysates obtained from cells treated with indicating concentrations of paclitaxel in the presence or absence of 25 μ M TOS for 30 h. Immunoblotting with an antibody to β -actin was used to control for equal loading of proteins per lane. (B) Cell lysates obtained after 36-h treatment with 50 nM paclitaxel in the presence or absence of 25 μ M TOS were incubated with the reaction buffer containing the appropriate caspase fluorogenic substrate. Reactions were proceeded as described in the text, and fluorescence was determined. Bar represents the ratio of the fluorescence in cells treated as indicated and in control (DMSO) cells. Data represent the average of at least three separate experiments. Significant differences are indicated by asterisks: * $P < 0.05$, compared with cells treated with paclitaxel alone.

ficant decreases in DNA fragmentation (Figure 5A) and PARP cleavage (Figure 5B). z-LEHD-FMK pretreatment only slightly inhibited the cotreatment-induced DNA fragmentation and PARP cleavage (to much less extent compared with z-IETD-FMK or z-VAD-FMK). In addition, pan-caspase inhibition and specific inhibition of caspase-8, but not that of caspase-9, led to the inhibition of caspase 3 activation (Figure 5B), suggesting that paclitaxel/TOS combination induced activation of caspase-8, an initiator caspase, leading to the subsequent activation of caspase-3, an effector caspase. These findings collectively indicate that paclitaxel/TOS cotreatment-induced apoptosis is mediated by and dependent upon caspase-8 activation in H460 cells. We conclude that TOS cotreatment lowered the paclitaxel dose required to induce caspase 8-activation in H460 cells.

TOS also augmented paclitaxel response in other lung cancer cell lines

We sought to examine whether potentiation by TOS on paclitaxel response was also observed in other human NSCLC cells. To statistically evaluate whether the combination of TOS and paclitaxel is synergistic in NSCLC cells, we determined the combination index at two fixed doses of paclitaxel in three NSCLC cell lines. The concentration of TOS for assessing the combination index with paclitaxel was adjusted to a dose only slightly affecting the growth (< 10%) of corresponding cells by itself. Under these conditions, the combination index was less than 1.0 in all three cell lines, suggesting that TOS and paclitaxel are synergistic (Table 1). This finding thus suggests that TOS synergistically acts with paclitaxel to suppress the growth of human NSCLC cells.

Table 1. Combination index of NSCLC cell lines. Synergism was assessed by the method of Chou and Talalay using a nonfixed ratio experimental design and the CalcuSyn software (Biosoft, Ferguson, MO). Combination index values < 1.0 indicates synergism.

Paclitaxel Dose	Combination Index		
	H460	A549	H358
IC ₅₀ ^a	0.784 ± 0.067	0.755 ± 0.082	0.789 ± 0.068
IC ₇₀ ^b	0.419 ± 0.064	0.775 ± 0.049	0.605 ± 0.102

^{a,b}IC₅₀ and IC₇₀ indicates the concentrations inhibiting the cell growth by 50 and 70%, respectively.

Discussion

Our earlier data showed that TOS, a vitamin E analogue and promising, pharmacologically acceptable anticancer agent, enhanced the exisulind-induced growth suppression in human colorectal cancer cells (Lim *et al.*, 2007). In the present report, we show for the first time that TOS, synergistically enhanced paclitaxel response in human NSCLC cell lines. Our findings suggest that TOS/paclitaxel combination treatment, by improving paclitaxel drug efficacy and lowering its toxic side effects through reduced drug dosage, may provide the basis for new combinatory strategies for the effective treatment of NSCLC that have shown poor response to chemotherapy.

Although paclitaxel has also shown to exhibit its anticancer activity mainly by inducing tubulin polymerization and mitotic arrest, our data show that TOS combination did not affect the extent of tubulin polymerization and G₂/M arrest induced by paclitaxel. However, augmentation by TOS of paclitaxel response was correlated with enhanced apoptosis, suggesting that TOS combination affected paclitaxel response mainly at the apoptosis level. Enhanced apoptosis induced by TOS/paclitaxel combination was accompanied by changes in the expression profile of Bcl-XL antiapoptotic protein. Overexpression of antiapoptotic proteins including Bcl-XL in cancer cells has been recognized as one plausible mechanism of developing resistance to paclitaxel (Lebedeva *et al.*, 2000; Williams *et al.*, 2005), and the inhibition of Bcl-XL by genetic or pharmacological approaches has been shown to sensitize cancer cells to paclitaxel (Shoemaker *et al.*, 2006). With this regards, the significance of Bcl-XL downregulation by TOS/paclitaxel combination observed here deserves further speculation.

Since previous studies from our lab and other groups have shown that the accumulation of intracellular reactive oxygen species (ROS) was an early and crucial step for paclitaxel- or TOS-induced apoptotic cell death in human cancer cells




Figure 5. Caspase-8 inhibition blocked the apoptosis induced by paclitaxel and TOS combination. (A) Effect of pretreatment with caspase-9 inhibitor z-LEHD-FMK, caspase-8 inhibitor z-IETD-FMK or the pan-caspase inhibitor z-VAD-FMK on the apoptosis induced by TOS and paclitaxel combination. After 2-h pretreatment with 25 μ M of each caspase inhibitor, H460 cells were treated with paclitaxel (100 nM) and TOS (25 μ M) for 30 h, and subjected to apoptosis ELISA assay. Bar represents the net absorbances at the suggested wavelengths ($A_{405\text{ nm}} - A_{490\text{ nm}}$) (mean \pm SD of four experiments performed in triplicate). Data represent the average \pm SD of two separate experiments. Significant differences are indicated by asterisks: ** $P < 0.01$, compared with cells treated with TOS plus taxol. (B) Effect of pretreatment with caspase inhibitors on the PARP cleavage and caspase-3 activation induced by combination treatment with TOS and paclitaxel. After 2-h pretreatment with 25 μ M of each inhibitor, H460 cells were treated with paclitaxel (100 nM) and TOS (25 μ M) for 30 h, and subjected to immunoblot analysis. Two additional studies yielded similar results. Paclitaxel and TOS combination treatment was abbreviated as T+T in the Figure.

(Kang *et al.*, 2004; Alexandre *et al.*, 2006), we further examined whether TOS cooperated with paclitaxel in the ROS production to increase apoptosis. However, we found that TOS had no effect on paclitaxel-induced ROS generation and that preincubation of cells with antioxidants did not

block the TOS/paclitaxel-induced apoptosis (unpublished data), suggesting that that ROS generation does not play a role in the synergistic apoptosis induced by TOS/paclitaxel cotreatment.

Our data indicate that the potentiation by TOS of paclitaxel-induced apoptosis was mediated through enforced activation of caspases, in particular that of caspase-8. In our study, paclitaxel induced the activation of caspase-8 and -3 but not that of caspase-9 in H460 cells regardless of TOS combination. This is consistent with previous data from other groups using H460 cells (Ferreira *et al.*, 2000). Considering that caspase-9 is also generally involved in mediating paclitaxel-induced apoptosis in many other cell lines, we cannot exclude a possibility that specific TOS effect on caspase-8 activation may be limited to H460 cell lines. Future work will be required to address whether TOS also enhances paclitaxel-induced caspase 9 activation by using other cell lines.

Although the activity of TOS has been translated to cancer suppression in a variety of preclinical models (Hahn *et al.*, 2006), the *in vivo* antitumor activity of TOS have shown to be compromised if orally administered due to the hydrolysis by esterases in the gastrointestinal tract. Currently, researchers are seeking to develop structural analogues of tocopheryl succinate having more potent anticancer activity after oral administration (Dong *et al.*, 2009). Since our data suggest that TOS holds promising therapeutic potential as an anticancer agent to combine with paclitaxel to improve chemotherapy of NSCLC, it would be of interest to investigate whether TOS or its structural analogue in combination with paclitaxel exert synergistic growth inhibitory effect *in vivo*.

Methods

Cell lines and cultures

The human H460 and A549 lung cancer cell lines were purchased from the American Type Culture Collection (ATCC, Manassas, VA), and the human H358 cell line was purchased from the Korean Cell Line Bank (Seoul, Korea). Cells were cultured in RPMI 1640 medium supplemented with 10% heat-inactivated FBS (Hyclone, Logan, UT) and 100 units/ml each of penicillin and streptomycin. All cells were grown in incubators in a humid atmosphere of 95% air/5% CO₂.

Reagents and antibodies

α -tocopheryl succinate (TOS) and MTT were obtained from Sigma (St. Louis, MO). paclitaxel was obtained from Tocris (Tocris Cookson, MO). z-VAD-FMK (pan-caspase inhibitor), z-IETD-FMK (caspase-8 inhibitor) and z-LEHD-FMK (cas-

pase-9 inhibitor) were obtained from Enzyme Systems Products (Livermore, CA). Ac-IETD-AFC (caspase-8 substrate) and Ac-LEHD-AFC (caspase-9 substrate) were purchased from Alexis Biochemicals (Lausen, Switzerland). Antibodies to human poly(ADP-ribose)polymerase (PARP) and Bcl-X were purchased from BD Pharmingen (San Diego, CA) and antibodies to active caspase 3 and procaspase-3, -8 and -9 were obtained from Cell Signaling Technologies (Beverly, MA). Antibody to β -tubulin was purchased from Oncogene Research Products (San Diego, CA). Antibodies to Bcl-2 and β -actin were obtained from Santa Cruz Biotechnology (CA) and Sigma (St. Louis, MO).

All other chemicals were of reagent grade and used without further purification.

Cell viability analysis

Cells ($2.5-5 \times 10^3$ /well in 0.1 ml medium) were seeded into 96-well plates. Starting 24 h later, cells were treated with various doses of paclitaxel or DMSO vehicle in the presence or absence of 25 μ M TOS (stock in DMSO at 50 mM). At 72 h postincubation, the growth and viability of cells were determined by using MTT (Sigma). The ability of cells to form formazan crystals by active mitochondrial respiration was determined using a microplate spectrophotometer (Molecular Devices) after dissolving the crystals in DMSO.

Colony formation assay

500 cells were seeded in 5 cm diameter dishes and incubated for 16 h at 37°C in an atmosphere containing 5% CO₂. Cells were then treated with increasing concentrations of paclitaxel in the presence or absence of TOS. The medium containing treatments was replaced every three days. After 10 days, surviving cells forming visible colonies (containing > 50 cells) were counted after staining with crystal violet in 60% methanol.

Immunoblotting

At time of harvest, treated cells were scraped from the plates, washed twice with ice-cold PBS, and pelleted at $2,400 \times g$ for 5 min. The cell pellet was then incubated for 15-30 min on ice in the lysis buffer containing 150 mM NaCl, 10 mM Tris, 0.2% Triton X-100, 0.3% NP-40, 0.2 mM Na₃VO₄ and protease inhibitors (Roche Diagnostics), pH 7.4. After centrifugation at 14,000 rpm for 15 min at 4°C, the supernatants were collected, and the protein concentration in each was measured by the Bradford method. Aliquots of supernatants containing equal amounts of protein were boiled in SDS-reducing buffer for 5 min, electrophoresed on SDS-polyacrylamide gels, and transferred to nitrocellulose membranes. The membranes were blocked with 5% nonfat dry milk and probed with specific primary antibodies, followed by incubation with appropriate peroxidase-conjugated secondary antibodies. The blots were developed with ECL Plus reagent (Amersham, Arlington Heights, IL) according to the manufacturer's protocol.

Tubulin polymerization assay

To evaluate tubulin polymerization, polymerized and non-polymerized tubulins were extracted as described elsewhere (Sackett *et al.*, 1997). Cells were plated in six-well plates in RPMI supplemented with 10% FBS and allowed to adhere overnight. Indicated compounds were added to the medium and cells were grown for 36 h. Cells were washed twice with PBS and lysed at 37°C for 5 min in the dark with 100 μ l of hypotonic buffer (1 mM MgCl₂, 2 mM EGTA, 0.5% Nonidet P-40, 2 mM PMSF, 200 units/ml aprotinin, 100 μ g/ml soybean trypsin inhibitor, 5 mM ϵ -smino capronic acid, 1 mM benzamidine, 20 mM Tris-HCl, pH 6.8). The lysates were rinsed with 100 μ l of hypotonic buffer and centrifuged at 14,000 rpm for 10 min at room temperature. Supernatants containing soluble (cytosolic) tubulin were separated from the pellets containing polymerized (cytoskeletal) tubulin. The pellets were resuspended in 200 μ l of hypotonic buffer. Soluble and polymerized tubulins were extracted, and respective levels were assessed by immunoblotting as described above using anti- α -tubulin antibodies.

Cell cycle analysis

After treatment, cells were harvested, washed, fixed in 80% ethanol for 30 min, and resuspended in PBS (pH 7.4) containing 0.1% Triton X-100, 5 μ g/ml propidium iodide, and 50 μ g/ml ribonuclease A for DNA staining. The cell-cycle distribution was analyzed using FACScan (Becton-Dickinson, San Jose, CA) and Modfit 3.0 (Verity Software, Topsham, ME) cell-cycle analysis software.

Apoptosis ELISA assay

Cytoplasmic histone-associated DNA fragments (mono- and oligo nucleosomes) were quantified by a photometric enzyme immunoassay using Cell Death Detection ELISA^{plus} (Roche Applied Bioscience) following the manufacturer's protocol. Briefly, after treatment, cells were pooled and lysed. Cytoplasmic fractions containing histone/DNA fragments were extracted and adhered to an immobilized anti-histone antibody plate. A peroxidase-conjugated anti-DNA antibody was used for detection of adhered histone/DNA fragments. A colorimetric substrate for peroxidase was then added to each well containing each sample. The net absorbances at the suggested wavelengths ($A_{405\text{ nm}}$ minus $A_{490\text{ nm}}$) were obtained for the lysates derived from treated as well as control cells.

Caspase activity assay

Caspase-8- and -9-like activity was measured by cleavage of fluorogenic caspase substrates, Ac-IETD-AFC or Ac-LEHD-AFC. Briefly, 2×10^5 cells were seeded in 60 cm² dishes, adhered for 24 h. After treatment, detached and attached cells were harvested and lysed on ice in lysis buffer (50 mM Tris/HCl (pH 7.4), 1 mM EDTA, and 10 mM EGTA, 10 μ M digitonin) to generate soluble whole cell lysates. The protein concentrations in cell lysates were determined using Bio-Rad protein assay (Bio-Rad Laboratories). Approximately 200 μ g of total protein were added to

the reaction buffer (20 mM HEPES, 50 mM NaCl, 10 mM dithiothreitol, pH 7.5) containing Ac-IETD-AFC or Ac-LEHD-AFC (50 μ M). Reactions were incubated for 2 h at 37°C, and the levels of released AFC was determined using the Gemini EM microplate spectrofluorometer (Molecular Devices) that allowed for light excitation at 400 nm wavelength and emission at a wavelength of 505 nm. The specific caspase activity, normalized for total proteins of cell lysates, was then expressed as fold of the baseline caspase activity of vehicle-treated control cells.

Statistical analysis

Statistically significant differences between values obtained under different experimental conditions were determined using two-tailed unpaired Student's *t*-tests. Synergism was assessed by the method of Chou and Talalay (1984) using a nonfixed ratio experimental design and the Calcsyn software (Biosoft, Ferguson, MO). Combination index values < 1.0 indicates synergism.

Acknowledgements

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2006-311-E00621).

References

- Alexandre J, Batteux F, Nicco C, Chereau C, Laurent A, Guillemin L, Weill B, Goldwasser F. Accumulation of hydrogen peroxide is an early and crucial step for TAX-induced cancer cell death both *in vitro* and *in vivo*. *Int J Cancer* 2006;119:41-8
- Chou TC, Talalay P. Quantitative analysis of dose-effect relationships: the combined effects of multiple drugs or enzyme inhibitors. *Adv Enzyme Regul* 1984;22:27-55
- Dong LF, Low P, Dyason JC, Wang XF, Prochazka L, Witting PK, Freeman R, Swettenham E, Valis K, Liu J, Zabalova R, Turanek J, Spitz DR, Domann FE, Scheffler IE, Ralph SJ, Neuzil J. α -tocopheryl succinate induces apoptosis by targeting ubiquinone-binding sites in mitochondrial respiratory complex II. *Oncogene* 2008;27:4324-35
- Dong LF, Freeman R, Liu J, Zabalova R, Marin-Hernandez A, Stantic M, Rohlena J, Valis K, Rodriguez-Enriquez S, Butcher B, Goodwin J, Brunk UT, Witting PK, Moreno-Sanchez R, Scheffler IE, Ralph SJ, Neuzil J. Suppression of tumor growth *in vivo* by the mitocan α -tocopheryl succinate requires respiratory complex II. *Clin Cancer Res* 2009;15:1593-600
- Ferreira CG, Span SW, Peters GJ, Krzyt FA, Giaccone G. Chemotherapy triggers apoptosis in a caspase-8-dependent and mitochondria-controlled manner in the non-small cell lung cancer cell line NCI-H460. *Cancer Res* 2000;60:7133-41
- Hahn T, Szabo L, Gold M, Ramanathapuram L, Hurley LH, Akporiaye ET. Dietary administration of the proapoptotic vitamin E analogue α -tocopheryloxyacetic acid inhibits metastatic murine breast cancer. *Cancer Res* 2006;66:

9374-8

Honore S, Pasquier E, Braguer D. Understanding microtubule dynamics for improved cancer therapy. *Cell Mol Life Sci* 2005;62:3039-56

Ihde DC, Minna JD. Non-small cell lung cancer. Part II: Treatment. *Curr Probl Cancer* 1991;15:105-54

Israel K, Yu W, Sanders BG, Kline K. Vitamin E succinate induces apoptosis in human prostate cancer cells: role for Fas in vitamin E succinate-triggered apoptosis. *Nutr Cancer* 2000;36:90-100

Kang YH, Lee E, Choi MK, Ku JL, Kim SH, Park YG, Lim SJ. Role of reactive oxygen species in the induction of apoptosis by α -tocopheryl succinate. *Int J Cancer* 2004;112: 385-92

Lebedeva I, Rando R, Ojwang J, Cossum P, Stein CA. Bcl-xL in prostate cancer cells: effects of overexpression and down-regulation on chemosensitivity. *Cancer Res* 2000;60: 6052-60

Lee E, Choi MK, Han IO, Lim SJ. Role of p21CIP1 as a determinant of SC-560 response in human HCT116 colon carcinoma cells. *Exp Mol Med* 2006a;38:325-31

Lee E, Choi MK, Youk HJ, Kim CH, Han IO, Yoo BC, Lee MK, Lim SJ. 5-(4-chlorophenyl)-1-(4-methoxyphenyl)-3-trifluoromethylpyrazole acts in a reactive oxygen species-dependent manner to suppress human lung cancer growth. *J Cancer Res Clin Oncol* 2006b;132:223-33

Lim SJ, Lee YJ, Park DH, Lee E, Choi MK, Park W, Chun KH, Choi HG, Cho JS. α -tocopheryl succinate sensitizes human colon cancer cells to exisulind-induced apoptosis. *Apoptosis* 2007;12:423-31

Neuzil J, Dong LF, Ramanathapuram L, Hahn T, Chladova M, Wang XF, Zabalova R, Prochazka L, Gold M, Freeman R, Turanek J, Akporiaye ET, Dyason JC, Ralph SJ. Vitamin E analogues as a novel group of mitocans: anti-cancer agents that act by targeting mitochondria. *Mol Aspects Med* 2007;28:607-45

Park SJ, Wu CH, Gordon JD, Zhong X, Emami A, Safa AR. Taxol induces caspase-10-dependent apoptosis. *J Biol Chem* 2004;279:51057-67

Reagan-Shaw S, Nihal M, Ahsan H, Mukhtar H, Ahmad N. Combination of vitamin E and selenium causes an induction of apoptosis of human prostate cancer cells by enhancing Bax/Bcl-2 ratio. *Prostate* 2008;68:1624-34

Sackett DL, Giannakakou P, Poruchynsky M, Fojo A. Tubulin from paclitaxel-resistant cells as a probe for novel anti-microtubule agents. *Cancer Chemother Pharmacol* 1997; 40:228-32

Shoemaker AR, Oleksijew A, Bauch J, Belli BA, Borre T, Bruncko M, Deckwirth T, Frost DJ, Jarvis K, Joseph MK, Marsh K, McClellan W, Nellans H, Ng S, Nimmer P, O'Connor J M, Oltersdorf T, Qing W, Shen W, Stavropoulos J, Tahir SK, Wang B, Warner R, Zhang H, Fesik SW, Rosenberg SH, Elmore SW. A small-molecule inhibitor of Bcl-XL potentiates the activity of cytotoxic drugs *in vitro* and *in vivo*. *Cancer Res* 2006;66:8731-9

Swettenham E, Witting PK, Salvatore BA, Neuzil J. α -tocopheryl succinate selectively induces apoptosis in neuroblastoma cells: potential therapy of malignancies of the nervous system? *J Neurochem* 2005;94:1448-56

Turley JM, Ruscetti FW, Kim SJ, Fu T, Gou FV, Birchenall-Roberts MC. Vitamin E succinate inhibits proliferation of BT-20 human breast cancer cells: increased binding of cyclin A negatively regulates E2F transactivation activity. *Cancer Res* 1997;57:2668-75

von Haefen C, Wieder T, Essmann F, Schulze-Osthoff K, Dorken B, Daniel PT. Paclitaxel-induced apoptosis in BJAB cells proceeds via a death receptor-independent, caspases-3/-8-driven mitochondrial amplification loop. *Oncogene* 2003;22:2236-47

Wang TH, Wang HS, Soong YK. Paclitaxel-induced cell death: where the cell cycle and apoptosis come together. *Cancer* 2000;88:2619-28

Weber T, Lu M, Andera L, Lahm H, Gellert N, Fariss MW, Korinek V, Sattler W, Ucker DS, Terman A, Schroder A, Erl W, Brunk UT, Coffey RJ, Weber C, Neuzil J. Vitamin E succinate is a potent novel antineoplastic agent with high selectivity and cooperativity with tumor necrosis factor-related apoptosis-inducing ligand (Apo2 ligand) *in vivo*. *Clin Cancer Res* 2002;8:863-9

Williams J, Lucas PC, Griffith KA, Choi M, Fogoros S, Hu YY, Liu JR. Expression of Bcl-xL in ovarian carcinoma is associated with chemoresistance and recurrent disease. *Gynecol Oncol* 2005;96:287-95

Zanetta G, Fei F, Mangioni C. Chemotherapy with paclitaxel, ifosfamide, and cisplatin for the treatment of squamous cell cervical cancer: the experience of Monza. *Semin Oncol* 2000;27(1 Suppl 1):23-7