

REVIEW

Regulatory T-cell vaccination independent of auto-antigen

David W Pascual¹, Xinghong Yang¹, Kathryn Holderness², SangMu Jun¹, Massimo Maddaloni¹ and Irina Kochetkova²

To date, efforts to treat autoimmune diseases have primarily focused on the disease symptoms rather than on the cause of the disease. In large part, this is attributed to not knowing the responsible auto-antigens (auto-Ags) for driving the self-reactivity coupled with the poor success of treating autoimmune diseases using oral tolerance methods. Nonetheless, if tolerogenic approaches or methods that stimulate regulatory T (T_{reg}) cells can be devised, these could subdue autoimmune diseases. To forward such efforts, our approach with colonization factor antigen I (CFA/I) fimbriae is to establish bystander immunity to ultimately drive the development of auto-Ag-specific T_{reg} cells. Using an attenuated *Salmonella* vaccine expressing CFA/I fimbriae, fimbriae-specific T_{reg} cells were induced without compromising the vaccine's capacity to protect against travelers' diarrhea or salmonellosis. By adapting the vaccine's anti-inflammatory properties, it was found that it could also dampen experimental inflammatory diseases resembling multiple sclerosis (MS) and rheumatoid arthritis. Because of this bystander effect, disease-specific T_{reg} cells are eventually induced to resolve disease. Interestingly, this same vaccine could elicit the required T_{reg} cell subset for each disease. For MS-like disease, conventional $CD25^{+}$ T_{reg} cells are stimulated, but for arthritis $CD39^{+}$ T_{reg} cells are induced instead. This review article will examine the potential of treating autoimmune diseases without having previous knowledge of the auto-Ag using an innocuous antigen to stimulate T_{reg} cells via the production of transforming growth factor- β and interleukin-10.

Experimental & Molecular Medicine (2014) 46, e82; doi:10.1038/emm.2014.4; published online 14 March 2014

Keywords: autoimmunity; bacteria; fimbriae; mucosal immunity; regulatory T cells; vaccine

COLONIZATION FACTOR ANTIGEN I (CFA/I) FIMBRIAE AND ENTEROTOXIGENIC *ESCHERICHIA COLI* (ETEC)

ETEC, the causative agent of travelers' diarrhea, is the most common bacterial diarrheal disease of children in Latin America, Asia and Africa,¹ and it is contracted upon ingestion of contaminated food or water. It is estimated that this disease is responsible for 400 million cases annually resulting in 300 000 deaths of preschool children.^{2,3} *E. coli* becomes enterotoxigenic upon acquisition of a plasmid or plasmids containing the heat-stable enterotoxin⁴ or the cholera-like exotoxin, which is commonly termed the heat-labile enterotoxin (LT).^{5,6} Both toxins are responsible for inducing fluid loss and electrolyte imbalance in the host. Facilitating infection and subsequent colonization, ETEC also acquires a plasmid encoding for the pili or fimbriae referred to as colonization factor antigens (CFAs), which mediate the colonization of *E. coli* in the gastrointestinal tract. The CFA pili

are a heterogeneous group of fimbrial adhesins responsible for adherence to small intestinal epithelial cells via their fimbriae or long, hairlike projections extending from the bacterial cell surface to epithelial mannose-containing glycoproteins.⁷ This adherence is generally host-specific for intestinal epithelium.⁸ While a specific natural receptor for CFA/I fimbriae has yet to be identified in the small intestine, some studies suggest that in eukaryotes a sialylated glycoprotein is the receptor,^{9,10} although others suggest that epithelial mannose-containing glycoproteins and/or glycosphingolipids may also serve as receptors.^{11,12} The low incidence rates in adults from ETEC-endemic regions have correlated with the presence of anti-LT and anti-CFA antibodies (Abs), suggesting that acquired immunity to these virulence factors are protective.¹³ Epidemiological studies show that children aged <3 years from these endemic regions are susceptible to multiple ETEC infections, which may provide for broad-spectrum immunity

¹Department of Infectious Diseases and Pathology, College of Veterinary Medicine, University of Florida, Gainesville, FL, USA and ²Department of Immunology and Infectious Diseases, Montana State University, Bozeman, MT, USA

Correspondence: Dr DW Pascual, Department of Infectious Diseases and Pathology, College of Veterinary Medicine, University of Florida, P.O. Box 110880, Gainesville, FL 32611, USA.

E-mail: pascuald@ufl.edu

Received 22 November 2013; accepted 6 December 2013

later in life,^{14,15} while adult travelers to these endemic regions are unprotected and remain susceptible to infection.^{2,3,16,17}

CFA/I fimbriae are an archetype of class 5 fimbriae characterized by a common four-gene operon.^{18,19} For CFA/I, this is *cfaABCE*.¹¹ This operon contains four genes in the following order: periplasmic chaperone *cfaA*, major fimbrial subunit *cfaB*, outer membrane usher protein *cfaC*, and minor fimbrial subunit *cfaE*.^{11,18} CfaA functions as a chaperone to facilitate proper folding of other components of the operon to the outer membrane, while CfaC 'ushers' the structural fimbrial proteins and orchestrates their assembly at the cell surface. The extracellular portion of CFA/I fimbriae comprises two proteins, CfaB and CfaE, and assembles with a single copy of CfaE followed by multiple copies of CfaB.^{19,20} In fact, CfaB is the major pilin subunit, and it is present as approximately 1000 copies on the cell surface per single copy of the minor subunit, CfaE.¹⁹

CFA/I FIMBRIAE ARE HIGHLY IMMUNOGENIC AND CONFER PROTECTION AGAINST ETEC

Attempts to successfully vaccinate against ETEC have met with varied success. Oral vaccination of human volunteers with CFA/I or CFA/II fimbriae failed to induce significant serum immunoglobulin G (IgG) or secretory IgA (SIgA) Abs.²¹ As a result of poor anti-fimbriae Ab titers,^{22,23} the human volunteers were not protected against virulent ETEC.²² Despite neutralization of their stomach acidity,²³ poor SIgA anti-CFA Ab responses were obtained in these volunteers. Subsequent work showed that gastric proteases altered the CFA fimbriae antigenicity even at a neutral pH.²⁴ Overcoming the deleterious effects of the gastrointestinal tract, oral vaccination of rabbits with microencapsulated CFA/I fimbriae still failed to induce serum and fecal IgA anti-CFA/I Abs.²¹ Although in a separate study, microencapsulated CFA/II fimbriae when directly intubated into the rabbit duodenum revealed that CFA/II fimbriae-specific Ab-forming cell responses could be induced in the Peyer's patches and spleens.²⁵ Once elicited, Abs to these fimbriae do protect against ETEC infection.^{26,27} An effective vaccine for ETEC still remains elusive, but promising results have been obtained from recent human trials using heat-killed ETEC plus recombinant LT-B/CT-B,¹ suggesting the fimbriae are optimally immunogenic when associated with the bacilli.

Attenuated, live *Salmonella* vectors have been extensively used as a means to vaccinate against salmonellosis^{28–31} and heterologous diseases.³² Such attenuated *Salmonella* strains have been shown to be effective in delivering heterologous antigens (Ags) because of their ability to stimulate both mucosal and systemic immune compartments^{29,33} most likely via infection of Peyer's patches and followed by subsequent spread into systemic immune compartment.^{31,34} Therefore, an ETEC vaccine adapted as an attenuated, balanced-lethal *ΔaroA Δasd* *S. Typhimurium* vaccine carrying an *asd*⁺ plasmid encoding the CFA/I operon.³⁵ The expression of this fimbriae appears similar^{36,37} to wild-type ETEC with long, hairlike projections extending from the bacterial cell surface.²³ Upon

oral immunization, *Salmonella*-CFA/I was quite adept in stimulating elevated mucosal IgA and serum IgG Abs to the fimbriae.^{35,38} Interestingly, the CFA/I fimbriae stimulated a biphasic T helper (Th) cell response with a rapid induction of Th2 cells within the first week of vaccination followed by a progressively increasing Th1 cell response to eliminate the salmonellae.³⁸ This was corroborated by the stimulation of elevated serum IgG1 Abs relative to IgG2a shortly after vaccination.³⁸ This is atypical to immune responses to *Salmonella* vaccines, which generally are Th1 cell-dependent promoting serum IgG2a Abs.^{39–48} Despite this obvious difference in Th cell profiles, the expression of CFA/I fimbriae did not alter its capacity to protect against wild-type *Salmonella* challenge.⁴⁹ Given the highly proinflammatory nature of *Salmonella*,^{39,42,50} subsequent analysis assessed whether the CFA/I fimbriae interfered with the normal recognition of the bacilli. Upon infection of RAW264.7 or thioglycolate-induced macrophages with low infection ratios of *Salmonella*-CFA/I, minimal-to-no interleukin (IL)-1 α , IL-1 β , IL-6 and tumor necrosis factor (TNF)- α production was observed in contrast to its isogenic *Salmonella* vector strain eliciting all of these proinflammatory cytokines with as few as one bacterium/80 macrophages.⁵¹ To ascertain why such a profound disparity in proinflammatory cytokine production, subsequent analysis could not find any differences in *Salmonella* colonization or increased susceptibility to macrophage cell death nor were there increases in anti-inflammatory IL-10 or IL-12p40 cytokines.⁵¹ Although the mechanism for the stealth-like qualities of *Salmonella*-CFA/I was not discerned, a possible explanation for these observations may be that the CFA/I fimbriae thwart innate immune responses by hindering detection by individual or a combination of pathogen-recognition receptors, including Toll-like receptor 4 (TLR4), TLR5, CD14, MD2 and lipopolysaccharide-binding protein,⁵² to indicate *Salmonella*'s presence. Hence, we hypothesized that the *Salmonella*-CFA/I may be an anti-inflammatory vaccine.

SALMONELLA-CFA/I AS AN ANTI-INFLAMMATORY VACCINE FOR AN ANIMAL MODEL OF MULTIPLE SCLEROSIS (MS)

The surprising results from the macrophage infection studies indicated that *Salmonella*-CFA/I was not eliciting the proinflammatory arm of immunity. Moreover, the stimulation of Th2 cells and associated anti-inflammatory cytokines suggests that this vaccine may also be therapeutic in treating autoimmune diseases, such as MS or arthritis. MS is an inflammatory demyelinating disease of the central nervous system (CNS) with destruction of white matter by autoreactive T cells.^{53–56} This neurodegenerative disease affects as many as 400 000 people in United States and >2 million people worldwide.⁵⁷ The frequency of MS is thought to be age and gender dependent, as it most frequently affects young and middle-aged adults and occurs twice as often in females as in males.^{54,57,58} Although not considered fatal, MS can progress into considerable neurological disability, impacting the quality of life, and resulting in a shortened lifespan.⁵⁹

Experimental autoimmune encephalomyelitis (EAE) is one of the best and most frequently studied rodent model that mimics the neuropathology and clinical disease of MS.^{60–65} EAE manifests as an ascending disease in the spinal cord where initial symptoms begin as a limp or paralyzed tail, followed by rear leg paralysis that can eventually progress into forearm paralysis.⁶⁶ It is induced upon immunization with restricted CNS peptides, such as myelin oligodendrocyte glycoprotein,⁶⁷ myelin basic protein⁶⁷ or proteolipid protein (PLP),⁶⁸ into susceptible mice. This injection results in the activation of myelin-specific CD4⁺ T cells in naive animals where in the CNS inflammatory cells are recruited that secrete interferon (IFN)- γ , TNF- α and IL-1, resulting in perpetuation of inflammation along with tissue damage, including axonal damage, demyelination and perivascular inflammatory lesions.⁶⁹ Due to demyelination, new epitopes become exposed, and exposure of these neoAgs acts as an immunization process, thereby causing further epitope spreading.^{62,70–72} Studies have shown that the proinflammatory-promoting cytokine, IL-23, is primarily responsible for encephalitogenic T-cell development in EAE.^{73–75} IL-17 is the principal mediator of the inflammation observed in EAE^{76–79} and in large part induced by IL-23.^{73–76} IL-17 is cross-regulated by both IL-4 and IFN- γ .⁷⁶ Neutralization of IL-17 has been shown to be protective,^{76–79} and protection can also be mediated via regulatory T (T_{reg}) cells.^{80–83} The proinflammatory cytokines IFN- γ ^{84–87} and TNF- α ^{73,74,82,85,88} can also impact disease primarily via Th1 cells. Protection to EAE correlates with Th2/T_{reg} cell dependence as IL-4,^{80,89–91} IL-10^{71,72,85,92–94} and transforming growth factor (TGF)- β ^{80,82,89,95–97} can reverse or prevent EAE. IL-27 has also been shown to be important for induction of protective T_{reg} cells.^{98–100}

As for many autoimmune diseases, patients have lost the capacity to be tolerized to self, and consequently, the patient begins to mount an immune response to self much like recognition of foreign Ags. This T-cell dysfunction exhibited in these patients results in chronic activation of inflammatory Th1 and Th17 cells.^{101,102} Often the course of treatment involves anti-inflammatory drugs and does not address the cause of the disease. Certainly, an approach that has had experimental success is the induction of tolerance. Tolerance is the inability to recognize self or to defined Ags (reviewed in Faria and Weiner,¹⁰³ Mueller¹⁰⁴ and Bilate and Lafaille¹⁰⁵) and past efforts to induce tolerance in humans by delivering Ags have proven successful for treating allergies.^{106–108} The major obstacle in human tolerance is in part attributed to the requirement for large doses of Ag or repeated administrations of therapeutics.¹⁰⁶ Although feeding auto-Ags is effective in treating experimental autoimmune diseases, when applied to patients, oral feeding with auto-Ags has been deemed unsuccessful.^{109,110}

As noted, the *Salmonella*-CFA/I vaccine elicited Th2-type responses to the fimbriae³⁸ much like an effect of soluble Ag with adjuvant.¹² The observed Th2 cell bias was directed to the fimbriae but not as much as to the *Salmonella* vector, which

still retained the capacity to stimulate Th1-type responses against *Salmonella* Ags.¹¹¹ With this backdrop, subsequent studies began to analyze the potential of *Salmonella*-CFA/I to augment disease, first with PLP_{139–151}-induced EAE.¹¹² As *Salmonella*-CFA/I elicits a biphasic Th cell response, groups of mice were orally vaccinated 1 week or 4 weeks before EAE challenge to coincide with a more Th2 cell- or Th1 cell-prone environment, respectively. In either case, *Salmonella*-CFA/I still conferred protection recovering completely from disease unlike those mice vaccinated with an isogenic *Salmonella* vaccine lacking CFA/I fimbriae or those mice treated with phosphate-buffered saline (PBS).¹¹² However, mice vaccinated with the isogenic *Salmonella* vector did show reduced disease relative to PBS-treated mice but still bore greater disease than those mice vaccinated with *Salmonella*-CFA/I. In addition to the clinical scores, increased disease pathology of the CNS was evident from the enhanced demyelination and inflammatory cell infiltration relative to *Salmonella*-CFA/I-vaccinated mice. This protection was supported by increased production of IL-4, IL-10 and IL-13 and diminished production of IFN- γ by PLP_{139–151}- or CFA/I fimbriae-restimulated CD4⁺ T cells.¹¹² In contrast, CD4⁺ T cells isolated from unprotected or *Salmonella* vector-immunized mice¹¹² displayed elevated IFN- γ and minimal-to-no Th2-type cytokines. Thus, *Salmonella*-CFA/I was successful in reducing EAE severity in a bystander fashion when administered before disease induction.

Although these prophylactic results were clearly promising, often with human autoimmune diseases, it is unknown which patients will develop disease and at what time point. Therefore, to address the therapeutic potential of oral *Salmonella*-CFA/I, additional studies were conducted to determine whether it could impact ongoing disease.⁶⁸ Adapting the same PLP_{139–151} challenge model, mice were subjected to intervention with *Salmonella*-CFA/I, *Salmonella* vector or PBS 6 days post-EAE induction. As per the prophylactic studies, both the *Salmonella*-CFA/I and *Salmonella* vector were able to subdue disease to different degrees, but only *Salmonella*-CFA/I was able to inhibit CNS inflammation unlike the *Salmonella* vector-treated mice, which showed extensive neutrophil, macrophage and T-cell infiltration into the CNS.⁶⁸ Protection to EAE by *Salmonella*-CFA/I was attributed to the stimulation of anti-inflammatory cytokines, IL-4, IL-10 and IL-13 with concomitant reductions in IFN- γ and IL-17.⁶⁸

Because of the observed anti-inflammatory cytokines, we queried whether these were generated by T_{reg} cells, possibly induced by *Salmonella*-CFA/I. T_{reg} cells were originally delineated in thymectomized neonatal mice that showed increased manifestations of autoimmune disorders.^{113,114} As a result, T_{reg} cells have been shown to maintain peripheral tolerance and are responsible for protection against a variety of autoimmune diseases, including colitis,¹¹⁵ arthritis¹¹⁶ and EAE.^{81,83,105,117} T_{reg}-cell subsets are varied in their expression of phenotypic markers, but natural T_{reg} cells are identified by the expression of IL-2 receptor α -chain.¹¹³ These are also transcriptionally regulated by forkhead box transcription factor (Foxp3),^{117–120}

and when Foxp3 function is ablated, CD4⁺CD25⁺ T_{reg} cells are abrogated producing a wasting disease and inflammatory bowel disease in mice.^{118,119} Co-inhibitory molecules, activation-induced cytotoxic T lymphocyte associated protein-4 and glucocorticoid-induced TNF receptor may also be expressed and contribute in mediating T_{eff} cell suppression.^{121,122} In lieu of the phenotypic markers, earlier studies relied on functional characterization of these T_{reg} cells, and these were distinguished by the cell surface expression or secretion of TGF-β^{120,123,124} referred to as Th3 cells¹²⁵ or production of IL-10 referred to as T_{regulatory} 1 cells.^{126,127} TGF-β production by CD25⁺CD4⁺ T_{reg} cells has been shown to be responsible for recovery from EAE,^{68,89,95–97,123,128} and the presence of TGF-β has been shown to be necessary for conversion of CD4⁺CD25[−] T cells into CD4⁺CD25⁺ FoxP3⁺ T_{reg} cells.^{120,121,124,129} More recently, IL-35, a member of the IL-12 family and produced by T_{reg} cells, was found to have inhibitory activity capable of potentially suppressing arthritis,^{130,131} colitis^{132,133} and EAE.¹³⁴ IL-35 can mediate its effects via the stimulation of IL-10.^{130,131,135}

To examine the possible role for induction of T_{reg} cells by *Salmonella*-CFA/I, a kinetic analysis was done. It was discovered that both the *Salmonella* vector and *Salmonella*-CFA/I could stimulate the induction of CD25⁺CD4⁺ T cells, but the percentage of Foxp3⁺ T_{reg} cells was particularly augmented in mice treated with *Salmonella*-CFA/I.⁶⁸ To assess their relative contribution, *in vivo* CD25 neutralization was performed resulting in the diminution of *Salmonella*-CFA/I's protective response demonstrating the importance of these T_{reg} cells to abate autoimmune disease.⁶⁸ To assess their relative potency, adoptive transfer of T_{reg} cells from each of the treatment groups (*Salmonella*-CFA/I, *Salmonella* vector and naive) was tested.⁶⁸ T_{reg} cells derived from mice vaccinated with *Salmonella*-CFA/I showed the greatest potency in PLP_{139–151}-challenged mice exhibiting nearly complete protection. Conversely, although T_{reg} cells obtained from *Salmonella* vector-vaccinated mice did confer protection, some disease was still evident that was subsequently protracted. Recipients adoptively transferred with naive T_{reg} cells only showed a delayed onset of disease, and all mice developed EAE. Interestingly, partial protection could be achieved with CD25[−]CD4⁺ T cells adoptively transferred from *Salmonella*-CFA/I-vaccinated mice unlike those same cells from *Salmonella* vector-immunized mice that were unable to confer protection. The adoptively transferred T_{reg} cells from *Salmonella*-CFA/I-vaccinated mice produced both TGF-β and IL-10, but the majority of the IL-10 was derived from CD25[−]CD4⁺ T cells. Moreover, these CD25[−]CD4⁺ T cells produced IL-4 and IL-13, suggesting these are potentially Th2 cells. T_{reg} cells from *Salmonella* vector-vaccinated mice produced little TGF-β and IL-10, and their CD25[−]CD4⁺ T cells did not produce IL-4, IL-10 or IL-13, which accounts for the lesser potency in reducing EAE.⁶⁸ As TGF-β1^{−/−} mice succumb to death *in utero* and those surviving succumb early in life,¹³⁶ additional adoptive transfer studies were performed to test the dependence on TGF-β for protection to EAE.¹³⁷

Upon adoptive transfer of T_{reg} or CD25[−]CD4⁺ T cells from *Salmonella*-CFA/I-vaccinated mice, recipients induced with EAE were neutralized of their TGF-β using a monoclonal Ab (mAb), and this abrogated much of the protective effect by *Salmonella*-CFA/I's T_{reg} cells. Complete abrogation was observed upon treating recipients with *Salmonella*-CFA/I's CD25[−]CD4⁺ T cells.¹³⁷ As with this latter finding, TGF-β neutralization resulted in the complete loss of any partial protection conferred by *Salmonella* vector T_{reg} cells, and no protection was evident in recipients given CD25[−]CD4⁺ T cells from *Salmonella* vector-immunized mice.¹³⁷ TGF-β neutralization also impacted the expression of Foxp3 by the T_{reg} cells enabling IL-17 to be augmented, which would account for the loss of function and ultimately protection. Thus, TGF-β is an essential regulatory element induced by *Salmonella*-CFA/I therapeutic.

In an attempt to disrupt the protective capacity of CFA/I fimbriae, a mutant was developed to alter the cell surface expression of the fimbriae. This mutant bears only the *cfaAB* portion of the operon and lacks the genes for the outer membrane usher protein *cfaC*, and minor fimbrial subunit *cfaE*, and is referred to as *Salmonella*-CFA/I_(intracellular) (*Salmonella*-CFA/I_{IC}).¹³⁸ This restricted the major subunit CfaB primarily to the *Salmonella*'s periplasm. Upon oral vaccination, mice showed reduced fecal SIgA and reduced serum IgG Abs to the fimbriae but remained as effective in protecting mice against EAE as did adoptive transfer of their T_{reg} cells into EAE recipients.¹³⁸ What was particularly interesting of this finding was the difference in cytokine profiles from *Salmonella*-CFA/I-vaccinated mice. Examination of cytokine production by T_{reg} cells from *Salmonella*-CFA/I_{IC}-treated EAE mice revealed no change in TGF-β production but considerably augmented IFN-γ and IL-13 production relative to similarly treated mice with *Salmonella*-CFA/I.¹³⁸ The percentage of Foxp3⁺ T_{reg} cells was similar. To test the relevance of the observed IFN-γ and IL-13, *in vivo* Ab neutralization studies were conducted. Adoptive transfer of *Salmonella*-CFA/I_{IC}'s T_{reg} cells into EAE recipients neutralized of their IFN-γ using a mAb resulted in no differences in susceptibility to EAE, and both groups of mice given T_{reg} cells with normal rat IgG or rat anti-mouse IFN-γ mAb showed protection against disease. In contrast, EAE recipients neutralized of their IL-13 using a polyclonal anti-IL-13 Ab and adoptively transferred with *Salmonella*-CFA/I_{IC}'s T_{reg} cells lost their protection further showing the importance of IL-13 in defense against EAE.¹³⁸ This further implicated the ability of IL-13 to directly affect T_{reg} cell function or enhance the action of T_{reg} cells.¹³⁸ Thus, these collective studies show the effectiveness of *Salmonella*-CFA/I and *Salmonella*-CFA/I_{IC} as therapeutics to defend against EAE.

SALMONELLA-CFA/I AS AN ANTI-INFLAMMATORY VACCINE FOR AN INFLAMMATORY MODEL OF ARTHRITIS

Rheumatoid arthritis (RA) is an autoimmune disorder and chronic inflammatory disease of the joints impacting ~1% of

the population in North America and the United Kingdom,¹³⁹ with women being three times more likely to be afflicted than men.^{139,140} Although the etiology of this disease remains to be discerned, it is manifested as a chronic synovitis and progressive destruction of the joints, leukocyte infiltrates and cartilage destruction and bone erosion. This destruction is believed to be supported and perpetuated by proinflammatory cytokines. Past studies have indeed shown that proinflammatory cytokines are overexpressed in RA joints (reviewed in Feldmann *et al.*,¹⁴¹ Brennan *et al.*¹⁴² and Kannan *et al.*¹⁴³). To understand how such cytokines are regulated, a rodent model sharing many of the same features for human disease (reviewed in Kannan *et al.*¹⁴³) was developed. This autoimmune disease, collagen-induced arthritis (CIA), is induced by immunizing rodents, typically with heterologous (bovine or chick) type II collagen in combination with adjuvant, to elicit immune attack of the host's native collagen. Thus, components of both the innate and adaptive immune systems are involved. Emphasis on regulating proinflammatory cytokines, particularly TNF- α , is key to minimizing disease as it can be detected in joints of RA patients.^{144–146} Treatment with TNF- α antagonists decreases inflammation and attenuates the destruction of cartilage and bone.^{141,147–151} Such treatment is also believed to inhibit other inflammatory cytokines, including IL-1, IL-6, IL-8 and granulocyte macrophages colony-stimulating factor.^{141,142} Components of both the adaptive and innate immune systems contribute as a source for TNF- α and other proinflammatory cytokines.^{143,152,153} IFN- γ also contributes to disease, but it is phase dependent^{154–157}. Th17 cells, which can regulate both Th1 and Th2 cells,^{158,159} may be important for mediating RA disease as ICOS^{-/-} DBA/1 mice showed depressed IL-17 production, but not TNF- α or IFN- γ , and still conferred complete resistance to CIA.¹⁶⁰ IL-17 is also expressed by the human synovium and is particularly elevated in patients with RA^{161,162} and is IL-23 dependent.¹⁶³ *In vivo* neutralization of IL-17 results in significantly reduced CIA and could also lessen the progression of the established disease.¹⁶⁴

Given the results from the EAE studies, we queried whether *Salmonella*-CFA/I would be effective in treating CIA, a rodent model for RA.¹⁵³ DBA/1 mice are susceptible to chick or bovine collagen II (CII) challenge, and develop a progressive disease affecting multiple joints.^{153,165} As such, DBA/1 mice were orally dosed with *Salmonella*-CFA/I, *Salmonella* vector or PBS 7 days before CII challenge. Mice were followed for a course of 42 days, and *Salmonella*-CFA/I protected against CIA as evidenced by minimal clinical disease and significantly reduced incidence unlike mice treated with the *Salmonella* vector or PBS.¹⁶⁶ The observed reduction in disease was supported by the production of the cytokines IL-4, IL-10 and TGF- β by CD4⁺ T cells. In addition, mononuclear cells from *Salmonella*-CFA/I-treated animals had decreased levels of TNF- α , IL-1 β , IL-6 and IL-27.¹⁶⁶ To distill which CD4⁺ T cells were responsible for the regulatory and anti-inflammatory cytokines in DBA/1 mice void of disease, mice were orally vaccinated with *Salmonella*-CFA/I, and cytokine profiles from

CD25⁺CD4⁺ and CD25⁻CD4⁺ T cells were assessed. The CD25⁺CD4⁺ T cells produced significantly more IL-4, IL-10 and TGF- β than the CD25⁻ T cells. Interestingly, IFN- γ and IL-17 were significantly elevated relative to CD25⁻CD4⁺ T cells but less than CD25⁺CD4⁺ T cells from *Salmonella* vector-immunized mice.¹⁶⁶ As the CD25⁺CD4⁺ T cells from similarly vaccinated mice were highly protective against EAE,⁶⁸ adoptive transfer studies were performed to measure the potency of these T_{reg} cells in conferring protection against CIA. Surprisingly, neither the individual CD25⁺CD4⁺ nor CD25⁻CD4⁺ T-cell subset was adequate in treating CIA relative to whole CD4⁺ T-cell isolates with respect to disease onset and mice with reduced clinical disease.¹⁶⁶ Thus, it appeared that both CD25⁺CD4⁺ and CD25⁻CD4⁺ T cells were required for protection against CIA. Subsequently, adoptive transfer studies using total CD4⁺ T cells in combination with anti-IL-4 mAb or anti-TGF- β mAb were performed and revealed that inhibition of either cytokine resulted in disease and loss of the protective response.¹⁶⁶ Collectively, these data showed that *Salmonella*-CFA/I could treat CIA via the induction of diverse populations of T_{reg} cells.

As neither CD25⁺ nor CD25⁻CD4⁺ T cells could completely protect following adoptive transfer into DBA/1 mice, our data suggested that perhaps a different T_{reg}-cell subset was being induced. To investigate such a possibility, the *Salmonella*-CFA/I-induced CD4⁺ T cells were screened for expression of alternative T_{reg}-cell markers other than CD25. One such alternative is CD39. These CD39⁺ T_{reg} cells are also able to suppress Th17 cells, and their absence has been linked to MS.¹⁶⁷ Specifically, CD39 is an ectonucleoside triphosphate diphosphohydrolase, and it is expressed on the cell surface of Foxp3⁺ T_{reg} cells, dampening proinflammatory cells by ultimately converting proinflammatory extracellular ATP to anti-inflammatory adenosine.^{166,168,169} Subsequent evaluation of CD39⁺ expression was conducted in C57BL/6 male mice, which also are susceptible to arthritis.¹⁷⁰ Thus, CIA mice were orally dosed with *Salmonella*-CFA/I, and it was revealed that only half of the CD39⁺CD4⁺ T cells was Foxp3⁺CD25⁺CD4⁺, although CD39 was also expressed on the CD25⁻CD4⁺ T cells.¹⁷⁰ To determine whether these CD39⁺ T cells were protective against disease, C57BL/6 mice were adoptively transferred with CD39⁺CD4⁺ or CD39⁻CD4⁺ T cells from *Salmonella*-CFA/I- or *Salmonella* vector-dosed mice into recipients challenged 14 days earlier with CIA. Only mice receiving CD39⁺CD4⁺ T cells from *Salmonella*-CFA/I-dosed mice were protected against CIA, indicating that again the CFA/I fimbriae are essential in stimulating this therapeutic subset of T_{reg} cells. As these CD39⁺CD4⁺ T cells were composed of both Foxp3⁺ and Foxp3⁻ cells, further analysis was performed to determine whether Foxp3⁺CD39⁺CD4⁺ T cells were protective against CIA. Notably, CIA recipients given either Foxp3⁺CD39⁺CD4⁺ or Foxp3⁻CD39⁺CD4⁺ T cells protected equally to disease but not as effectively as total CD39⁺CD4⁺ T cells.¹⁷⁰ These subsets were further teased to discern how they differ. Cytokine analysis revealed that IL-10 and TGF- β segregated

with Foxp3⁺CD39⁺CD4⁺ and Foxp3⁻CD39⁺CD4⁺ T cells, respectively.¹⁷⁰ Neutralization of TGF- β reduced the percentage of CD39 expression, implicating the importance of TGF- β for induction of CD39.¹⁷⁰

CONCLUSIONS

Outside of our studies, there have only been a few reports where bacterial infections have been used to subdue EAE,^{171,172} and even less for CIA. However, it is important to emphasize that the induction of T_{reg} cells is mediated not so much by the attenuated *Salmonella* vaccine strain as these cells are induced by the fimbriae or the combination of bacteria with fimbriae. Current studies are addressing these possibilities. Nonetheless, our studies demonstrate the feasibility of a simple oral treatment with *Salmonella*-CFA/I to render protection against EAE and CIA without having previous knowledge of the auto-Ag. Consequently, in a bystander fashion, both fimbriae- and PLP₁₃₉₋₁₅₁-specific, TGF- β -producing, FoxP3⁺CD25⁺CD4⁺ T cells were induced for EAE and CII-specific, IL-10-producing FoxP3⁺CD39⁺ and TGF- β -producing FoxP3⁻CD39⁺CD4⁺ T cells. For the protective T_{reg} cells in CIA, co-expression of CD25 did not specifically segregate with either subset.¹⁷⁰ The advantage of using *Salmonella*-CFA/I is that intervention of autoimmunity can be achieved upon vaccination with an innocuous Ag, and in this case the side-effect would be the additional protection against the diarrheal diseases, ETEC and salmonellosis. Moreover, this therapeutic can be administered orally enabling patient compliance. Additionally, this approach does not polyclonally activate T_{reg} cells, which have been shown to have a cataclysmic outcome.¹⁷³

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by the Public Health Service Grants AI-41123 and P01 AT-004986-05.

- Svennerholm AM, Lundgren A. Recent progress toward an enterotoxigenic *Escherichia coli* vaccine. *Expert Rev Vaccines* 2012; **11**: 495–507.
- Qadri F, Svennerholm AM, Faruque AS, Sack RB. Enterotoxigenic *Escherichia coli* in developing countries: epidemiology, microbiology, clinical features, treatment, and prevention. *Clin Microbiol Rev* 2005; **18**: 465–483.
- Hill DR, Beeching NJ. Travelers' diarrhea. *Curr Opin Infect Dis* 2010; **23**: 481–487.
- Field M, Graf LH, Laird WJ, Smith PL. Heat-stable enterotoxin of *Escherichia coli*: in vitro effects on guanylate cyclase activity, cyclic GMP concentration and ion transport in the small intestine. *Proc Natl Acad Sci USA* 1978; **75**: 2800–2804.
- Clements JD, Finkelstein RA. Demonstration of shared and unique immunologic determinants in enterotoxins from *V. cholerae* and *E. coli*. *Infect Immun* 1978; **22**: 709–713.
- Spangler BD. Structure and function of cholera toxin and the related *Escherichia coli* heat-labile enterotoxin. *Microbiol Rev* 1992; **56**: 622–647.
- Evans DJ Jr, Evans DG. Colonization factor antigens of human pathogens. *Curr Top Microbiol Immunol* 1990; **151**: 129–145.
- de Graaf FK, Gastra W. Fimbriae of enterotoxigenic *Escherichia coli*. In: Klemm P (ed) *Fimbriae: Adhesins, Biogenics, Genetics, and Vaccines*. CRC Press: Boca Raton, FL, USA, 1994. pp 53–83.
- Tchesnokova V, McVeigh AL, Kidd B, Yakovenko O, Thomas WE, Sokurenko EV *et al*. Shear-enhanced binding of intestinal colonization factor antigen I of enterotoxigenic *Escherichia coli*. *Mol Microbiol* 2010; **76**: 489–502.
- Li YF, Poole S, Rasulova F, McVeigh AL, Savarino SJ, Xia D. A receptor-binding site as revealed by the crystal structure of CfaE, the colonization factor antigen I fimbrial adhesin of enterotoxigenic *Escherichia coli*. *J Biol Chem* 2007; **282**: 23970–23980.
- Jansson L, Tobias J, Lebens M, Svennerholm AM, Teneberg S. The major subunit, CfaB, of colonization factor antigen I from enterotoxigenic *Escherichia coli* is a glycosphingolipid binding protein. *Infect Immun* 2006; **74**: 3488–3497.
- Pascual DW, Ochoa-Repáraz J, Rynda A, Yang X. Tolerance in the absence of auto-antigen. *Endocr Metab Immune Disord Drug Targets* 2007; **7**: 203–210.
- Black RE, Merson MH, Rowe B, Taylor PR, Abdul Alim AR, Gross RJ *et al*. Enterotoxigenic *Escherichia coli* diarrhoea: acquired immunity and transmission in an endemic area. *Bull World Health Organ* 1981; **59**: 263–268.
- Black RE, Brown KH, Becker S, Alim AR, Huq I. Longitudinal studies of infectious diseases and physical growth of children in rural Bangladesh. II. Incidence of diarrhea and association with known pathogens. *Am J Epidemiol* 1982; **115**: 315–324.
- Cravioto A, Reyes RE, Ortega R, Fernández G, Hernández R, López D. Prospective study of diarrhoeal disease in a cohort of rural Mexican children: incidence and isolated pathogens during the first two years of life. *Epidemiol Infect* 1988; **101**: 123–134.
- Levine MM. *Escherichia coli* that cause diarrhea: enterotoxigenic, enteropathogenic, enteroinvasive, enterohemorrhagic, and enteroadherent. *J Infect Dis* 1987; **155**: 377–389.
- Levine MM, Girón JA, Noriega FR. Fimbrial vaccines. In: Klemm P (ed) *Fimbriae: Adhesins, Biogenics, Genetics, and Vaccines*. CRC Press: Boca Raton, FL, USA, 1994. pp 255–270.
- Anantha RP, McVeigh AL, Lee LH, Agnew MK, Cassels FJ, Scott DA *et al*. Evolutionary and functional relationships of colonization factor antigen I and other class 5 adhesive fimbriae of enterotoxigenic *Escherichia coli*. *Infect Immun* 2004; **72**: 7190–7201.
- Li YF, Poole S, Nishio K, Jang K, Rasulova F, McVeigh A *et al*. Structure of CFA/I fimbriae from enterotoxigenic *Escherichia coli*. *Proc Natl Acad Sci USA* 2009; **106**: 10793–10798.
- Li YF, Poole S, Rasulova F, Esser L, Savarino SJ, Xia D. Crystallization and preliminary X-ray diffraction analysis of CfaE, the adhesive subunit of the CFA/I fimbriae from human enterotoxigenic *Escherichia coli*. *Acta Crystallogr Sect F Struct Biol Cryst Commun* 2006; **62**(Pt 2), 121–124.
- Edelman R, Russell RG, Losonsky G, Tall BD, Tacket CO, Levine MM *et al*. Immunization of rabbits with enterotoxigenic *E. coli* colonization factor antigen (CFA/I) encapsulated in biodegradable microspheres of poly (lactide-co-glycolide). *Vaccine* 1993; **11**: 155–158.
- Evans DG, Graham DY, Evans DJ Jr. Administration of purified colonization factor antigens (CFA/I, CFA/II) of enterotoxigenic *Escherichia coli* to volunteers. Response to challenge with virulent enterotoxigenic *Escherichia coli*. *Gastroenterology* 1984; **87**: 934–940.
- Levine M, Morris JG, Losonsky G, Boedeker E, Rowe B. Fimbriae (pili) adhesins as vaccine. In: Lark D L, Normark S, Uhlin B-E, Wolf-Watz H (eds) *Protein-Carbohydrate Interactions in Biological Systems: Molecular Biology of Microbial Pathogenicity*. Academic Press: London, UK, 1986. pp 143–145.
- Schmidt M, Kelly EP, Tseng LY, Boedeker EC. Towards an oral *E. coli* pilus vaccine for travelers diarrhea: susceptibility to proteolytic digestion. *Gastroenterology* 1985; **82**: A1575.
- Reid RH, Boedeker EC, McQueen CE, Davis D, Tseng LY, Kodak J *et al*. Preclinical evaluation of microencapsulated CFA/II oral vaccine against enterotoxigenic *E. coli*. *Vaccine* 1993; **11**: 159–167.
- Sánchez J, Holmgren J. Virulence factors, pathogenesis and vaccine protection in cholera and ETEC diarrhea. *Curr Opin Immunol* 2005; **17**: 388–398.
- Boedeker EC. Vaccines for enterotoxigenic *Escherichia coli*: current status. *Curr Opin Gastroenterol* 2005; **21**: 15–19.
- Galen JE, Pasetti MF, Tennant S, Ruiz-Olvera P, Szein MB, Levine MM. *Salmonella enterica* serovar Typhi live vector vaccines finally come of age. *Immunol Cell Biol* 2009; **87**: 400–412.

- 29 McGregor AC, Waddington CS, Pollard AJ. Prospects for prevention of *Salmonella* infection in children through vaccination. *Curr Opin Infect Dis* 2013; **26**: 254–262.
- 30 Desin TS, Köster W, Potter AA. *Salmonella* vaccines in poultry: past, present and future. *Expert Rev Vaccines* 2013; **12**: 87–96.
- 31 Pasetti MF, Levine MM, Sztein MB. Animal models paving the way for clinical trials of attenuated *Salmonella enterica* serovar Typhi live oral vaccines and live vectors. *Vaccine* 2003; **21**: 401–418.
- 32 Curtiss R 3rd, Xin W, Li Y, Kong W, Wanda SY, Gunn B *et al*. New technologies in using recombinant attenuated *Salmonella* vaccine vectors. *Crit Rev Immunol* 2010; **30**: 255–270.
- 33 Mestecky J, Nguyen H, Czerkinsky C, Kiyono H. Oral immunization: an update. *Curr Opin Gastroenterol* 2008; **24**: 713–719.
- 34 Curtiss R 3rd. Bacterial infectious disease control by vaccine development. *J Clin Invest* 2002; **110**: 1061–1066.
- 35 Wu S, Pascual DW, VanCott JL, McGhee JR, Maneval DR Jr, Levine MM *et al*. Immune responses to novel *Escherichia coli* and *Salmonella typhimurium* vectors that express colonization factor antigen I (CFA/I) of enterotoxigenic *E. coli* in the absence of the CFA/I positive regulator cfaR. *Infect Immun* 1995; **63**: 4933–4938.
- 36 Yang X, Suo Z, Thornburg T, Holderness K, Walters N, Kellerman L *et al*. Expression of *Escherichia coli* virulence usher protein attenuates wild-type *Salmonella*. *Virulence* 2012; **3**: 29–42.
- 37 Cao L, Suo Z, Lim T, Jun S, Deliorman M, Riccardi C *et al*. Role of overexpressed CFA/I fimbriae in bacterial swimming. *Phys Biol* 2012; **9**: 036005.
- 38 Pascual DW, Hone DM, Hall S, van Ginkel FW, Yamamoto M, Walters N *et al*. Expression of recombinant enterotoxigenic *Escherichia coli* colonization factor antigen I by *Salmonella typhimurium* elicits a biphasic T helper cell response. *Infect Immun* 1999; **67**: 6249–6256.
- 39 Nauciel C, Espinasse-Maes F. Role of gamma interferon and tumor necrosis factor alpha in resistance to *Salmonella typhimurium* infection. *Infect Immun* 1992; **60**: 450–454.
- 40 Killar LM, Eisenstein TK. Delayed-type hypersensitivity and immunity to *Salmonella typhimurium*. *Infect Immun* 1986; **52**: 504–508.
- 41 Pope M, Kotlarski I. Detection of *Salmonella*-specific L3T4+ and LYT-2+ T cells which can proliferate in vitro and mediate delayed-type hypersensitivity reactivity. *Immunol* 1994; **81**: 183–191.
- 42 Mastroeni P, Villareal-Ramos B, Hormaeche CE. Role of T cells, TNF alpha and IFN gamma in recall of immunity to oral challenge with virulent salmonellae in mice vaccinated with live attenuated aro- *Salmonella* vaccines. *Microb Pathog* 1992; **13**: 477–491.
- 43 VanCott JL, Staats HF, Pascual DW, Roberts M, Chatfield SN, Yamamoto M *et al*. Regulation of mucosal and systemic antibody responses by T helper cell subsets, macrophages, and derived cytokines following oral immunization with live recombinant *Salmonella*. *J Immunol* 1996; **156**: 1504–1514.
- 44 Ravindran R, Foley J, Stoklasek T, Glimcher LH, McSorley SJ. Expression of T-bet by CD4 T cells is essential for resistance to *Salmonella* infection. *J Immunol* 2005; **175**: 4603–4610.
- 45 Yang DM, Fairweather N, Button LL, McMaster WR, Kahl LP, Liew FY. Oral *Salmonella typhimurium* (AroA-) vaccine expressing a major leishmanial surface protein (gp63) preferentially induces T helper 1 cells and protective immunity against leishmaniasis. *J Immunol* 1990; **145**: 2281–2285.
- 46 Fouts TR, Tuskan RG, Chada S, Hone DM, Lewis GK. Construction and immunogenicity of *Salmonella typhimurium* vaccine vectors that express HIV-1 gp120. *Vaccine* 1995; **13**: 1697–1705.
- 47 Xu D, McSorley SJ, Chatfield SN, Dougan G, Liew FY. Protection against Leishmania major infection in genetically susceptible BALB/c mice by gp63 delivered orally in attenuated *Salmonella typhimurium* (AroA- AroD-). *Immunol* 1995; **85**: 1–7.
- 48 Hess J, Gentschev I, Miko D, Welzel M, Ladel C, Goebel W *et al*. Superior efficacy of secreted over somatic antigen display in recombinant *Salmonella* vaccine induced protection against listeriosis. *Proc Natl Acad Sci USA* 1996; **93**: 1458–1463.
- 49 Walters N, Trunkle T, Sura M, Pascual DW. Enhanced immunoglobulin A response and protection against *Salmonella enterica* serovar Typhimurium in the absence of the substance P receptor. *Infect Immun* 2005; **73**: 317–324.
- 50 Tite JP, Dougan G, Chatfield SN. The involvement of tumor necrosis factor in immunity to *Salmonella* infection. *J Immunol* 1991; **147**: 3161–3164.
- 51 Pascual DW, Trunkle T, Sura J. Fimbriated *Salmonella enterica* serovar Typhimurium abates initial inflammatory responses by macrophages. *Infect Immun* 2002; **70**: 4273–4281.
- 52 Miller SI, Ernst RK, Bader MW. LPS, TLR4 and infectious disease diversity. *Nat Rev Microbiol* 2005; **3**: 36–46.
- 53 Swanborg RH. Experimental autoimmune encephalomyelitis in rodents as a model for human demyelinating disease. *Clin Immunol Immunopathol* 1995; **77**: 4–13.
- 54 Smith ME, Eller NL, McFarland HF, Racke MK, Raine CS. Age dependence of clinical and pathological manifestations of autoimmune demyelination. Implications for multiple sclerosis. *Am J Pathol* 1999; **155**: 1147–1161.
- 55 Hafler DA. Multiple sclerosis. *J Clin Invest* 2004; **113**: 788–794.
- 56 Batoulis H, Addicks K, Kuerten S. Emerging concepts in autoimmune encephalomyelitis beyond the CD4/TH1 paradigm. *Ann Anat* 2010; **192**: 179–193.
- 57 Prat E, Martin R. The immunopathogenesis of multiple sclerosis. *J Rehab Res Dev* 2002; **39**: 187–200.
- 58 Steinman L, Martin R, Bernard C, Conlon P, Oksenberg JR. Multiple sclerosis: deeper understanding of its pathogenesis reveals new targets for therapy. *Annu Rev Neurosci* 2002; **25**: 491–505.
- 59 Inglesse M. Multiple sclerosis: new insights and trends. *AJNR Am J Neuroradiol* 2006; **27**: 954–957.
- 60 Fujinami RS. Can virus infections trigger autoimmune disease? *J Autoimmun* 2001; **16**: 229–234.
- 61 von Büdingen HC, Tanuma N, Villoslada P, Ouallet JC, Hauser SL, Genain CP. Immune responses against the myelin/oligodendrocyte glycoprotein in experimental autoimmune demyelination. *J Clin Immunol* 2001; **21**: 155–170.
- 62 Miller SD, Katz-Levy Y, Neville KL, Vanderlugt CL. Virus-induced autoimmunity: epitope spreading to myelin autoepitopes in Theiler's virus infection of the central nervous system. *Adv Virus Res* 2001; **56**: 199–217.
- 63 Swanborg RH. Experimental autoimmune encephalomyelitis in the rat: lessons in T-cell immunology and autoreactivity. *Immunol Rev* 2001; **184**: 129–135.
- 64 Kuchroo VK, Anderson AC, Waldner H, Munder M, Bettelli E, Nicholson LB. T cell response in experimental autoimmune encephalomyelitis (EAE): role of self and cross-reactive antigens in shaping, tuning, and regulating the autopathogenic T cell repertoire. *Annu Rev Immunol* 2002; **20**: 101–123.
- 65 Martin Mdel P, Monson NL. Potential role of humoral immunity in the pathogenesis of multiple sclerosis (MS) and experimental autoimmune encephalomyelitis (EAE). *Front Biosci* 2007; **12**: 2735–2749.
- 66 St Louis J, Uniyal S, Xu L, Chan E, Singh B, Chan BM *et al*. Tolerance induction by acylated peptides: suppression of EAE in the mouse with palmitoylated PLP peptides. *J Neuroimmunol* 2001; **115**: 79–90.
- 67 Hall SW, Cooke A. Autoimmunity and inflammation: murine models and translational studies. *Mamm Genome* 2011; **22**: 377–389.
- 68 Ochoa-Repáraz J, Riccardi C, Rynda A, Jun S, Callis G, Pascual DW. Regulatory T cell vaccination without autoantigen protects against experimental autoimmune encephalomyelitis. *J Immunol* 2007; **178**: 1791–1799.
- 69 Stromnes IM, Goverman JM. Active induction of experimental allergic encephalomyelitis. *Nat Protoc* 2006; **1**: 1810–1819.
- 70 McRae BL, Vanderlugt CL, Dal Canto MC, Miller SD. Functional evidence for epitope spreading in the relapsing pathology of experimental autoimmune encephalomyelitis. *J Exp Med* 1995; **182**: 75–85.
- 71 Tuohy VK, Yu M, Yin L, Mathisen PM, Johnson JM, Kawczak JA. Modulation of the IL-10/IL-12 cytokine circuit by interferon-beta inhibits the development of epitope spreading and disease progression in murine autoimmune encephalomyelitis. *J Neuroimmunol* 2000; **111**: 55–63.
- 72 Yin L, Yu M, Edling AE, Kawczak JA, Mathisen PM, Nanavati T *et al*. Pre-emptive targeting of the epitope spreading cascade with genetically modified regulatory T cells during autoimmune demyelinating disease. *J Immunol* 2001; **167**: 6105–6112.
- 73 Cua DJ, Sherlock J, Chen Y, Murphy CA, Joyce B, Seymour B *et al*. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* 2003; **421**: 744–748.
- 74 Becher B, Durell BG, Noelle RJ. Experimental autoimmune encephalitis and inflammation in the absence of interleukin-12. *J Clin Invest* 2002; **110**: 493–497.

- 75 Zhang GX, Gran B, Yu S, Li J, Siglienti I, Chen X *et al*. Induction of experimental autoimmune encephalomyelitis in IL-12 receptor- β -deficient mice: IL-12 responsiveness is not required in the pathogenesis of inflammatory demyelination in the central nervous system. *J Immunol* 2003; **170**: 2153–2160.
- 76 Park H, Li Z, Yang XO, Chang SH, Nurieva R, Wang YH *et al*. A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17. *Nat Immunol* 2005; **6**: 1133–1141.
- 77 Hofstetter HH, Ibrahim SM, Koczan D, Kruse N, Weishaupt A, Toyka KV *et al*. Therapeutic efficacy of IL-17 neutralization in murine experimental autoimmune encephalomyelitis. *Cell Immunol* 2005; **237**: 123–130.
- 78 Komiyama Y, Nakae S, Matsuki T, Nambu A, Ishigame H, Kakuta S *et al*. IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *J Immunol* 2006; **177**: 566–573.
- 79 Sutton C, Brereton C, Keogh B, Mills KH, Lavelle EC. A crucial role for interleukin (IL)-1 in the induction of IL-17-producing T cells that mediate autoimmune encephalomyelitis. *J Exp Med* 2006; **203**: 1685–1691.
- 80 Inobe J, Slavin AJ, Komagata Y, Chen Y, Liu L, Weiner HL. IL-4 is a differentiation factor for transforming growth factor- β secreting Th3 cells and oral administration of IL-4 enhances oral tolerance in experimental allergic encephalomyelitis. *Eur J Immunol* 1998; **28**: 2780–2790.
- 81 Zhang X, Koldicz DN, Izikson L, Reddy J, Nazareno RF, Sakaguchi S *et al*. IL-10 is involved in the suppressing growth factor- β secreting Th3 cells and oral administration of IL-4 enhances oral tolerance in experimental allergic encephalomyelitis. *Eur J Immunol* 1998; **28**: 2780–2790.
- 82 Sun JB, Xiao BG, Lindblad M, Li BL, Link H, Czerkinsky C *et al*. Oral administration of cholera toxin B subunit conjugated to myelin basic protein protects against experimental autoimmune encephalomyelitis by inducing transforming growth factor- β -secreting cells and suppressing chemokine expression. *Int Immunol* 2000; **12**: 1449–1457.
- 83 Kohm AP, Carpentier PA, Anger HA, Miller SD. Cutting edge: CD4 + CD25 + regulatory T cells suppress antigen-specific autoreactive immune responses and central nervous system inflammation during active experimental autoimmune encephalomyelitis. *J Immunol* 2002; **169**: 4712–4716.
- 84 Tran EH, Prince EN, Owens T. IFN- γ shapes immune invasion of the central nervous system via regulation of chemokines. *J Immunol* 2000; **164**: 2759–2768.
- 85 Ito A, Bebo BF Jr, Matejuk A, Zamora A, Silverman M, Fyfe-Johnson A *et al*. Estrogen treatment down-regulates TNF- α production and reduces the severity of experimental autoimmune encephalomyelitis in cytokine knockout mice. *J Immunol* 2001; **167**: 542–552.
- 86 Willenborg DO, Fordham SA, Staykova MA, Ramshaw IA, Cowden WB. IFN- γ is critical to the control of murine autoimmune encephalomyelitis and regulates both in the periphery and in the target tissue: a possible role for nitric oxide. *J Immunol* 1999; **163**: 5278–5286.
- 87 Chu CQ, Wittmer S, Dalton DK. Failure to suppress the expansion of the activated CD4 T cell population in interferon γ -deficient mice leads to exacerbation of experimental autoimmune encephalomyelitis. *J Exp Med* 2000; **192**: 123–128.
- 88 Körner H, Lemckert FA, Chaudhri G, Etteldorf S, Sedgwick JD. Tumor necrosis factor blockade in actively induced experimental autoimmune encephalomyelitis prevents clinical disease despite activated T cell infiltration to the central nervous system. *Eur J Immunol* 1997; **27**: 1973–1981.
- 89 Khoury SJ, Hancock WW, Weiner HL. Oral tolerance to myelin basic protein and natural recovery from experimental autoimmune encephalomyelitis are associated with downregulation of inflammatory cytokines and differential upregulation of transforming growth factor β , interleukin 4, and prostaglandin E expression in the brain. *J Exp Med* 1992; **176**: 1355–1364.
- 90 Falcone M, Rajan AJ, Bloom BR, Brosnan CF. A critical role for IL-4 in regulating disease severity in experimental allergic encephalomyelitis as demonstrated in IL-4-deficient C57BL/6 mice and BALB/c mice. *J Immunol* 1998; **160**: 4822–4830.
- 91 Mendel I, Natarajan K, Ben-Nun A, Shevach EM. A novel protective model against experimental allergic encephalomyelitis in mice expressing a transgenic TCR-specific for myelin oligodendrocyte glycoprotein. *J Neuroimmunol* 2004; **149**: 10–21.
- 92 Cua DJ, Groux H, Hinton DR, Stohlman SA, Coffman RL. Transgenic interleukin 10 prevents induction of experimental autoimmune encephalomyelitis. *J Exp Med* 1999; **189**: 1005–1010.
- 93 Legge KL, Min B, Bell JJ, Caprio JC, Li L, Gregg RK *et al*. Coupling of peripheral tolerance to endogenous interleukin 10 promotes effective modulation of myelin-activated T cells and ameliorates experimental allergic encephalomyelitis. *J Exp Med* 2000; **191**: 2039–2052.
- 94 Slavin AJ, Maron R, Weiner HL. Mucosal administration of IL-10 enhances oral tolerance in autoimmune encephalomyelitis and diabetes. *Int Immunol* 2001; **13**: 825–833.
- 95 Karpus WJ, Swanborg RH. CD4 + suppressor cells inhibit the function of effector cells of experimental autoimmune encephalomyelitis through a mechanism involving transforming growth factor- β . *J Immunol* 1991; **146**: 1163–1168.
- 96 Racke MK, Dhib-Jalbut S, Cannella B, Albert PS, Raine CS, McFarlin DE. Prevention and treatment of chronic relapsing experimental allergic encephalomyelitis by transforming growth factor-beta 1. *J Immunol* 1991; **146**: 3012–3017.
- 97 Miller A, Lider O, Roberts AB, Sporn MB, Weiner HL. Suppressor T cells generated by oral tolerization to myelin basic protein suppress both *in vitro* and *in vivo* immune responses by the release of transforming growth factor beta after antigen-specific triggering. *Proc Natl Acad Sci USA* 1992; **89**: 421–425.
- 98 Pot C, Apetoh L, Awasthi A, Kuchroo VK. Induction of regulatory Tr1 cells and inhibition of TH17 cells by IL-27. *Semin Immunol* 2011; **23**: 438–445.
- 99 Fitzgerald DC, Ciric B, Touil T, Harle H, Grammatikopoulou J, Das Sarma J *et al*. Suppressive effect of IL-27 on encephalitogenic Th17 cells and the effector phase of experimental autoimmune encephalomyelitis. *J Immunol* 2007; **179**: 3268–3275.
- 100 Fitzgerald DC, Zhang GX, El-Behi M, Fonseca-Kelly Z, Li H, Yu S *et al*. Suppression of autoimmune inflammation of the central nervous system by interleukin 10 secreted by interleukin 27-stimulated T cells. *Nat Immunol* 2007; **8**: 1372–1379.
- 101 Bettini M, Vignali DA. Regulatory T cells and inhibitory cytokines in autoimmunity. *Curr Opin Immunol* 2009; **21**: 612–618.
- 102 Paust S, Cantor H. Regulatory T cells and autoimmune disease. *Immunol Rev* 2005; **204**: 195–207.
- 103 Faria AMC, Weiner HL. Oral tolerance. *Immunol Rev* 2005; **206**: 232–259.
- 104 Mueller DL. Mechanisms maintaining peripheral tolerance. *Nat Immunol* 2010; **11**: 21–27.
- 105 Bilate AM, Lafaille JJ. Induced CD4 + Foxp3 + regulatory T cells in immune tolerance. *Annu Rev Immunol* 2012; **30**: 733–758.
- 106 Canonica GW, Passalacqua G. Sublingual immunotherapy in the treatment of adult allergic rhinitis patients. *Allergy* 2006; **61**(Suppl 81), 20–23.
- 107 Dahl R, Kapp A, Colombo G, de Monchy JG, Rak S, Emminger W *et al*. Efficacy and safety of sublingual immunotherapy with grass allergen tablets for seasonal allergic rhinoconjunctivitis. *J Allergy Clin Immunol* 2006; **118**: 434–440.
- 108 Didier A, Malling HJ, Worm M, Horak F, Jäger S, Montagut A *et al*. Optimal dose, efficacy, and safety of once-daily sublingual immunotherapy with a 5-grass pollen tablet for seasonal allergic rhinitis. *J Allergy Clin Immunol* 2007; **120**: 1338–1345.
- 109 Faria A, Weiner HL. Oral tolerance: therapeutic implications for autoimmune disease. *Clin Dev Immunol* 2006; **13**: 143–157.
- 110 Skyler JS, Krischer JP, Wolfsdorf J, Cowie C, Palmer JP, Greenbaum C *et al*. Effects of oral insulin in relatives of patients with type 1 diabetes: the Diabetes Prevention Trial—Type 1. *Diabetes Care* 2005; **28**: 1068–1076.
- 111 Pascual DW, White MD, Larson T, Walters N. Impaired mucosal immunity in L-selectin-deficient mice orally immunized with a *Salmonella* vaccine vector. *J Immunol* 2001; **167**: 407–415.
- 112 Jun S, Gilmore W, Callis G, Rynda A, Haddad A, Pascual DW. A live diarrheal vaccine imprints a Th2 cell bias and acts as an anti-inflammatory vaccine. *J Immunol* 2005; **175**: 6733–6740.
- 113 Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor α -chains (CD25): breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* 1995; **155**: 1151–1164.
- 114 Hori S, Takahashi T, Sakaguchi S. Control of autoimmunity by naturally arising regulatory CD4 + T cells. *Adv Immunol* 2003; **81**: 331–371.
- 115 Huber S, Schramm C, Lehr HA, Mann A, Schmitt S, Becker C *et al*. Cutting edge: TGF- β signaling is required for the *in vivo* expansion and immunosuppressive capacity of regulatory CD4 + CD25 + T cells. *J Immunol* 2004; **173**: 6526–6531.

- 116 Morgan ME, Flierman R, van Duivenvoorde LM, Witteveen HJ, van Ewijk W, van Laar JM *et al*. Effective treatment of collagen-induced arthritis by adoptive transfer of CD25+ regulatory T cells. *Arthritis Rheum* 2005; **52**: 2212–2221.
- 117 Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science* 2003; **99**: 1057–1061.
- 118 Fontenot JD, Gavin MA, Rudensky AY. FoxP3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol* 2003; **4**: 330–336.
- 119 Vieira PL, Christensen JR, Minaee S, O'Neill EJ, Barrat FJ, Boonstra A *et al*. IL-10-secreting regulatory T cells do not express FoxP3 but have comparable regulatory function to naturally occurring CD4+CD25+ regulatory T cells. *J Immunol* 2004; **172**: 5986–5993.
- 120 Chen W, Jin W, Hardegen N, Lei KJ, Li L, Marinos N *et al*. Conversion of peripheral CD4+CD25- naive T cells into CD4+CD25+ regulatory T cells by TGF- β induction of transcription factor FoxP3. *J Exp Med* 2003; **198**: 1875–1886.
- 121 Oida T, Xu L, Weiner HL, Kitani A, Strober W. TGF- β -mediated suppression by CD4+CD25+ T cells is facilitated by CTLA-4 signaling. *J Immunol* 2006; **177**: 2331–2339.
- 122 Shimizu J. Stimulation of CD4+CD25+ regulatory T cells through GITR breaks immunological self-tolerance. *Nat Immunol* 2002; **3**: 135–142.
- 123 Faria AM, Maron R, Ficker SM, Slavin AJ, Spahn T, Weiner HL. Oral tolerance induced by continuous feeding: enhanced up-regulation of transforming growth factor- β /interleukin-10 and suppression of experimental autoimmune encephalomyelitis. *J Autoimmun* 2003; **20**: 135–145.
- 124 Zheng S, Wang J, Horwitz DA. Cutting edge: FoxP3+CD4+CD25+ regulatory T cells induced by IL-2 and TGF- β are resistant to Th17 conversion by IL-6. *J Immunol* 2008; **180**: 7112–7116.
- 125 Weiner HL. Oral tolerance: immune mechanisms and the generation of Th3-type TGF- β -secreting regulatory cells. *Microbes Infect* 2001; **3**: 947–954.
- 126 Mowat AM. Anatomical basis of tolerance and immunity to intestinal antigens. *Nat Rev Immunol* 2003; **3**: 331–341.
- 127 Wildbaum G, Netzer N, Karin N. Tr1-dependent active tolerance blunts the pathogenic effects of determinant spreading. *J Clin Invest* 2002; **110**: 701–710.
- 128 Santos LM, al-Sabbagh A, Londono A, Weiner HL. Oral tolerance to myelin basic protein induces regulatory TGF- β -secreting T cells in Peyer's patches of SJL mice. *Cell Immunol* 1994; **157**: 439–447.
- 129 Chen ML, Yan BS, Bando Y, Kuchroo VK, Weiner HL. Latency-associated peptide identifies a novel CD4+CD25+ regulatory T cell subset with TGF- β -mediated function and enhances suppression of experimental autoimmune encephalomyelitis. *J Immunol* 2008; **180**: 7327–7337.
- 130 Niedbala W, Wei XQ, Cai B, Hueber AJ, Leung BP, McInnes IB *et al*. IL-35 is a novel cytokine with therapeutic effects against collagen-induced arthritis through the expansion of regulatory T cells and suppression of Th17 cells. *Eur J Immunol* 2007; **37**: 3021–3029.
- 131 Kochetkova I, Golden S, Holderness K, Callis G, Pascual DW. IL-35 stimulation of CD39+ regulatory T cells confers protection against collagen II-induced arthritis via the production of IL-10. *J Immunol* 2010; **184**: 7144–7153.
- 132 Collison LW, Workman CJ, Kuo TT, Boyd K, Wang Y, Vignali KM *et al*. The inhibitory cytokine IL-35 contributes to regulatory T-cell function. *Nature* 2007; **450**: 566–569.
- 133 Wirtz S, Billmeier U, Mchedlidze T, Blumberg RS, Neurath MF. Interleukin-35 mediates mucosal immune responses that protect against T-cell-dependent colitis. *Gastroenterol* 2011; **141**: 1875–1886.
- 134 Collison LW, Chaturvedi V, Henderson AL, Giacomini PR, Guy C, Bankoti J *et al*. IL-35-mediated induction of a potent regulatory T cell population. *Nat Immunol* 2010; **11**: 1093–1101.
- 135 Collison LW, Pillai MR, Chaturvedi V, Vignali DA. Regulatory T cell suppression is potentiated by target T cells in a cell contact, IL-35- and IL-10-dependent manner. *J Immunol* 2009; **182**: 6121–6128.
- 136 Kulkarni AB, Ward JM, Yaswen L, Mackall CL, Bauer SR, Huh CG *et al*. Transforming growth factor- β 1 null mice: an animal model for inflammatory disorders. *Am J Pathol* 1995; **146**: 264–275.
- 137 Jun S, Ochoa-Repáraz J, Zlotkowska D, Hoyt T, Pascual DW. Bystander-mediated stimulation of proteolipid protein-specific regulatory T (Treg) cells confers protection against experimental autoimmune encephalomyelitis (EAE) via TGF- β . *J Neuroimmunol* 2012; **245**: 39–47.
- 138 Ochoa-Repáraz J, Rynda A, Ascón MA, Yang X, Kochetkova I, Riccardi C *et al*. IL-13 production by regulatory T cells protects against experimental autoimmune encephalomyelitis independently of autoantigen. *J Immunol* 2008; **181**: 954–968.
- 139 Carmona L, Cross M, Williams B, Lassere M, March L. Rheumatoid arthritis. *Best Pract Res Clin Rheumatol* 2010; **24**: 733–745.
- 140 Scott DL, Wolfe F, Huizinga TW. Rheumatoid arthritis. *Lancet* 2012; **376**: 1094–1108.
- 141 Feldmann M, Brennan FM, Maini RN. Role of cytokines in rheumatoid arthritis. *Annu Rev Immunol* 1996; **14**: 397–440.
- 142 Brennan FM, Maini RN, Feldmann M. Role of pro-inflammatory cytokines in rheumatoid arthritis. *Springer Semin Immunopathol* 1998; **20**: 133–147.
- 143 Kannan K, Ortmann RA, Kimpel D. Animal models of rheumatoid arthritis and their relevance to human disease. *Pathophysiology* 2005; **12**: 167–181.
- 144 Di Giovine FS, Nuki G, Duff GW. Tumor necrosis factor in synovial exudates. *Ann Rheum Dis* 1988; **47**: 768–772.
- 145 Saxne T, Palladino MA Jr, Heinegard D, Talal N, Wollheim FA. Detection of tumor necrosis factor alpha but not tumor necrosis factor beta in rheumatoid arthritis synovial fluid and serum. *Arthritis Rheum* 1988; **31**: 1041–1045.
- 146 Firestein GS, Alvaro-Gracia JM, Maki R. Quantitative analysis of cytokine gene expression in rheumatoid arthritis. *J Immunol* 1990; **144**: 3347–3353.
- 147 Piguet PF, Grau GE, Vesin C, Loetscher H, Gentz R, Lesslauer W. Evolution of collagen arthritis in mice is arrested by treatment with anti-tumor necrosis factor (TNF) antibody or a recombinant soluble TNF receptor. *Immunol* 1992; **77**: 510–514.
- 148 Thorbecke GJ, Shah R, Leu CH, Kuruvilla AP, Hardison AM, Palladino MA. Involvement of endogenous tumor necrosis factor α and transforming growth factor β during induction of collagen type II arthritis in mice. *Proc Natl Acad Sci USA* 1992; **89**: 7375–7379.
- 149 Williams RO, Feldmann M, Maini RN. Anti-tumor necrosis factor ameliorates joint disease in murine collagen-induced arthritis. *Proc Natl Acad Sci USA* 1992; **89**: 9784–9788.
- 150 Moreland LW, Baumgartner SW, Schiff MH, Tindall EA, Fleischmann RM, Weaver AL *et al*. Treatment of rheumatoid arthritis with a recombinant human tumor necrosis factor receptor (p75)-Fc fusion protein. *N Engl J Med* 1997; **337**: 141–147.
- 151 Lipsky PE, van der Heijde DM, St. Clair EW, Furst DE, Breedveld FC, Kalden JR *et al*. Infliximab and methotrexate in the treatment of rheumatoid arthritis. Anti-Tumor Necrosis Factor Trial in Rheumatoid Arthritis with Concomitant Therapy Study Group. *N Engl J Med* 2000; **343**: 1594–1602.
- 152 Fava RA, Olsen NJ, Postlethwaite AE, Broadley KN, Davidson JM, Nanney LB *et al*. Transforming growth factor beta 1 (TGF- β 1) induced neutrophil recruitment to synovial tissues: implications for TGF- β -driven synovial inflammation and hyperplasia. *J Exp Med* 1991; **173**: 1121–1132.
- 153 Cassatella MA, Gasperini S, Russo MP. Cytokine expression and release by neutrophils. *Ann NY Acad Sci* 1997; **832**: 233–242.
- 154 Manoury-Schwartz B, Chiochia G, Bessis N, Abeshira-Amar O, Batteux F, Muller S *et al*. High susceptibility to collagen-induced arthritis in mice lacking IFN- γ receptors. *J Immunol* 1997; **158**: 5501–5506.
- 155 Vermeire K, Heremans H, Vandeputte M, Huang S, Billiau A, Matthys P. Accelerated collagen-induced arthritis in IFN- γ receptor-deficient mice. *J Immunol* 1997; **58**: 550713.
- 156 Kelchtermans H, De Klerck B, Mittra T, Van Balen M, Bullens D, Billiau A *et al*. Defective CD4+CD25+ regulatory T cell functioning in collagen-induced arthritis: an important factor in pathogenesis, counter-regulated by endogenous IFN- γ . *Arthritis Res Ther* 2005; **7**: R402–R415.
- 157 Finnegan A, Kaplan CD, Cao Y, Eibel H, Glant TT, Zhang J. Collagen-induced arthritis is exacerbated in IL-10-deficient mice. *Arthritis Res Ther* 2003; **5**: R18–R24.
- 158 Weaver CT, Harrington LE, Mangan PR, Gavrieli M, Murphy KM. Th17: an effector CD4 T cell lineage with regulatory T cell ties. *Immunity* 2006; **24**: 677–688.
- 159 Ley K, Smith E, Stark MA. IL-17A-producing neutrophil-regulatory Tn lymphocytes. *Immunol Res* 2006; **34**: 229–242.
- 160 Nurieva RI, Treuting P, Duong J, Flavell RA, Dong C. Inducible costimulator is essential for collagen-induced arthritis. *J Clin Invest* 2003; **111**: 701–706.
- 161 Chabaud M, Durand JM, Buchs N, Fossiez F, Page G, Frappart L *et al*. Human interleukin-17: a T cell-derived proinflammatory cytokine

- produced by the rheumatoid synovium. *Arthritis Rheum* 1999; **42**: 963–970.
- 162 Ziolkowska M, Koc A, Luszczkiewicz G, Ksiezopolska-Pietrzak K, Klimczak E, Chwalinska-Sadowska H *et al*. High levels of IL-17 in rheumatoid arthritis patients: IL-15 triggers in vitro IL-17 production via cyclosporin A-sensitive mechanism. *J Immunol* 2000; **164**: 2832–2838.
 - 163 Kim HR, Cho ML, Kim KW, Juhn JY, Hwang SY, Yoon CH *et al*. Up-regulation of IL-23p19 expression in rheumatoid arthritis synovial fibroblasts by IL-17 through PI3-kinase-, NF- κ B- and p38 MAPK-dependent signalling pathways. *Rheumatol* 2007; **46**: 57–64.
 - 164 Lubberts E, Koenders MI, OppersWalgreen B, van den Bersselaar L, Coenen-Roo CJJ, Joosten LA *et al*. Treatment with a neutralizing anti-murine interleukin-17 antibody after the onset of collagen-induced arthritis reduces joint inflammation, cartilage destruction, and bone erosion. *Arthritis Rheum* 2004; **50**: 650–659.
 - 165 Courtenay JS, Dallman MJ, Dayan AD, Martin A, Mosedale B. Immunization against heterologous type II collagen induces arthritis in mice. *Nature* 1980; **283**: 666–668.
 - 166 Kochetkova I, Trunkle T, Callis G, Pascual DW. Vaccination without autoantigen protects against collagen II-induced arthritis via immune deviation and regulatory T cells. *J Immunol* 2008; **181**: 2741–2752.
 - 167 Fletcher JM, Loneragan R, Costelloe L, Kinsella K, Moran B, O'Farrelly C *et al*. CD39+ Foxp3+ regulatory T Cells suppress pathogenic Th17 cells and are impaired in multiple sclerosis. *J Immunol* 2009; **183**: 7602–7610.
 - 168 Borsellino G, Kleinewietfeld M, Mitri DD, Sternjak A, Diamantini A, Giometto R *et al*. Expression of ectonucleotidase CD39 by Foxp3+ Treg cells: hydrolysis of extracellular ATP and immune suppression. *Blood* 2007; **110**: 1225–1232.
 - 169 Bynoe MS, Viret C. Foxp3+ CD4+ T cell-mediated immunosuppression involves extracellular nucleotide catabolism. *Trends Immunol* 2008; **29**: 99–102.
 - 170 Kochetkova I, Thornburg T, Callis G, Pascual DW. Segregated regulatory CD39+ CD4+ T cell function: TGF- β -producing Foxp3- and IL-10-producing Foxp3+ cells are interdependent for protection against collagen-induced arthritis. *J Immunol* 2011; **187**: 4654–4666.
 - 171 Lehmann D, Ben-Nun A. Bacterial agents protect against autoimmune disease. I. Mice pre-exposed to *Bordetella pertussis* or *Mycobacterium tuberculosis* are highly refractory to induction of experimental autoimmune encephalomyelitis. *J Autoimmun* 1992; **5**: 675–690.
 - 172 Lee J, Reinke EK, Zozulya AL, Sandor M, Fabry Z. *Mycobacterium bovis* bacille Calmette-Guérin infection in the CNS suppresses experimental autoimmune encephalomyelitis and Th17 responses in an IFN- γ -independent manner. *J Immunol* 2008; **181**: 6201–6212.
 - 173 Marshall E. Drug trials. Violent reaction to monoclonal antibody therapy remains a mystery. *Science* 2006; **311**: 1688–1689.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>