Patterns of Visual Loss in Untreated Sick Cell Retinopathy

B J MORIARTY, R W ACHESON, P I CONDON and G R SERJEANT

Jamaica

Summary

Ophthalmic assessments of 120 patients with homozygous sickle cell (SS) disease and of 222 with sickle cell haemoglobin-C (SC) disease were conducted over a period of ten years. Visual acuity loss (V.A. $\leq 6/18$) attributable to sickle cell retinopathy occurred in 10% of untreated eyes during a mean observation period of 6.9 years. Visual loss was strongly associated with proliferative sickle retinopathy ($p < 0.001$) and most commonly resulted from vitreous haemorrhage, tractional retinal detachment and epiretinal membranes. The incidence of visual loss was 31 per 1000 eye-years observation among eyes with proliferative disease compared to 1.4 per 1000 eye-years observation among eyes with non-proliferative disease.

The natural history and ocular complications of sickle cell retinopathy are extensively described,¹⁻⁴ but little attention has been focussed on the frequency or causes of visual loss associated with this condition. The present study describes visual loss in patients with untreated sickle cell retinopathy and discusses the causes of visual morbidity.

Materials and Methods

Over 3000 patients have attended the sickle cell clinic of the University Hospital of the West Indies or a group of peripheral sickle cell clinics operated by staff of the Medical Research Council Laboratories. This represents approximately one third of all Jamaicans with sickling haemoglobinopathies.

The study group consisted of all those patients with SS or SC disease, aged 15 years and over who underwent ophthalmic screening on at least two occasions separated by 3 months or more during the 10 year period 1st October 1976 to 30th September 1986. Cataracts preventing fundoscopy caused ex-

clusion of 16 eyes. Dense vitreous haemorrhages prevented fundoscopy at the start of the study in 14 eyes but these eyes were included after spontaneous resolution of the haemorrhage. No eyes had received treatment prior to entry into the study. The final study group included 342 patients (120 SS, 222 SC) aged 15-60 years at enrolment. The age and sex distribution of patients is summarised in Table I. The average duration of follow-up of patients was 6.9 years (Fig 1).

Assessment included visual symptoms, corrected visual acuity, slit lamp biomicroscopy and intraocular pressure. Direct and indirect ophthalmoscopy were carried out after mydriasis. Red free and colour photography and fluorescein angiography were performed where appropriate.

'Moderate' visual loss was defined as best corrected visual acuity of 6/18 - 6/60 and 'severe' visual loss as best corrected visual acuity $< 6/60$, on at least two occasions separated by at least 3 months. Those eyes showing 'severe' visual loss after a preceding episode of 'moderate' visual loss (e.g. vitre-

From: The Medical Research Council Laboratories (Jamaica), University of the West Indies, Kingston, Jamaica

Correspondence to: B J Moriarty, MA(Cantab), FRCS, Moorfields Eye Hospital, High Holborn, London WC1

Table I Patients and eyes examined according to sex and age when last seen

AGE	MALES		FEMALES		TOTAL	
	PATIENTS	EYES	PATIENTS	EYES	PATIENTS	EYES
15-29	73	105	82	145	155(45%)	250(45%)
30-49	72	112	93	148	165(48%)	260(47%)
50+	8	16	14	28	22(7%)	44(8%)
TOTAL	153	233	189	321	342	554
	(45%)	(42%)	(55%)	(58%)		

ous haemorrhages, tractional retinal detachment) were counted as 'severe'. If any eye developed proliferative changes after a preceding period of non-proliferative retinopathy, the eye was classified as 'proliferative' and follow-up time was measured from the time of onset of proliferative disease.

A randomised controlled trial of Xenon feeder coagulation of proliferative sickle cell retinopathy (PSR^{5,6} conducted between 1978 and 1980 resulted in the treatment of 53 eyes of 78 patients. A randomised controlled trial of Argon scatter photocoagulation conducted since 1982 has resulted in the treatment on 61 eyes of 67 patients. This report describes visual loss in the control eyes of these trials and in both eyes of all other patients where fundoscopy was possible.

Results

Visual loss (V.A. $\leq$ 6/18) occurred in 72 eyes during the study period.

In 52 (10%) eyes of 49 (15%) patients, this was attributable to sickle cell retinopathy, being associated with proliferative changes in 45 eyes and non-proliferative changes in 7 eyes. Visual loss was 'severe' in 27/52 (52%) of eyes involved. The prevalence of visual loss was 45/134 (34%) among eyes affected by PSR which was significantly greater ($X^2 = 114.2$, $p < 0.001$) than among the 7/420 (2%) eyes with non-proliferative disease. The incidence of visual loss was 31 per 1000 eye-years of observation among eyes affected by PSR compared to 1.4 per 1000 eye-years among eyes with non-proliferative changes ($X^2 = 110.2$, $p < 0.001$) giving a relative risk of 23 (95% confidence interval = 8-66).

The commonest causes of visual loss among eyes affected by PSR were vitreous

Fig. 1. Numbers and duration of patient follow up.

haemorrhage and traction retinal detachment (Table II). Analysis of 16 eyes with vitreous haemorrhages examined within two weeks of onset of symptoms and observed over one year indicated that first symptomatic haemorrhages were associated with transient (< 3 months) and generally moderate visual loss whereas subsequent symptomatic haemorrhages generally caused prolonged (> 3 months) and severe visual loss (Table III). The commonest cause of visual loss among eyes without proliferative change was angiod streaks and disciform degeneration (Table IV).

In 20 eyes, visual loss was not attributable to sickle cell disease but to cataract (8 eyes), glaucoma (5 eyes), uveitis (5 eyes), myopia and trauma (1 eye each).

Discussion

There have been few assessments on the prevalence of visual loss in patients with sickle cell disease. An earlier study³ described severe visual loss in 12% of eyes with PSR over an average follow-up of 4 years but since eyes severely affected by PSR were enrolled in a trial of Xenon arc photocoagulation,⁷ this may have underestimated the prevalence

of visual loss among untreated patients. The present study observed severe visual loss in 17% of PSR affected eyes over a mean follow-up period of 6.9 years.

Earlier studies on non-selected patients

demonstrated a higher prevalence of PSR in the SC patients (32%) compared to SS patients (3%).^{2,3} Screening of asymptomatic patients is biased towards SC patients. This is reflected in the study group described, by the

Table II Causes of 'moderate' and 'severe' visual loss in proliferative sickle retinopathy

CAUSES	(Eyes) N= 134					
	'MODERATE'		'SEVERE'		TOTAL	
	Nos.	(%)	Nos.	(%)	Nos.	(%)
Vitreous haemorrhage	10	(46)	10	(43)	20	(44)
Tractional retinal detachment	5	(23)	9	(39)	14	(31)
Epiretinal membranes	6	(27)	2	(9)	8	(18)
Mixed vitreous haemorrhage + traction detach.	—		2	(9)	2	(4)
Rhegmatogenous retinal detachment	1	(4)	—		1	(2)
TOTAL	22		23		45	

Table III Characteristics of first and subsequent vitreous haemorrhages (V.H.)

	First symptomatic V.H. (N = 8)	Subsequent V.H. (N = 8)
PSR > 60°	8 (100%)	8 (100%)
SC Genotype	8 (100%)	8 (100%)
Visual loss > 3/12	0	7 (88%)
Severe visual loss within 2/52	2 (25%)	8 (100%)
Complications	0	8 (100%)
Persistence of V.H. > 6/12 = 6 cases		
Traction detachment = 2 cases		

Table IV Causes of 'moderate' and 'severe' visual loss in non-proliferative sickle cell retinopathy

CAUSES	(Eyes) (N = 420)			
	'MODERATE'		'SEVERE'	
	Nos.	Nos.	Nos.	(%)
Angioid streaks with disciform degeneration	—	4	4	(57%)
Sickling maculopathy	2	—	2	(29%)
Branch retinal artery occlusion	1	—	1	(14%)
TOTAL	3	4	7	

preponderance of SC patients (222) compared to SS patients (120) – whereas nationally the SS genotype (0.4% live births) is more prevalent than SC genotype (0.2% live births). Thus, although the study group is not representative of patients with sickling haemoglobinopathies *per se*, the prevalence, incidence and causes of visual loss in PSR and non-PSR groups is valid.

Visual loss in sickle cell retinopathy is principally determined by PSR which predisposes to vitreous haemorrhage, vitreous traction, retinal detachment, and epiretinal membrane formation.⁸ Auto-infarction of PSR may reduce the chance of visual loss but the present study indicates that a considerable risk of visual loss remains. Vitreous haemorrhage was the most frequent cause of both 'moderate' and 'severe' visual loss.

Risk factors for vitreous haemorrhage have been identified as the SC genotype, presence of previous vitreous haemorrhage, and PSR involving more than 60° of the retinal circumference.⁶ Iron and blood products from previous haemorrhage may accelerate vitreous degeneration, causing further traction and predisposing to further more persistent, vitreous haemorrhage and/or a tractional detachment.

Tractional retinal detachment (Fig 2) results from elevation of the retina by adherent vitreous bands, and the majority of such

Fig. 2. Tractional retinal detachment. Red-free photograph of left eye of 36 year old male SC patient with tractional vitreous band extending from PRS lesion (at 4 o'clock) to supero-temporal vessels and tractional retinal detachment involving posterior pole. Visual acuity: 6/36.

detachments (85%) occurred in eyes with pre-existing vitreous haemorrhage. Involvement of the posterior pole usually results in severe visual loss difficult to treat by conventional surgery⁹ although occasionally amenable to therapy with the Neodymium:YAG laser. Rhegmatogenous retinal detachment (Fig 3) is uncommon in sickle cell retinopathy but may result from traction on thin atrophic retina.

Epiretinal membranes may be responsible for moderate visual loss in sickle cell disease, and occasionally a localised associated tractional retinal detachment (Fig 4) may result in severe visual loss. Risk factors for ERM

Fig. 3. Rhegmatogenous retinal detachment. Red-free photograph of left eye of 35 year old male SC patient showing a small retinal hole resulting from traction to base of fibrotic 'sea-fan' lesion.

Fig. 4. Epiretinal membrane. Red-free photograph of left eye of 20 year old female SC patient showing an epiretinal membrane with detachment of supero-temporal arcade and a pseudo-macular hole. Visual acuity: 3/60.

Fig. 5. Angioid streaks with disciform degeneration. Fluorescein angiogram of right eye of 59 year old female SS patient demonstrating angioid streaks and disciform macular degeneration. Visual acuity: counting fingers at 1 metre.

Fig. 6. 'Sickling maculopathy'. Fluorescein angiogram of right eye of 47 year old female SS patient depicting an enlarged foveal avascular zone and pathological avascular zones. Visual acuity: 6/18.

formation including the presence and extent of PSR involvement, pre-existing vitreous haemorrhage, and non-treatment of PSR.⁸

Angioid streaks in sickle cell disease are usually of the mild peripapillary type and rarely cause visual loss,¹⁰ although severe visual loss from untreatable disciform macular degeneration (Fig 5) secondary to a presumed subretinal neovascular membrane (SRNVM) occurred in 4 eyes.

Posterior pole vaso-occlusion is a rare cause of prolonged visual loss, although central retinal artery occlusion,^{2,11} branch arteriolar occlusion¹² and macular infarction¹³ have all been reported. Sickling maculopathy results from discrete occlusions and remodelling of the capillary network causing pathological avascular zones and enlargement of the foveal avascular zone¹⁴⁻¹⁶ (Fig 6).

The visual loss in sickle cell retinopathy affects a relatively young group of patients and is largely associated with proliferative disease. Using our incidence figures in conjunction with prevalence rates of PSR in SS and SC disease of 3% and 32% respectively, it can be estimated that 73% of visual loss in sickle cell retinopathy is attributable to proliferative disease (95% confidence interval = 47-89). Avoidance of such visual loss must therefore depend on the early detection and effective treatment of proliferative disease.

We thank Ms G H Maude and Mr D Dunn for statistical help and Mrs S Grant for secretarial assistance.

References

- Goldberg MF: Natural history of untreated proliferative sickle retinopathy. *Arch Ophthalmol* 1971; **85**: 428-37.
- Condon PI and Serjeant GR: Ocular findings in homozygous sickle cell anaemia in Jamaica. *Am J Ophthalmol* 1972; **73**: 533-43.
- Condon PI and Serjeant GR: Ocular findings in hemoglobin SC disease in Jamaica. *Am J Ophthalmol* 1972; **74**: 921-31.
- Condon PI and Serjeant GR: Behaviour of untreated proliferative sickle retinopathy. *Br J Ophthalmol* 1980; **64**: 404-11.
- Jampol LM, Condon PI, Farber MD, *et al.*: A randomized clinical trial of feeder vessel photocoagulation of proliferative sickle cell retinopathy. I. Preliminary results. *Ophthalmol* 1983; **90**: 540-5.
- Condon PI, Jampol LM, Farber MD, *et al.*: A randomized clinical trial of feeder vessel photocoagulation of proliferative sickle cell retinopathy. II. Update and analysis of risk factors. *Ophthalmol* 1984; **91**, 2: 1496-8.
- Condon PI and Serjeant GR: Photocoagulation in proliferative sickle retinopathy: Results of a 5-year study. *Br J Ophthalmol* 1980; **64**: 832-40.
- Moriarty BJ, Acheson RW, Serjeant GR: Epiretinal membranes in sickle cell disease. *Br J Ophthalmol* 1987; **71**: 466-9.

- ⁹ Goldberg MF and Jampol LM: Treatment of neovascularization, vitreous hemorrhage, and retinal detachment in sickle cell retinopathy. In: Symposium on Medical and Surgical Disease of the Retina and Vitreous. St Louis, C V Mosby 1983; 53-81.
- ¹⁰ Hamilton AM, Pope FM, Condon PI, *et al.*: Angioid streaks in Jamaican patients with homozygous sickle cell disease. *Br J Ophthalmol* 1981; **65**: 341-7.
- ¹¹ Acacio I and Goldberg MF: Peripapillary and macular vessel occlusions in sickle cell anemia. *Am J Ophthalmol* 1973; **74**: 861-6.
- ¹² Condon PI, Whitelocke RAF, Bird AC, *et al.*: Recurrent visual loss in homozygous sickle cell disease. *Br J Ophthalmol* 1985; **64**: 700-6.
- ¹³ Knapp JW: Isolated macular infarction in sickle cell (SS) disease. *Am J Ophthalmol* 1972; **73**: 857-9.
- ¹⁴ Stevens TS, Busse B, Chang-Box L, *et al.*: Sickling hemoglobinopathies. Macular and perimacular vascular abnormalities. *Arch Ophthalmol* 1974; **92**: 455-63.
- ¹⁵ Asdourian GK, Nagpal KC, Busse B, *et al.*: Macular and Perimacular vascular remodeling in sickling haemoglobinopathies. *Br J Ophthalmol* 1976; **60**: 431-53.
- ¹⁶ Marsh RJ, Ford SM, Rabb MF, *et al.*: Macular vasculature, visual acuity, and irreversible sickled cell in homozygous sickle cell disease. *Br J Ophthalmol* 1981; **66**: 155-60.